

Micronutrient Biofortification in Pulses: an Agricultural Approach

ABSTRACT

More than two billion of the global populations are malnourished. In India, 25% of the total children are suffering from protein energy malnutrition. India ranks 103rd among the 119 country in Global Hunger Index (GHI). Though world level hunger declined from 29.2 in 2000 to 20.9 in 2018, the figures still holds a deadly future. Micronutrient malnutrition is a very serious problem mostly affecting children and women in the country. The impact is highly seen in poor and landless rural people who can't afford diverse foods or supplements in their diets with needed nutritionnutritients. Among consumed foods, pulses are the cheapest sources of proteins, vitamins and micronutrients and can be supplied to the people through daily diet. Biofortification in pulses through agronomic, breeding and microbial intervention can increase the level of bioavailable micronutrients especially Zn and Fe in the final food products. This paper focuses on the role of micronutrients on human health and various mechanisms to get nutrient rich staple food along with main emphasis on biofortification.

The structure of this abstract is not good.

Please you must show in this paragraph 1) the background which is here, but should be reduced, 2) The objective which must be SMART, 3) the methodologie use to collect or to prepare and analyse the data, 4) The main results, 5) the main conclusion or recommendation.

Keywords: Biofortification, Pulse, Micronutrient, Malnutrition, Hunger

1. INTRODUCTION

Worldwide, more than two billion of people or one in every three persons is spotted to be troubled with multiple micronutrient deficiencies (FAO, 2015). Growing children are grievously affected by nutrient deficiencyiesy compared to adults, as their nutrients requirement changes according to growth and developmental phages (Prieto and Cid, 2011). In Kolhapur district, 40% children between the age group of 8-9 years are micronutrients deficient (iron in 38.8% and fluoride in 36.6% respectively) (*Bharati et al 2018*) and globally were ?? it is 22% (GNR, 2018). ~~At-In the wholea#~~ India, level-18 percent% of infants had a birth weight of less than 2.5kg, 38% children below five years were under-weight, 28% mild, 29%

moderately and 2% severely anaemic (NFHS- 4, 2015-16). Malnutrition caused by vitamins and minerals is also known as “Hidden hunger”, which don’t give any visual symptom usually. As per GHI 2018, India ranked 103rd among 119 countries and while world-wide level of hunger declines from 29.2 in 2000 to 20.9 in 2018. Micronutrient deficiencies are the fountainhead of various health issues like poor neurological function, impaired eye sight, diabetes, hypertension, weak immunity, diarrhea, food allergies, thinning hair, leaky gut, acne or rashes (Lynch and Green 2001; Beard 2001; Shankar and Prasad 1998; Gilbert and Foster 2001; Stein *et al.*, 2005). Those deficiencies are attributable to less-low intake of quality diet, ~~enriched~~ with proteins, vitamins and minerals (Bhatnagar *et al.*, 2011 and Bouis and Saltzman, 2017). Increased price of non staple commodities is one of the important reasons of decreasing dietary quality, especially to resource poor people (Bouis *et al.*, 2011). In developing countries agricultural products are the prime source of ~~nutrition-nutritrients~~ (Graham *et al.*, 2001; Schneeman, 2001). Main concern of green revolution was laid on yield increase not on quality food production. And it scale down soil productivity accompanied by less nutritive food grain production (Bhatnagar *et al.*, 2011). Micronutrient rich vegetables, pulses and animal products have also not been increased in last fifty years (Bouis and Saltzman, 2017). Possible ways to combat those deficiencies encircle dietary diversification (healthy balance diet), food fortification, biofortification and supplementation (Allen *et al.* 2006). Biofortification is the process of increasing nutrient concentration in plant edible parts by fertilization (agronomic intervention), breeding approaches or microbes (White and Broadley, 2005), whereas fortification is nutrient enrichment during processing (https://en.wikipedia.org/wiki/Food_fortification). Biofortification is an effective strategy in long run to overcome the current situation as it is more cost effective, sustainable and practical one to reach poorest of the poor population (Meenakshi *et al.*, 2010, Hoddinott *et al.*, 2013, Garcia-Banuelos *et al.*, 2014). Besides quality enhancement, micronutrient has some added advantages like yield increase, biomass enhancement and disease control in micronutrient deficient soils (Hussain *et al.*, 2010). A healthy balance diet must include pulses as they are rich source of energy, protein, dietary fibre and also content considerable amount of vitamins and minerals like thiamin, riboflavin, pyridoxine, folic acid, vitamin E and K, zinc, iron etc (Ofuya and Akhidue, 2005; Thavarajah *et al.*, 2011; Johnson *et al.*, 2005). So, pulses can be considered as good option for biofortification to provide nutritious food sustainably (Thavarajah *et al.*, 2011).

2. ROLE OF MICRONUTRIENTS ON HUMAN HEALTH

Iron plays key role in haemoglobin formation and oxygen transport (Underwood and Suttle, 1999). Iron deficiency exerts influence on learning ability (CDC 2010), immune system (Fiall, 2003), ability to work (Viteri, 1974) and cognitive development (Bread and Connor, 2003). Its deficiency is also associated with anemia and pregnancy related issues like mortality, low birth weight etc (CDC 2010).

Zinc requirement get larger during pregnancy and puberty. Zinc deficiency curtails physical growth and development of children (Brown *et al.*, 2002). Gastrointestinal, central nervous, epidermal, immune, skeletal, and reproductive systems are known to be affected by zinc deficiency (Hambidge and Walravens, 1982). The daily requirement of Zn and Fe varies with the age of people (Table 1).

Selenium is a good source of antioxidant which narrow down heart and skin diseases, cancer, alzheimer, (Elahi *et al.* 2009; Marksbery and Lovell, 2006; Klaunig and Kamendulis 2004; Cui *et al.*, 2012; Shirley *et al.*, 2014), thyroid (Ventura *et al.*, 2017), asthma (Norton and Hoffmann, 2012). Patients having tuberculosis, influenza and hepatitis C delineated to be benefited by selenium (Steinbrenner *et al.*, 2015).

Table 1. Daily requirements of Zn and Fe in Indian context (ICMR, 2010)

	Group	Recommended Daily Allowance (mg day ⁻¹)	
		Zinc	Iron
Adult men		12	21
Adult women	Normal	10	17
	Pregnant	12	35
Children	1-3 Years	5	9
	4-6 Years	7	13
	7-9 Years	8	16
Adolescents	Boys	11-12	21-28
	Girls	9-12	26-27

3. CRITERIA OF BIOFORTIFIED CROP

Bouis and Welch (2010) suggested the following criteria to be a potential biofortified crop.

- High Yielding: Crop productivity must be maintained.
- Effective: The increased level of micronutrient must have significant positive impact on human.
- Stable: Increased level of micronutrients in crop must be stable year after year.
- Good Taste And Cooking Quality

4. POTENTIAL WAYS OF BIOFORTIFICATION

- Agronomic intervention
- Breeding intervention
- Microbial intervention

4.1 AGRONOMIC INTERVENTIONS

Agronomic biofortification is the application of micronutrients via chemical fertilizer with the aid of foliar application, soil application, seed priming and seed coating of fertilizers to increase the bioavailability of nutrients in edible plant parts (De Valena *et al.*, 2017). Several factors like source of fertilizer, quantity of fertilizer and time and methods of application regulate the nutrient intake to the edible plant parts and its bioavailability to the consumer (Singh and Prasad, 2014, Rietra *et al.*, 2015). Micronutrient amendment in soil is a useful strategy to increase micronutrient quantity in crop (Manzeke *et al.*, 2012; Vanlauwe *et al.*, 2015 and Voortman and Bindraban, 2015). Among the different methods of application, foliar application is more efficient (Lawson *et al.*, 2015) as it can manage soil immobilization (Garcia-Banuelos *et al.*, 2014) and quick availability of nutrients to the crop. Hidoto *et al.* (2017) reported 85 g ha⁻¹ grain zinc yield with foliar application in chickpea which was significantly higher than soil application (71 g ha⁻¹) and priming (68 g ha⁻¹). Combined application in both soil and foliar often showed better results (Phattarakul *et al.*, 2012). Other biofortification methods like seed priming and seed coating are spotted to give very infrequent result (Duffner *et al.*, 2014). Johnson *et al.* (2005) found that seed priming with both B and Zn increased the seed Zn and B content of chickpea and lentil respectively (table 2). Zinc and selenium biofortification is most fruitful with agronomic interventions (Cakmak, 2014).

Table 2. Effect of seed priming on Zn, B and Mo content of chickpea and lentil

Treatments	Seed content (mg kg ⁻¹)					
	Chickpea			Lentil		
	Zn	B	Mo	Zn	B	Mo
(purchased)	40	9	3	50	6	2
water	60	10	4	50	6	2
B	60	100	3	50	100	2
Zn	700	7	3	630	5	2
1/2(B + Zn)**	400	50	2	400	50	2
B + Zn	800	80	3	660	100	2
B, 12 h	40	100	3			
Zn, 12 h	500	8	2			

Mo	60	4	300			
----	----	---	-----	--	--	--

(Source: Johnson *et al.*, 2005) **Priming times were 8 h and 12 h for chickpea and lentil respectively. Solutions used were 0.004M ZnSO₄·7H₂O (for Zn), 0.008 M H₃BO₃(for B), 0.0026M Na₂MoO₄·2H₂O (for Mo).

4.1.1 ZINC FORTIFICATION

Application of zinc to the pulse crops greatly helps in enhancing the level of zinc in harvested (economic) plant parts. Zinc fertilization increases bioavailability of Zn in human by increasing phytate content (Hussain *et al.*, 2013). Guillén-Molina *et al.* (2016) concluded that application of zinc chelate (7 and 14 mM L⁻¹ of Zn-EDTA) increase grain zinc and iron concentration in cowpea. Shivay *et al.* (2015) reported that foliar spray of zinc at three different stages of chickpea had significant influence on zinc uptake both in grain and straw during 2011-12 and 2012-13 (Table 3). Foliar spray of Zn-EDTA at active vegetative, flowering and grain filling stages had greatest crop recovery of applied Zn (17.33%) during 2011-12 (table 2). Zinc fertilization improves zinc bioavailability in bean and pea (Cakmak *et al.* 2010, Zhang *et al.* 2010). Zinc content in seed helps in significant liner increase of protein biosynthesis (Martre *et al.* 2003). Maximum Fe content was recorded with application of 50µM Zn-DTPA (183.7±2.16 ppm) and 100 µM ZnSO₄ (197.9±3.45 ppm) whereas highest Zn with 100µM Zn-DTPA (46.3±3.87 ppm) and 100 µM ZnSO₄ (49.6±2.54 ppm) of bean in hydroponic situation (Table 4). Hidoto *et al.* (2016) stated that maximum grain Zn content and Zn yield in chickpea were noted in soil application of 25 kg ha⁻¹ Zn which had an advantage of 7% over control (table 5).

Table 3. Zinc content by grain and straw of Chickpea

Treatment	Zn uptake in grain (g ha ⁻¹)		Zn uptake in straw (g ha ⁻¹)	
	2011-12	2012-13	2011-12	2012-13
Check (no Zn)	78.5	71.3	78.0	68.5
ZnSHH soil at 5 kg Zn ha ⁻¹	102.3	93.9	104.2	93.9
ZnSHH one spray (V)	96.3	87.9	103.3	92.8
ZnSHH two sprays (V + F)	112.3	103.2	128.6	116.2
ZnSHH, three sprays (V + F + G)	124.9	114.8	166.8	152.0
Zn-EDTA soil at 2.5 kg Zn ha ⁻¹	102.7	93.9	114.5	103.5

Zn-EDTA one spray (V)	98.8	90.9	117.0	106.0
Zn-EDTA two sprays (V + F)	125.4	115.8	139.2	126.6
Zn-EDTA three sprays (V + F + G)	162.8	135.4	181.0	148.9
LSD (P = 0.05)	14.93	15.52	10.45	20.25

ZnSHH= Zn sulfate hepta hydrate V= active vegetative stage, F= flowering stage, G= grain filling stage (Source:

Shivay *et al.*, 2015)

Table 4. Iron and zinc concentration of bean in hydroponic situation

Dose	Micronutrient concentration	
Zn-DTPA (μM)	Fe	Zn
0	146.5 \pm 0.41	28.4 \pm 1.12
25	174.4 \pm 1.45	45.7 \pm 2.35
50	183.7 \pm 2.16	42.8 \pm 3.55
100	153.0 \pm 1.63	46.3 \pm 3.87
ZnSO ₄ (μM)	Fe	Zn
0	146.5 \pm 0.41	28.4 \pm 1.12
25	189.2 \pm 2.89	42.3 \pm 3.11
50	162.1 \pm 2.03	42.6 \pm 2.87
100	197.9 \pm 3.45	49.6 \pm 2.54

Source: (Sida-Arreola *et al.*, 2017)

Table 5. Effect of zinc sulphate soil application on Chickpea

Zn rate	Straw Zn	Grain Zn	Zn yield
ZnSO ₄ ·7H ₂ O (kg ha ⁻¹)	(mg kg ⁻¹)	(mg kg ⁻¹)	(g ha ⁻¹)
0	20.63	37.05	91.0
5	20.48	37.54	98.3
10	23.24	34.20	87.7
15	22.15	33.11	86.2
20	21.82	35.52	86.3
25	21.57	39.55	99.7
30	22.31	39.18	98.0

Source: Hidoto *et al.*, 2016.

4.1.2 IRON FORTIFICATION

Iron is another most important micronutrient which improves human health. Supply of iron through fortification of pulses is helpful and economic for major portion of Indian population. Iron content of cowpea bean seed increased 29.4% with application of 100 μM L⁻¹ ferrous sulphate and 32% with 50 μM L⁻¹

¹ ferrous chelate over control (Mirquez- Quiroz *et al.*, 2015). Ali *et al.* (2014) observed that application of 1.5% FeSO₄ at branching and flowering resulted 55%, 66% and 81% increase in iron content in leaf, stem and grain in mungbean over control respectively (Table 6). Khalid *et al.* (2015) reported that application of PGPR along with iron (@ 5.6 kg ha⁻¹) resulted grain, root and shoot iron content 4.6 mg, 3.16 mg and 1.7 mg in 100 g chickpea seed respectively (Table 7). According to Salih (2013), foliar fertilization of 2 ppm Fe and 2 ppm Zn reported maximum increase in Fe (154 mg kg⁻¹) and Zn (42 mg kg⁻¹) content of cowpea seed respectively (Table 8). Nandan *et al.* (2018) pointed out that foliar spray of Fe @ 0.05% along with recommended dose of fertilizer resulted significantly higher iron content in seed (66.46 mg kg⁻¹) and stover (66.83 mg kg⁻¹) whereas, maximum zinc content in seed (44.98 mg kg⁻¹) and straw (44.08 mg kg⁻¹) was noted with Zn (0.5%) and Fe (0.05%).

Table 6. Iron content in leaves, stems and grains in mungbean

Treatment	Iron content (mg kg ⁻¹)		
	Leaves	Stems	Grains
Control	511.37	380.07	78.50
0.5% FeSO ₄ at branching	601.73	470.42	90.43
0.5% FeSO ₄ at flowering	623.70	488.17	96.10
0.5% FeSO ₄ at branching + 0.5% FeSO ₄ at flowering	675.43	520.24	101.50
1.0% FeSO ₄ at branching	654.07	515.22	96.83
1.0% FeSO ₄ at flowering	668.37	505.16	99.60
1.0% FeSO ₄ at branching + 1.0% FeSO ₄ at flowering	717.17	585.54	127.80
1.5% FeSO ₄ at branching	672.60	550.33	115.73
1.5% FeSO ₄ at flowering	698.70	559.51	121.43
1.5% FeSO ₄ at branching + 1.5% FeSO ₄ at flowering	794.90	634.27	146.43

Source: Ali *et al.*, 2014

Table 7. Iron uptake in different plant parts of chickpea

Treatment	Fe Concentration (mg 100 g ⁻¹)
-----------	--

	Grains	Shoot	Root
Absolute control	1.20	0.66	0.14
Fe @ 5.6 kg ha ⁻¹	2.40	1.80	0.86
S1	3.26	2.23	1.40
S2	3.30	2.50	1.30
S3	3.36	2.26	1.33
S4	3.20	2.36	1.36
S5	3.40	2.40	1.30
S1+Fe @ 5.6 kg ha ⁻¹	3.60	2.73	1.70
S2+Fe @ 5.6 kg ha ⁻¹	4.36	3.16	1.56
S3+Fe @ 5.6 kg ha ⁻¹	3.50	2.80	1.50
S4+Fe @ 5.6 kg ha ⁻¹	3.53	2.70	1.50
S5+Fe @ 5.6 kg ha ⁻¹	3.63	2.63	1.46

Source: Khalid *et al.*, 2015

Table 8: Effect of foliar fertilization on Fe, B and Zn content of cowpea

	Treatment	Fe	B	Zn
		Mg kg ⁻¹		
	Control, 0 pmm	40.00	16.00	8.00
	Fe, 1 ppm	90.00	31.00	25.00
	Fe, 2 ppm	154.00	47.00	42.00
	B, 1 ppm	51.00	31.00	18.00
	B, 2 ppm	58.00	40.00	24.00
	Zn, 1 ppm	47.00	26.00	13.00
	Zn, 2 ppm	50.00	37.00	17.00
Tukey's HSD	Treatment and concentration	1.28	1.35	1.35
	Interaction	2.61	2.94	2.94

Source: Salih, 2013

4.1.3 SELENIUM FORTIFICATION

Selenium fertilization by means of inorganic fertilizer results increased selenium concentration in diet (White and Broadley, 2009; Alfthan *et al.*, 2015). Unlike selenite (SeO_3^{2-}), selenate (SeO_4^{2-}) provides immediate availability to plants when added to soil (Broadley *et al.*, 2006; Fordyce, 2013; Pilbeam *et al.*, 2015). Selenium foliar application increases concentration in pea and common bean from $21 \mu\text{g kg}^{-1}$ to $743 \mu\text{g kg}^{-1}$ (Smrkolj *et al.*, 2005) and 30 to $2379 \mu\text{g kg}^{-1}$ (Smrkolj *et al.*, 2007) respectively.

Further credibility of agronomic biofortification requires much more research on micronutrient bioavailability, including metabolic pathways that affect absorption and health benefits of different chemical forms of micronutrients.

4.2 BREEDING INTERVENTIONS

When utilizable genetic variability is present in a species then genetic biofortification is conductible, but when there is no variability, transgenic approaches are well qualified (Garg *et al.*, 2018). Initially reduction of Phytic acid and polyphenols are used to be the fundamental approach of biofortification as these compounds are known to narrow down iron bioavailability. But recent studies implies that priority should be given to increase iron concentration rather than Phytic acid and Polyphenol reduction because those also have some beneficial properties and resist cancer cell (Pixley *et al.*, 2011, Murgia *et al.*, 2012). Zein protein over expression on soybean increases methionine and cysteine content (Dinkins *et al.*, 2001) and methionine content by cystathionine γ -synthase (Song *et al.*, 2013, Hanafy *et al.*, 2013). Increase in beta carotene and oleic acid in soybean has been attended by introducing bacterial PSY gene (Schmidt *et al.*, 2015) and siRNA-mediated gene silencing had been used to reduce α -linolenic acids (Flores *et al.*, 2008). Similarly, linoleic acid and palmitic acid content of soybean was reduced by antisense RNA technology (Zhang *et al.*, 2014). Storage albumin of Brazil nut which is rich source of methionine has been used to increase common bean methionine content (Aragao *et al.*, 1999) whereas, lupines methionine has been intensified by albumin of Sunflower (Molvig *et al.*, 1997). A sensitive approach to understand the escalated zinc uptake is DNA strand breakage (King *et al.*, 2015).

Field trials regarding genetic effect on selenium concentration reported significant difference among genotypes (Thavarajah *et al.*, 2010; Garrett *et al.*, 2013; Ray *et al.*, 2014). 94 pea genotypes were grown in Saskatchewan field (University of Saskatchewan) and not a single nucleotide polymorphism (SNP) marker was noted to affect seed Se concentration (Diapari *et al.*, 2015). In contrast, lentil and chickpea revealed genotypic variation associated with selenium concentration in Saskatchewan (Thavarajah *et al.*, 2008 ; Thavarajah, 2012; Ray *et al.*, 2014; Rahman *et al.*, 2015). Field experiments conducted in Morocco, Nepal, Syria, Australia and Turkey were also ensured significant genetic variance in lentil Se concentration (Thavarajah *et al.*, 2011). Mungbean (Nair *et al.*, 2015) and soybean (Yang *et al.*, 2003) also shown genetic variation. Bean has a potential to increase zinc content

by 50% and iron by 60-80% as it evidence high heritability in zinc and iron content (Blair *et al.*, 2009; Beebe *et al.*, 2000; Petry *et al.*, 2015).

4.3 MICROBIAL INTERVENTIONS

Phytoavailability of micronutrients can be increased by soil microorganisms like *Rhizobium*, *Bacillus*, *Pseudomonas* etc (Rengel *et al.*, 1999; Smith, 2007). PGPR can be an alternate approach to biofortify pulses as it increases disease resistance (Phi *et al.*, 2010; Dary *et al.*, 2010), solubility of phosphorus (Richardson, 2001; Wani, 2007) and root growth (Glick, 1995, Zhang *et al.*, 2010). But the implication of PGPR and other microorganisms in biofortification of pulses are sparse (De *et al.*, 2011). Rhizobacteria produce siderophores which promote iron fortification in crop as well as revamps soil fertility directly by enhancing iron availability at rhizosphere or indirectly by reducing pathogen effect (Rana *et al.*, 2012; Srivastava *et al.*, 2013).

Grain protein concentration of chickpea ranged from 180 to 309 mg g⁻¹ with inoculation of *Bacillus* PSB1 and *M. ciceri* RC3 + *A. chroococcum* A4 + *Bacillus* PSB10 respectively with 25% yield advantage (Wani, 2007).

Fungi and bacteria improves bioavailability of zinc at rhizosphere zone (Fasim *et al.*, 2002; Biari *et al.*, 2008) due to decline in soil pH (Koide and Kabir, 2000; Subramanian *et al.*, 2000), chelation (Whiting *et al.*, 2001) and increased root sphere (Burkert and Robson, 1994).

Some biofortified pulse crop varieties were released across the world helping to combat the present situation of malnutrition and hidden hunger of mineral nutrients among the people (table 9 and 10).

Table 9. Several Lentil released varieties that possess high iron and zinc levels (The 2nd Global Conference on Biofortification: Getting Nutritious Foods to People, Ashutosh Sarker (ICARDA))

Country	Variety	Content (ppm)	
		Fe	Zn
Bangladesh	Barimusur-4	86.2	---
	Barimusur-5	86	59
	Barimusur-6	86	63
	Barimusur-7	81	---
Nepal	Sisir	98	64
	Khajurah-2	100.7	59
	Khajurah-1	---	58

	Shekhar	83.4	---
India	Pusa Vaibhav	102	---
	L4704	125	74
	IPL 220	73-114	51-64
	Pusa Ageti Masoor	65.0	---
Syria	Idlib-2	73	---
	Idlib-3	72	---
Ethiopia	Alemaya	82	66

Table 10. Iron biofortified bean variety released by Harvest Plus (Garg *et al.*, 2018)

Rwanda	Democratic Republic of Congo
RWR 2245, RWR 2154, MAC 42, MAC 44, CAB 2, RWV 1129, RWV 3006, RWV 3316, RWV 3317, and RWV 2887	COD MLB 001, COD MLB 032, HM 21-7, RWR 2245, PVA 1438, COD MLV 059, VCB 81013, Nain de Kyondo, Cuarentino, Namulenga.

5. CONCLUSION

Largest number of hungry people especially children and women live in India which is quite alarming. In a developing country like India, where maximum people does not have sufficient access to afford commercially fortified food, diversified diet and food suppliments, biofortification is an acceptable cost effective way to eliminate malnutrition. And evidences revealed that a nutritious food like pulse is one of the good options to fortify.

6. REFERENCES

- Alfthan G, Eurola M, Ekholm P. Effects of nationwide addition of selenium to fertilizers on foods, and animal and human health in Finland: from deficiency to optimal selenium status of the population. *Journal of Trace Elements in Medicine and Biology*, 2015;31: 142–147.
- Ali B, Ali A, Tahir M and Ali S. Growth, Seed yield and quality of mungbean as influenced by foliar application of iron sulfate. *Pakistan Journal of Life and Social Sciences*, 2014;12(1): 20-25.
- Allen L, Benoist B, Dary O and Hurrell R. Guidelines on food fortification with micronutrients. World Health Organization and Food and Agriculture Organization of the United Nations. 2006.
- Aragão FJL, Barros LMG, De Sousa MV, Grossi de Sá MF, Almeida ERP, Gander et al. Expression of a methionine-rich storage albumin from the Brazil nut (*Bertholletia excelsa* HBK, Lecythidaceae) in

transgenic bean plants (*Phaseolus vulgaris* L., Fabaceae). *Genetics and Molecular Biology*, 1999;22(3): 445-449.

Beard JL. Iron biology in immune function, muscle metabolism and neuronal functioning. *The Journal of nutrition*, 2001;131(2): 568S-580S.

Beard JL & Connor JR. Iron status and neural functioning. *Annual review of nutrition*, 2003;23(1): 41-58.

Beebe S, Gonzalez AV and Rengifo J. Research on trace minerals in the common bean. *Food and Nutrition Bulletin*, 2000;21(4): 387-391.

Bharati HP, Kavthekar SO, Kavthekar SS and Kurane AB. Prevalence of micronutrient deficiencies clinically in rural school going children. *International Journal of Contemporary Pediatrics*, 2018;5(1):234-238

Bhatnagar M, Bhatnagar-Mathur P, Reddy SD, Anjaiah V and Sharma K K. Crop Biofortification Through Genetic Engineering: Present Status and Future Directions. Institute of Biotechnology, Acharya NG Ranga Agricultural University, Hyderabad 500 030 India. 2011. <file:///H:/Papers/Bio/2011Bhatnagaretal.BiofortRev.392-407.pdf>.

Biari A, Gholami A and Rahmani HA. Growth promotion and enhanced nutrient uptake of maize (*Zea mays* L.) by application of plant growth promoting rhizobacteria in arid region of Iran. *Journal of Biological Science*, 2008;8: 1015–1020.

Blair MW, Astudillo C, Grusak MA, Graham R and Beebe SE. Inheritance of seed iron and zinc concentrations in common bean (*Phaseolus vulgaris* L.). *Molecular Breeding*, 2009;23(2): 197-207.

Bouis HE and Saltzman A. Improving nutrition through biofortification: A review of evidence from Harvest Plus, 2003 through 2016. *Global Food Security*, 2017;12:49–58.

Bouis HE and Welch RM. Biofortification—a sustainable agricultural strategy for reducing micronutrient malnutrition in the global south. *Crop Science*, 2010;50:S-20.

Bouis HE, Eozenou P and Rahman A. Food prices, household income, and resource allocation: socioeconomic perspectives on their effects on dietary quality and nutritional status. *Food and Nutrition Bulletin*, 2011;32 (1): S14–S23.

Broadley MR, White PJ, Bryson RJ, Meacham MC, Bowen HC, Johnson SE et al.. Biofortification of UK food crops with selenium. *Proceedings of the Nutrition Society*, 2006;65(2): 169-181.

Brown KH, Pearson JM, Rivera J & Allen LH. Effect of supplemental zinc on the growth and serum zinc concentrations of prepubertal children: a meta-analysis of randomized controlled trials. *The American journal of clinical nutrition*, 2002;75(6): 1062-1071.

251 Bürkert B, and Robson A. ⁶⁵Zn uptake in subterranean clover (*Trifolium subterraneum* L.) by three
 252 vesicular-arbuscular mycorrhizal fungi in a root-free sandy soil. *Soil Biology and Biochemistry*, 1994;
 253 26(9):1117-1124.

254 Cakmak I. Agronomic biofortification. Conference brief #8, In: *Proceedings of the 2nd Global Conference*
 255 *on Biofortification: Getting Nutritious Foods to People*, Rwanda. 2014.

256 Cakmak I, Pfeiffer WH, McClafferty B. Biofortification of durum wheat with zinc and iron. *Cereal Chemistry*. 2010; 87(1):10-
 257 20.

258 CDC. Breastfeeding Report Card, United states: Outcome Indicators (Publication, from Centers for
 259 Disease Control and Prevention, National Immunization Survey. 2010.

260 Cui H, Kong Y & Zhang H. Oxidative stress, mitochondrial dysfunction, and aging. *Journal of signal*
 261 *transduction*, 2012; pp. 13.

262 Dary M, Chamber-Peerez MA, Palomares AJ, and Pajuelo E. “In situ” phytostabilisation of heavy metal
 263 polluted soils using *Lupinus luteus* inoculated with metal resistant plant-growth promoting rhizobacteria.
 264 *Journal of Hazardous Materials*, 2010; 177(1): 323-330.

265 De Santiago A, Quintero JM, Aviles M and Delgado A. Effect of *Trichoderma asperellum* strain T34 on
 266 iron, copper, manganese, and zinc uptake by wheat grown on a calcareous medium. *Plant and Soil*,
 267 2011;342(1-2): 97-104.

268 De Valença AW, Bake A, Brouwer I D and Giller K E. Agronomic biofortification of crops to fight hidden
 269 hunger in sub-Saharan Africa. *Global food security*, 2017;12: 8-14.

270 Diapari M, Sindhu A, Warkentin TD, Bett K and Tar'an B. Population structure and marker-trait
 271 association studies of iron, zinc and selenium concentrations in seed of field pea (*Pisum sativum*
 272 L.). *Molecular Breeding*, 2015;35(1): 30.

273 Dinkins RD, Reddy MS, Meurer CA, Yan B, Trick H, Thibaud-Nissen F et al.. Increased sulfur amino acids
 274 in soybean plants over expressing the maize 15 kDa zein protein. *In Vitro Cellular & Developmental*
 275 *Biology-Plant*, 2001;37(6): 742-747.

276 Duffner A, Hoffland E, Stomph TJ, Melse-Boonstra A, Bindraban PS. Eliminating zinc deficiency in rice-
 277 based systems. VFRC Report 2014/2. Virtual Fertilizer Research Center, Washington, D.C. 2014.

278 Elahi MM, Kong YX and Matata BM. Oxidative stress as a mediator of cardiovascular disease. *Oxidative*
 279 *medicine and cellular longevity*, 2009;2(5): 259-269.

280 Failla ML. Trace elements and host defense: recent advances and continuing challenges. *The Journal of*
 281 *nutrition*, 2003;133(5): 1443S-1447S.

282 FAO, IFAD, WFP. *The State of Food Insecurity in the World 2015*. FAO, Rome. 2015.

283 Fasim F, Ahmed N, Parsons R and Gadd GM. Solubilization of zinc salts by a bacterium isolated from
 284 the air environment of a tannery. FEMS microbiology letters, 2002;213(1):1-6.

285 Flores T, Karpova O, Su X, Zeng P, Bilyeu K, Sleper DA et al. Silencing of GmFAD3 gene by siRNA leads
 286 to low α -linolenic acids (18: 3) of fad3-mutant phenotype in soybean [*Glycine max* (Merr.)]. Transgenic
 287 research, 2008;17(5):839-850.

288 Fordyce FM. Selenium deficiency and toxicity in the environment. In Essentials of medical geology.
 289 Springer, Dordrecht. 2013;pp. 375-416

290 Garcia-Banuelos ML, Sida-Arreola JP and Sanches E. Biofortification – promising approach to increasing
 291 the content of iron and zinc in staple food crops. Journal of Elementology, 2014;19(3): 865–888.

292 Garg M, Sharma N, Sharma S, Kapoor P, Kumar A, Chunduri V et al. Biofortified crops generated by
 293 breeding, agronomy, and transgenic approaches are improving lives of millions of people around the
 294 world. Frontiers in nutrition, 2018;5: 12. doi.org/10.3389/fnut.2018.00012

295 Garrett RG, Gawalko E, Wang N, Richter A and Warkentin TD. Macro-relationships between regional-
 296 scale field pea (*Pisum sativum*) selenium chemistry and environmental factors in western
 297 Canada. Canadian Journal of Plant Science, 2013;93(6):1059-1071.

298 Gilbert C and Foster A. Childhood blindness in the context of VISION 2020: the right to sight. Bulletin of
 299 the World Health Organization, 2001;79:227-232.

300 Glick BR. The enhancement of plant growth by free-living bacteria. Canadian journal of
 301 microbiology, 1995;41(2): 109-117.

302 Global Hunger Index, 2018.

303 Global Nutrition report, 2018.

304 Graham RD, Welch RM & Bouis H. Addressing micronutrient malnutrition through enhancing the
 305 nutritional quality of staple foods: principles, perspectives and knowledge gaps. Advances in Agronomy,
 306 2001;70:77–142.

307 Hambidge KM and Walravens PA. Disorders of mineral metabolism. Clinics in
 308 gastroenterology, 1982;11(1):87-117.

309 Hanafy MS, Rahman SM, Nakamoto Y, Fujiwara T, Naito S, Wakasa K. et al. Differential response of
 310 methionine metabolism in two grain legumes, soybean and azuki bean, expressing a mutated form of
 311 *Arabidopsis* cystathionine γ -synthase. Journal of plant physiology, 2013;170(3): 338-345.

312 Hidoto L, Worku W, Mohammed H and Taran B. Agronomic Approach to Increase Seed Zinc Content and
 313 Productivity of Chickpea (*Cicer arietinum* L.) Varieties on Zinc Deficient Soils of Southern Ethiopia.
 314 *Advances in Life Science and Technology*, 2016;42.

315 Hidoto L, Worku W, Mohammed H and Taran B. Effects of zinc application strategy on zinc content and
 316 productivity of chickpea grown under zinc deficient soils. *Journal of Soil Science and Plant*
 317 *Nutrition*, 2017;17(1): 112-126.

318 Hidoto L, Tar'an B, Worku W and Mohammed H. Towards zinc biofortification in chickpea: performance of
 319 chickpea cultivars in response to soil zinc application. *Agronomy*, 2017;7(1): 11.

320 Hussain S, Maqsood M A and Rahmatullah M. Increasing grain zinc and yield of wheat for the
 321 developing world: A Review. *Emirates Journal of Food and Agriculture*, 2010;326-339.

322 Hussain S, Maqsood MA, Rengel Z, Aziz T and Abid M. Estimated zinc bioavailability in milling fractions
 323 of biofortified wheat grains and in flours of different extraction rates. *International Journal of Agriculture*
 324 *and Biology*, 2013;15(5): 921–926.

325 (ICMR (2010) <http://icmr.nic.in/final/rda-2010.pdf>)

326 Johnson S E, Lauren JG, Welch RM and Duxbury JM. A comparison of the effects of micronutrient seed
 327 priming and soil fertilization on the mineral nutrition of chickpea (*Cicer arietinum*), lentil (*Lens culinaris*),
 328 rice (*Oryza sativa*) and wheat (*Triticum aestivum*) in Nepal. *Experimental Agriculture*, 2005;41(4): 427-
 329 448.

330 Khalid S, Asghar HN, Akhtar MJ, Aslam A and Zahir ZA. Biofortification of iron in chickpea by plant
 331 growth promoting rhizobacteria. *Pakistan Journal of Botany*, 2015;47(3): 1191-1194.

332 King JC, Brown KH, Gibson RS, Krebs NF, Lowe NM, Siekmann JH et al. Biomarkers of Nutrition for
 333 Development (BOND)—zinc review. *The Journal of nutrition*, 2015;146(4): 858S-885S.

334 Klauni EJ and Kamendulis ML. The role of oxidative stress in carcinogenesis. *Annual Review of*
 335 *Pharmacology and Toxicology*, 2004;44: 239-267.

336 Koide RT, and Z Kabir. Extra radical hyphae of the mycorrhizal fungus *Glomus intraradices* can hydrolyse
 337 organic phosphate. *New Phytologist*, 2000;148: 511–517.

338 Lawson PG, Daum D, Czauderna R, Meuser H and Harling JW. Soil versus foliar iodine fertilization as a
 339 biofortification strategy for field-grown vegetables. *Frontiers in Plant Science*, 2015;6: 450.

340 Manzeke GM, Mapfumo P, Mtambanengwe F, Chikowo R, Tendayi T, Cakmak I. Soil fertility
 341 management effects on maize productivity and grain zinc content in smallholder farming systems of
 342 Zimbabwe. *Plant and Soil*, 2012;361(1-2), 57–69.

343 Markesbery WR and Lovell MA. DNA oxidation in Alzheimer's disease. *Antioxidants and redox*
 344 *signaling*, 2006;8(11-12): 2039-2045.

345 Márquez-Quiroz C, De-la-Cruz-Lázaro E, OsorioOsorio R, Sánchez-Chávez E. Biofortification of cowpea
 346 beans with iron: iron's influence on mineral content and yield. *Journal of Soil Science and Plant Nutrition*,
 347 2015;15(4): 839-847.

348 Martre P, Porter JR, Jamieson PD and Triboï E. Modeling grain nitrogen accumulation and protein
 349 composition to understand the sink/source regulations of nitrogen remobilization for wheat. *Plant*
 350 *physiology*, 2003;133(4): 1959-1967.

351 Meenakshi JV, Johnson NL, Manyong VM, DeGroote H, Javelosa J, Yanggen DR et al.. How cost-
 352 effective is biofortification in combating micronutrient malnutrition? An ex ante assessment. *World*
 353 *Development*, 2010;38(1):64-75.

354 Molina MG, Quiroz CM, de la Cruz Lazaro E, Martinez JRV, Parra JMS, Carrillo MG et al.. Biofortification
 355 of cowpea beans (*Vigna unguiculata* L. Walp) with iron and zinc. *Mexican Journal of Agricultural*
 356 *Sciences*, 2016;17: 3427-3438.

357 Molvig L, Tabe LM, Eggum BO, Moore AE, Craig S, Spencer D et al. Enhanced methionine levels and
 358 increased nutritive value of seeds of transgenic lupins (*Lupinus angustifolius* L.) expressing a sunflower
 359 seed albumin gene. *Proceedings of the National Academy of Sciences*, 1997;94(16): 8393-8398.

360 Murgia I, Arosio P, Tarantino D, & Soave C. Biofortification for combating 'hidden hunger' for iron. *Trends*
 361 *in plant science*, 2012;17(1): 47-55.

362 Nair RM, Thavarajah P, Giri RR, Ledesma D, Yang RY, Hanson P et al. Mineral and phenolic
 363 concentrations of mungbean [*Vigna radiata* (L.) R. Wilczek var. *radiata*] grown in semi-arid tropical
 364 India. *Journal of Food Composition and Analysis*, 2015;39: 23-32.

365 Nandan B, Sharma BC, Chand G, Bazgalia K, Kumar R and Banotra M. Agronomic Fortification of Zn and
 366 Fe in Chickpea an Emerging Tool for Nutritional Security – A Global Perspective. *Acta Scientific*
 367 *Nutritional Health*, 2018;2(4):12-19.

368 National Family Health Survey Report-4, M/o Health & Family Welfare (2015-16).

369 Norton LR and Hoffmann RP. Selenium and asthma. *Molecular Aspects of Medicine*, 2012;33(1): 98–
 370 106.

371 Ofuya ZM, & Akhidue V. The role of pulses in human nutrition: a review. *Journal of Applied Sciences and*
 372 *Environmental Management*, 2005;9(3): 99-104.

373 Petry N, Boy E, Wirth JP and Hurrell RF. Review: The potential of the common bean (*Phaseolus vulgaris*)
 374 as a vehicle for iron biofortification. *Nutrients*, 2015;7: 1144–1173.

375 Phattarakul N, Rerkasem B, Li LJ, Wu LH, Zou CQ, Ram H et al. Biofortification of rice grain with zinc
376 through zinc fertilization in different countries. *Plant and Soil*, 2012;361(1-2): 131–141.

377 Phi QT, Park YM, Seul KJ, Ryu CM, Park SH, Kim JGet al. Assessment of root-associated *Paenibacillus*
378 *polymyxa* groups on growth promotion and induced systemic resistance in pepper. *Journal of*
379 *Microbiology and Biotechnology*, 2010;20(12): 1605-1613.

380 Pilbeam DJ, Greathead HMR and Drihem K. Selenium. In: AV Barker, DJ Pilbeam, eds. *A handbook of*
381 *plant nutrition*, 2nd edn. Boca Raton, FL: CRC Press, 2015;165–198.

382 Pixley KV, Palacios-Rojas N and Glahn RP. The usefulness of iron bioavailability as a target trait for
383 breeding maize (*Zea mays* L.) with enhanced nutritional value. *Field Crops Research*, 2011;123(2): 153-
384 160.

385 Podder R, Tar'an B, Tyler R, Henry C, DellaValle D and Vandenberg A. Iron fortification of lentil (*Lens*
386 *culinaris* Medik.) to address iron deficiency. *Nutrients*, 2017;9(8), p.863.

387 Prieto MB and Cid JLH. Malnutrition in the critically ill child: the importance of enteral
388 nutrition. *International journal of environmental research and public health*, 2011;8(11): 4353-4366.

389 Rahman MM, Erskine W, Materne MA, McMurray LM, Thavarajah P, Thavarajah D et al. Enhancing
390 selenium concentration in lentil (*Lens culinaris* subsp. *culinaris*) through foliar application. *The Journal of*
391 *Agricultural Science*, 2015;153(4): 656-665.

392 Rana A, Joshi M, Prasanna R, Shivay YS, and Nain L. Biofortification of wheat through inoculation of
393 plant growth promoting rhizobacteria and cyanobacteria. *European Journal of Soil Biology*, 2012;50:
394 118-126.

395 Ray H, Bett K, Tar'an B, Vandenberg A, Thavarajah D and Warkentin T. Mineral micronutrient content of
396 cultivars of field pea, chickpea, common bean, and lentil grown in Saskatchewan, Canada. *Crop*
397 *Science*, 2014;54(4): 1698-1708.

398 Rengel Z, Batten GD and Crowley DD. Agronomic approaches for improving the micronutrient density in
399 edible portions of field crops. *Field crops research*, 1999;60(1-2): 27-40.

400 Richardson AE. Prospects for using soil microorganisms to improve the acquisition of phosphorus by
401 plants. *Functional Plant Biology*, 2001;28(9): 897-906.

402 Rietra RPJJ, Heinen M, Dimpka C, Bindraban PS. Effects of nutrient antagonism and synergism on
403 fertilizer use efficiency. VFRC Report 2015/5. Virtual Fertilizer Research Centre, Washington, DC. 2015;p
404 47.

405 Salih HO. Effect of Foliar Fertilization of Fe, B and Zn on nutrient concentration and seed protein of
406 Cowpea "*Vigna unguiculata*". *Journal of Agriculture and Veterinary Science*, 2013;6(3): 42-46.

407 Schmidt MA, Parrott WA, Hildebrand DF, Berg RH, Cooksey A, Pendarvis K et al. Transgenic soya bean
 408 seeds accumulating β -carotene exhibit the collateral enhancements of oleate and protein content
 409 traits. *Plant biotechnology journal*, 2015;13(4): 590-600.

410 Schneeman BO. Linking agricultural production and human nutrition. *Journal of the Science of Food and*
 411 *Agriculture*, 2001;81(1): 3-9.

412 Shankar AH and Prasad AS. Zinc and immune function: the biological basis of altered resistance to
 413 infection. *The American journal of clinical nutrition*, 1998;68(2): 447S-463S.

414 Shirley R, Ord E, and Work L. Oxidative stress and the use of antioxidants in
 415 stroke. *Antioxidants*, 2014;3(3): 472-501.

416 Shivay YS, Prasad R and Pal M. Effects of source and method of zinc application on yield, zinc
 417 biofortification of grain, and Zn uptake and use efficiency in chickpea (*Cicer arietinum*
 418 L.). *Communications in Soil Science and Plant Analysis*, 2015;46(17): 2191-2200.

419 Sida-Arreola JP, Sánchez E, Ojeda-Barrios DL, Ávila-uezada GD, Flores-Córdova MA, Márquez-Quiroz C
 420 et al. Can biofortification of zinc improve the antioxidant capacity and nutritional quality of
 421 beans? *Emirates Journal of Food and Agriculture*, 2017;29(3), p.237.

422 Singh MK and Prasad SK. Agronomic aspects of zinc biofortification in rice (*Oryza sativa*
 423 L.). *Proceedings of the national academy of sciences, India section B: biological Sciences*, 2014;84(3):
 424 613-623.

425 Smith SE, Read DJ. *Mycorrhizal Symbiosis*. 3rd ed. London, UK: Elsevier 2007.

426 Smrkolj P, Germ M, Kreft I, & Stibilj V. Respiratory potential and Se compounds in pea (*Pisum sativum*
 427 L.) plants grown from Se-enriched seeds. *Journal of Experimental Botany*, 2006;57(14): 3595-3600.

428 Smrkolj P, Osvald M, Osvald J, & Stibilj V. Selenium uptake and species distribution in selenium-
 429 enriched bean (*Phaseolus vulgaris* L.) seeds obtained by two different cultivations. *European Food*
 430 *Research and Technology*, 2007;225(2):233-237.

431 Song S, Hou W, Godo I, Wu C, Yu Y, Matityahu I et al.. Soybean seeds expressing feedback-insensitive
 432 cystathionine γ -synthase exhibit a higher content of methionine. *Journal of experimental*
 433 *botany*, 2013;64(7): 1917-1926.

434 Sperotto RA, Ricachenevsky FK, de Abreu Waldow V and Fett JP. Iron biofortification in rice: it's a long
 435 way to the top. *Plant Science*, 2012;190: 24-39.

436 Srivastava MP, Tewari R, and Sharma N. Effect of different cultural variables on siderophores produced
 437 by *Trichoderma* spp. *International Journal of Advance Research*, 2013;1: 1-6.

438 Stein AJ, Meenakshi JV, Qaim M, Nestel P, Sachdev HPS, Bhutta ZA. Technical Monograph
 439 4. Analysing the Health Benefits of Biofortified Staple Crops by Means of the Disability-Adjusted Life
 440 Years Approach: A Handbook Focusing on Iron, Zinc and Vitamin A. Washington, WA: HarvestPlus.
 441 2005.

442 Steinbrenner H, Al-Quraishy S, Dkhil MA, Wunderlich F and Sies H. Dietary selenium in adjuvant
 443 therapy of viral and bacterial infections. *Advances in nutrition*, 2015;6(1): 73-82.

444 Subramanian KS, Tenshia V, Jayalakshmi K, & Ramach V. Role of arbuscular mycorrhizal fungus
 445 (*Glomus intraradices*)(fungus aided) in zinc nutrition of maize. *Journal of Agricultural Biotechnology and*
 446 *Sustainable Development*, 2009;1(1): 029-038.

447 Thavarajah D, Ruszkowski J, Vandenberg A. High potential for selenium biofortification of lentils (*Lens*
 448 *culinaris* L.). *Journal of Agricultural and Food Chemistry*, 2008;57: 10747–10753.

449 Thavarajah D, Warkentin T, Vandenberg A.. Natural enrichment of selenium in Saskatchewan field peas
 450 (*Pisum sativum* L.). *Canadian Journal of Plant Science*, 2010;90: 383–389.

451 Thavarajah P. Evaluation of chickpea (*Cicer arietinum* L.) micronutrient composition: Biofortification
 452 opportunities to combat global micronutrient malnutrition. *Food research international*, 2012;49(1): 99-
 453 104.

454 Thavarajah P, Sarker A, Materne M, Vandemark G, Shrestha R, Idrissi O et al. A global survey of effects
 455 of genotype and environment on selenium concentration in lentils (*Lens culinaris* L.): implications for
 456 nutritional fortification strategies. *Food Chemistry*, 2011;125(1):72-76.

457 Underwood EJ and Suttle NF. 3rd ed. Wallingford: CABI International Publishin. The mineral nutrition of
 458 livestock, 1999;p. 614.

459 Vanlauwe B, Descheemaeker K, Giller KE, Huising J, Merckx R, Nziguheba G et al. Integrated soil fertility
 460 management in sub-Saharan Africa: unravelling local adaptation. *Soil*, 2015;1(1): 491–508.

461 Ventura M, Melo M and Carrilho F. Selenium and Thyroid Disease: From Pathophysiology to Treatment.
 462 *International Journal of Endocrinology*, 2017;9

463 Viteri FE. Anemia and physical work capacity. *Clinics in Haematolgy*, 1974;3:609-626.

464 Voortman R and Bindraban PS. Beyond N and P: Toward a land resource ecology perspective and
 465 impactful fertilizer interventions in sub-Saharan Africa. VFRC Report 2015/1. Virtual Fertilizer Research
 466 Center, Washington, DC. 2015.

467 Wani P, Khan M and Zaidi A. Co-inoculation of nitrogen-fixing and phosphate-solubilizing bacteria to
 468 promote growth, yield and nutrient uptake in chickpea. *Acta Agronomica Hungarica*, 2007;55(3): 315-323.

469 White PJ and Broadley MR. Biofortifying crops with essential mineral elements. Trends in plant
470 science, 2005;10(12): 586-593.

471 White PJ and Broadley MR. Biofortifying crops with essential mineral elements. Trends in plant
472 science, 2005;10(12): 586-593.

473 White PJ and Broadley MR. Biofortification of crops with seven mineral elements often lacking in human
474 diets—iron, zinc, copper, calcium, magnesium, selenium and iodine. New Phytologist, 2009;182(1): 49-84.

475 Whiting SN, de Souza MP, and Terry N. Rhizosphere bacteria mobilize Zn for hyperaccumulation by
476 *Thlaspi caerulescens*. Environmental Science and Technology, 2001;35: 3144–3150.

477 Yang F, Chen L, Hu Q and Pan G. Effect of the application of selenium on selenium content of soybean
478 and its products. Biological Trace Element Research, 2003;93(1-3): 249-256.

479 Zhang L, Yang XD, Zhang YY, Yang J, Qi GX Guo DQ, Xing GJ et al. Changes in oleic acid content of
480 transgenic soybeans by antisense RNA mediated posttranscriptional gene silencing. International journal
481 of genomics, 2014;8.

482 Zhang Y, Shi R, Rezaul KM, Zhang F and Zou C. Iron and zinc concentrations in grain and flour of winter
483 wheat as affected by foliar application. Journal of agricultural and food chemistry, 2010;58(23): 12268-
484 12274.