

Original Research Article

Chemical Composition of *Abrus precatorius* seeds

Abstract

Aims: *Abrus precatorius* (AP) seed powder is used as Ayurvedic tablet (*Sarivadi Vati*) in treating hearing problems. The objective of present work is to describe oil, starch, protein, polyphenol and mineral composition of *Abrus precatorius* seeds.

Methodology: The legume of AP was collected, and seeds were separated manually. The dried seeds in powered form ~~was-were~~ employed for the oil, starch, polyphenol and mineral analysis. ~~not clear, recast~~—The solvent extraction technique was used for elucidation of oil percentage value. The starch content was determined by the enzymatic method. The total polyphenol and flavonoid contents were analyzed spectrophotometrically using Folin-Ciocalteu reagent and aluminum chloride as color developing reagents, respectively. The X-ray fluorescence (XRF) spectrometry technique was used for monitoring of the minerals. at what wave length

Results: The seed kernel was composed of stored oil (3.2%), protein (92.0%) and starch (4.8%). The total polyphenols and flavonoid content were 24710 and 2520 mg/kg (dw). A remarkably high content of polyphenols in the seed coat and seed pot was observed. Mineral nutrients as P, S and (mainly) K were hyper-accumulated by the seed kernel. The seeds showed a glass transition at -21°C , two endothermic peaks at 109°C (dehydration and protein unfolding) and at 209°C , and a calorific value (~ 406 kcal/100

DM) which exceeds the food energy value of *Pisum sativum*, *Lens culinaris* and other common pulses.

Conclusions: The seed kernel was mainly composed of stored protein with lower content of oil and starch. The high content of polyphenol, K, Mg, Ca and Fe in the seeds was stored. The heavy metals were observed under limits for ~~the safe herbal~~ uses of seeds. i suggest abstract should be in single spacing, include a brief recommendation

Keywords: *Abrus precatorius* seed, Thermal properties, ~~Oil,~~ ~~Starch~~, Protein, Polyphenols, Mineral.

1. INTRODUCTION

Abrus precatorius L. is a perennial high-climbing, twining woody toxic vine commonly known as Gunja or Jequirity in what language, include the English Name, family of the climber, which can be abundantly found all throughout the plains of India as a weed. The roots, leaves and seeds of this plant have found medicinal uses ~~(~~ Always use this ~~1,~~ ~~2),~~ [1,2], and it has been reported to have anti-epileptic, anti-viral, anti-malarial, anti-fertility, antidiabetic, neuroprotective, neuromuscular, nephroprotective and immunomodulatory effects, immunostimulatory properties and anti-inflammatory activity ~~-(3, 4)~~ [3,4]. The seeds are considered abortifacient, aphrodisiac, antimicrobial, diuretic and poisonous due to presence of *abrin*, and have been found to be useful in affections of the nervous system and for external use in skin diseases, ulcers and affections of the hair source?. Their antinutritional factors (total free phenols, tannins,

trypsin inhibitor activity and ~~hemagglutinating-meaning~~ activity ~~of what~~ have also been investigated ~~(5,6)~~[5,6]. The seed proteins are rich in most of the essential amino acids, and are deficient only in cysteine and threonine, when compared to the ~~WHO/FAO~~ indicate year, also at first citing use the full names not accronyms requirement pattern.

In this work, nutritional potential and thermal characteristics of *A. precatorius* having scarlet coloured seeds are described. Please add more flesh to this background or include a literature

2. MATERIAL AND METHODS

2.1. Chemicals and Reagents

AR-grade Folin-Ciocalteu reagent obtained from where, aluminum chloride, tannic acid, gallic acid and quercetin ~~—del~~ supplied by Sigma-Aldrich— were used for the analysis of the phenols. AR grade sodium maleate buffer, sodium acetate buffer, potassium hydroxide, amyloglucosidase, pancreatic- α -amylase, glucoseoxidase—peroxidase — ~~del~~ purchased from Megazyme International Ireland Ltd.— ~~del~~ were used for the starch analysis. please provide lists of other equipment/materials, model number and source

2.2. Sample Collection

The *Abrus precatorius* plant was botanically authenticated with the aid of standard monographs (7) did the researcher identify the seed aside monographs? if yes provide the name of the Botanist and the index no.. The plant was ripened in the early summer and collected in May, 2017 from the area located in Pt. Ravishankar Shukla University, Raipur (21.25°N 81.63°E), India. The plants (0.5 ~~del space~~kg) and surface soil (0.5 ~~del~~

spacekg) were collected in separate polyethylene bags. They were transported to the laboratory and sundried for one week in a glass room. The seeds from the legume were manually separated. The *Abrus precatorius* ~~fruit parts and~~not part of your title or objective soil were further dried in an oven at 50 °C overnight. The seed pod, seed coat, kernel and soil samples were crushed into fine power and sieved at 0.1 mm mesh size. what about storage

2.2. Drying of Seeds

The moisture content of the seeds was determined by drying seeds at 105 °C in an air oven for 6 hrs prior to the analysis, and mean values are reported. All characterization results ~~are were~~ presented on a-the basis of dry weight (dw) ~~basisdel~~.

2.3. Measurement of Mass

The mass of seeds was weighed by using the Mettler Toledo electronic balance (AG245). Three seeds were randomly selected for the weighing and their mean mass was reported.

2.4. Thermal Characterization

A DSC 204 F1 Phoenix apparatus (Netzsch, Selb, Germany) was used for the differential scanning calorimetry (DSC) characterization. Data collection was conducted in the 25-300 °C range with constant heating rate of 10 ~~del space~~°C per min. The analytical parameters were determined using the in-built proprietary software name and version.

2.5. Oil Extraction

~~The ground seeds (5.0 g)~~ 5.0g of the powdered seed were agitated in *n*-hexane (25 mL) in a centrifuge for 1 min ~~provide the rpm~~, according to the procedure described by Gónas et al. (8). The combined supernatant liquids were evaporated in a vacuum rotary evaporator at 40 °C until constant weight was obtained. The oil content was expressed in % (w/w) on a dry weight (dw) of the seed basis.

2.6. Analysis of Starch and Protein

The starch in the seed kernel was determined by the enzymatic method (9). The seed kernel was made of stored oil. ~~Protein and starch del~~ The protein ~~and starch~~ contents was computed by subtracting the value of the oil and starch to a value of 100 to get result in the percentage (10).

2.7. Caloric Value

The energy content of the seeds was estimated ~~from del~~ by multiplying the percentages of protein, fat and carbohydrate by the factors ~~as~~ proposed by Meiners et al. (10).

2.8. Polyphenol Analysis

A sample in powder form (100 ~~del space~~mg) was dispersed in 5 ~~del space~~mL of an acetone and water mixture (70:30, v/v), which was sonicated in an ultra-sonic bath for 20 minutes at 20 °C, according to the procedure reported by Bertaud et al. (11). The total phenolic content (TPC) of each extract was determined as tannic acid by using the

Folin-Ciocalteu reagent as tannic acid (12). The flavonoid content was determined by the ~~aluminum~~aluminium chloride method as ~~quereetin~~Quercetin (13).

2.9. Mineral Characteristics

A Bruker (Billerica, MA, USA) III Tracer SD portable spectrometer ~~are all these 1~~ equipment equipped with a 4W rhodium anode and Xflash SDD 2028 channels was used for the X-ray fluorescence (XRF) elemental analysis of the samples. The calibration was carried out by using standard brown and ~~white cowpea (Vigna unguiculata (L.) Walp.) seeds~~ why use this seed. Is it a standard procedure to use this seed.. provide ref and standard soil sample (NCS DC 73382 CRM).

2.10. Statistics

Total polyphenol, flavonoid, resistant starch, soluble starch and oil content analytical variables were analyzed only for the seeds, while the contents of Cl, P, S, K, Rb, Mg, Ca, Sr, Mn, Fe, Cu and Zn elements were determined both soil and seeds. All analyses were ~~was~~ carried out in triplicate.

3. RESULTS AND DISCUSSION

3.1. Physical Characteristics of Seeds

A group of 6 *Abrus precatorius* seeds were enclosed in the oblong, flat and truncate shaped yellowish ~~colored~~ British or US spelling crosscheck seed pod (**Figure 1**). The seeds were scarlet colored with black spot in globose shape, and a 125 ± 3 mg per seed weight ($n = 3$) was obtained for the samples under study. The seed coat was found to be

relatively thick (representing $29 \pm 1\%$ of the seed weight), while the kernel fraction was $71 \pm 2\%$. The average water content of the seeds was found to be $3.2 \pm 0.1\%$.



Figure 1. Image of *Abrus precatorius* L. seed. [source](#)

3.2. Thermal Characteristics

The DSC thermogram of the kernel (**Figure 2**) showed a glass transition at $-21\text{ }^{\circ}\text{C}$ (onset at $-25.6\text{ }^{\circ}\text{C}$) and two endothermic peaks at $109\text{ }^{\circ}\text{C}$ and $209\text{ }^{\circ}\text{C}$. The glass transition probably reflects rotational mobility of side chains within seed glasses (β -transition or rotation of hydroxyl groups on sugars), although a melting of the stored proteins cannot be excluded. The main endotherm at $109\text{ }^{\circ}\text{C}$ corresponds to dehydration and protein denaturation. The second endotherm, with a peak at $209\text{ }^{\circ}\text{C}$, can be attributed to melting of the carbohydrates and other components. The enthalpy of these effects was found to be 174.4 and 36.17 J/g , respectively.- [link your result to previous studies](#)




Figure 2. *Abrus precatorius* L. seed kernel thermogram scanned.

3.3. Caloric Value

The energy content of the seeds, estimated by multiplying the percentages of protein, fat and carbohydrate by the factors proposed by Meiners et al. (10) was ca. 406 kcal/ 100 kg (DW).recast. present result first before linking to previous studies. Present table or fig to show result at the appendix

3.4. Oil, Starch, Protein and Polyphenol Contents

The seed reserve food consists of oil, starch, protein and other constituents, such as trace elements and bioactive compounds. The oil, starch and protein contents of the

seed kernel were found to be 2.2 ± 0.1 , 4.8 ± 0.2 and $93.0 \pm 1.8\%$, respectively. The concentrations of the soluble and resistant fractions of starch were $0.60 \pm 0.06\%$ and $4.2 \pm 0.2\%$. The TPC and flavonoids content (Fla) in the seed kernel were found to be relatively low: $14200 \pm$ and $1900 \pm$ mg/kg, respectively. In contrast, the TPC (Fla) concentration in the seed coat, seed pod and leaf were found to be 24710 ± 290 (2520 ± 51), 3082 ± 62 (3560 ± 66) and 10230 ± 21 (8000 ± 155) mg/kg, respectively. These values were higher than those reported by Jain et al (14) for seeds collected in the Mumbai region Present table or fig to show result at the appendix.

3.5. Mineral Composition

The concentrations of P, S, K, Rb, Mg, Ca, Sr, Mn, Fe, Cu, Zn, Mo and Pb in the kernel of *A. precatorius* seed were found to be 2302, 1841, 11132, 4.0, 1046, 975, 1.0, 25, 213, 13, 48, 1.0 and 1.0 mg/kg, respectively. The values for P, K, S, Mg and Ca in the seed kernel may be regarded as high, while those of other micronutrients (Cu, Mn, Zn and Fe) was moderate, and those of Rb, Sr, Mo and Pb was low. It is worth noting that the concentration of Mg was higher than that of Ca. The concentrations of Ca, Mn, Fe and Zn were higher than those reported for seeds from another region in India Pani et al. (15). Present table or fig to show result at the appendix

3.6. Bioaccumulation Factor

The soil was brown colored with a pH value of 7.7. The mean concentration of Cl, P, S, K, Rb, Mg, Ca, Sr, Mn, Fe, Cu and Zn was found to be 135, 147, 238, 1370, 7.2, 1480, 6330, 49, 1250, 16700, 48 and 24 mg/kg, respectively. The soil was found to be

dominated with elements i.e. K, Mg, Ca, Mn and Fe. Among them, the highest

concentration of Fe was detected. Present table or fig to show result at the appendix

also link your result to previous studies

The bioaccumulation factor (BC), which describes the accumulation and enrichment of an element in the seed kernel with respect to the soil, was found to be for 15.7, 7.7 and 8.1 for P, S and K, respectively, indicating an hyperaccumulation of these three nutrients. The BC values for Rb, Mg, Ca, Sr, Mn, Fe, Cu, Zn, Mo and Pb were 0.6, 0.7, 0.2, 0.02, 0.02, 0.01, 0.3, 0.9, 1 and 0.04, respectively.

4. CONCLUSIONS

The results of the present study ~~reveal~~revealed a low concentration of oil and starch in *A. precatorius* seed kernel ~~(featuring a lipid content similar to that of fenugreek seeds, and a carbohydrate content is similar to fava beans)~~del, recast your conclusion to

capture summary of research findings. The major fraction of the kernel is composed of protein. P, S and K mineral nutrients were found to be strongly bioaccumulated in the seed, and at least a 1.9-fold molar excess of Mg over Ca was detected in the kernel. As regards polyphenol and flavonoid contents, lower concentrations were present in the kernel than in the seed coat and seed pod. The calorific value of *A. precatorius* seeds exceeds the food energy value of other *Fabaceae*. The deleterious effects of antinutritional substances may be minimised by cooking, since they are heat labile.

provide recommendations

CONSENT

189 Not applicable.

190 ETHICS APPROVAL

191 Not applicable.

192 CONFLICT OF INTEREST

193 The authors declare no conflict of interest, financial or otherwise.

194 COMPETING INTERESTS DISCLAIMER:

195
196 Authors have declared that no competing interests exist. The products used for this
197 research are commonly and predominantly use products in our area of research and
198 country. There is absolutely no conflict of interest between the authors and producers
199 of the products because we do not intend to use these products as an avenue for any
200 litigation but for the advancement of knowledge. Also, the research was not funded by
201 the producing company rather it was funded by personal efforts of the authors.
202

203 REFERENCES

- 204 1. Das A, Jain V, Mishra A. A brief review on a traditional herb: *abrus precatorius* (L.).
205 International Journal of Forensic Medicine and Toxicological Sciall journals in
206 italicsences. 2016; 1 (1): 1-10.
- 207 2. Garaniya N, Bapodra A. Ethno botanical and phytopharmacological potential of
208 *Abrus precatorius* (L.), A review. Asian Pacific Journal of Tropical Biomedine.
209 2014; 4 (1): S27–S34. doi: 10.12980/APJTB.4.2014C1069.
- 210 3. Prathyusha P, Subramanium MS, Sivakumar R. Pharmacognostical studies on white
211 and red forms of *Abrus precatorius* Linn. Indian Journal of Natural Products and
212 Resources. 2010; 1 (4): 476–480.

- 213 4. Sivakumar R, Alagesaboopathi C. Studies on cytotoxicity and antitumor screening of
214 red and white forms of *Abrus precatorius* (L.). African Journal of Biotechnology.
215 2008; 7 (22): 3984–3988.
- 216 5. Desai VB, Sirsi M, Shankarappa M, Kasturibai AR. Chemical and pharmacological
217 investigations on the seeds of *Abrus precatorius* Linn. II. Effect of seeds on mitosis
218 and meiosis in grasshopper, *Poecilocera picta* and some ciliates. Indian Journal of
219 Experimental Biology. 1971; 9 (3): 369–371.
- 220 6. Maiti PC, Mukherjea S, Chatterjee A. Chemical examination of seeds of *Abrus*
221 *precatorius*. Journal of the Indian Academy of Forensic Sciences. 1970; 9:64–68.
- 222 7. Khare CP. Indian Medicinal Plants. Springer-Verlag New York, 2007.
- 223 8. Górnaś P, Rudzińska M, Seglińska D. Lipophilic composition of eleven apple seed
224 oils: A promising source of unconventional oil from industry by-products.
225 Industrial Crops and Products. 2014; 60: 86–91. doi:
226 10.1016/j.indcrop.2014.06.003.
- 227 9. AOAC Official Method. Resistant starch in starch and plant materials enzymatic
228 digestion, First action. 2002.02.
- 229 10. Meiners CR, Derise NL, Lau HC, Crews MG, Ritchey SJ, Murphy EW. 1976.
230 Proximate composition and yield of raw and cooked mature dry legumes. Journal
231 of Agriculture and Food Chemistry. 1976; 24(6): 1122-1126.

11. Bertaud F, Tapin-Lingua S, Pizzi A, Navarrete P, Petit-Conil M. Characterisation of industrial barks for their tannin contents for further green-wood based adhesives applications. InTech Fibre, COST FP0901-Hamburg, 2010.
12. Singleton VL, Orthofer R, Lamuela-Raventós RM. Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods in Enzymology*. 1999; 299: 152-178.
13. Chang CC, Yang MH, Wen HM, Chern JC. Estimation of total flavonoid content in propolis by two complementary colorimetric methods. *Journal of Food and Drug Analysis*. 2002; 10 (3):178-182.
14. Jain A, Sinha P, Jain A, Vavilala S. Estimation of flavonoid content, polyphenolic content and antioxidant potential of different parts of *Abrus precatorius* (L.). *International Journal of Pharmacy and Pharmaceutical Sciences*. 2015; 7(8): ISSN-0975-1491.
15. Pani D, Rath SK, Ray DK, Sahoo SL. Proton induced X-ray emission-based analysis of trace element composition of cotyledon derived in vitro callus culture of *Abrus precatorius* (L.), a multimedical wild legume. *Journal of Radioanalytical and Nuclear Chemistry*. 2016; 308 (1): 113-122.

please cross check the pattern of referencing specified by the journal body. Your references should be consistent, provide vol, no and pages for all.