

## **Original Research Article**

### **Identification, Antagonistic Potentials and Plasmid Profiling of Micro-Organisms Associated with Termitarium and Macerated dead termites from Cashew Trees in Ibule-Soro, Akure Nigeria**

#### **ABSTRACT**

This research was carried out to identify microorganisms associated with termitarium on cashew tree barks and macerated dead termites from Ibule-Soro, Akure, Nigeria. Bacterial isolates such as *Bacillus sp*, *Micrococcus sp*, *Corynebacterium sp*, *Streptococcus sp* were identified, while fungi isolates such as *Aspergillus niger*, *Fusarium solani* and *Penicillium notatum* were identified. The result of antimicrobial sensitivity patterns of the isolates showed that all the bacteria were susceptible to at least three of the antibiotics. However, *Micrococcus sp* and *Bacillus sp* were screened to be multiple antibiotic resistant isolates. Plasmid profiling of these multiple antibiotic resistant bacteria isolates were carried out to determine the size of the bacteria plasmids and genetic basis of their antimicrobial resistance. The isolates were cured of their plasmid and subjected to antibiotic treatments again to determine whether their susceptibility to antibiotic is chromosomal or extra-chromosomal. Antagonistic properties of the isolated bacteria and fungi were determined against known bacterial pathogens such as *Staphylococcus aureus*, *Shigella sp*, *Salmonella sp*, and *Escherichia coli*, the result showed that only the fungus *Penicillium notatum* showed positive and mild antagonistic potential against the selected pathogens. Findings from this research showed the potentials of termite nest as reservoirs for beneficial microorganisms with great antagonistic properties.

**Keywords:** Resistance; Antagonistic; Macerated; Plasmid Profiling; Termitarium; Cashew Tree

#### **INTRODUCTION**

Termitarium is the nest of termites composed of partly digested food materials and fecal matter of termites, containing minerals and other organic constituents that provides a suitable environment for the existence of a huge diversity of microorganisms (Longair, 2004). The microbial population of dual origins from both termites and neighbouring soil might result in greater microbial diversity in the termitarium than termite gut or termite-associated soil. However (Fall *et al.*, 2007).

Comment [R1]: g

Comment [R2]: give space

31 Termitarium are associated with cashew trees (*Anacardium occidentale*), with these  
32 termites boring holes through the plant and using it as a safe haven. *Anacardium occidentale*  
33 is a tropical plant that produces the cashew seed and the cashew apple. The cashew nut, often  
34 simply called a cashew, is widely consumed. It is eaten on its own, used in recipes, or  
35 processed into cashew cheese or cashew butter. The shell of the cashew seed yields  
36 derivatives that can be used in many applications including lubricants, waterproofing, paints,  
37 etc. In terms of uses, it is known that every part of the cashew plant is very useful such that  
38 they possess medicinal properties (Hamad and Mubofu, 2015).

39 The bark and the leaf of the tree possess medicinal benefits and have been used as remedy for  
40 both diarrhea and colic. Cashews leaf extract is utilized to reduce blood sugar and blood  
41 pressure levels. Oils extracted from the seeds prove effective in the preparation of  
42 insecticides. The infusion of the bark of the cashew tree has astringent properties and is used  
43 as a mouthwash for treating oral ulcers and as a remedy for sore throat and influenza. Leaves  
44 of the cashew tree, when boiled with water, serve as an anti-pyretic and are used for the  
45 treatment of aches and pains throughout the body (Hamad and Mubofu, 2015).

## 46 47 **Materials and Methods**

### 48 **Collection of samples**

49 Termite feeding tubes (Termitarium) containing live termites and cashew tree barks were  
50 collected from cashew tree into sterile sample collectors. These samples were collected at a farm  
51 settlement in Ibule-soro village, Ondo state, Nigeria. Samples were analyzed within 6hrs of collection.

52

53

54

55 **Preparation of samples for microbial isolation**

56 The method described in Fall *et al.*, (2007) was adopted for sample preparation. The diluent  
57 used for the samples was sterile distilled water. Using a sterile syringe, a 9ml of sterile distilled water  
58 was dispensed into 3 different test tubes under aseptic conditions and a 1g of the termitarium was  
59 poured into the first test tube, homogenized and a 1ml was taken out for a serial dilution procedure till  
60 the 5<sup>th</sup> dilution was obtained. A 1ml of the last dilution factor was seeded on already sterilize media  
61 for fungal and bacterial isolation (Fawole and Oso, 2007).

62 **Bacteria Isolation from termitarium**

63 A 1ml of dilution of choice from already prepared sample was pour plated into the Petri dish. The  
64 plates were swerved gently to allow proper mixture and were allowed to solidify. All the Petri dishes  
65 were stacked conveniently for storage in the incubator and were incubated at 37°C for 24 hours  
66 inverted (Fawole and Oso, 2007).

67 **Bacteria Isolation from cashew tree bark**

68 A 1ml of the prepared sample was pour plated in sterile Petri dish aseptically. The plates were  
69 swerved gently to ensure even mixture and then allowed to gel. The plates were incubated after  
70 solidifying at 37°C for 24 hours inverted (Fawole and Oso, 2007).

71 **Bacteria Isolation from macerated dead termite**

Comment [R3]: Check it

72 A 1ml of the suspension was dispensed in to the Petri dish and the prepared media were poured on it.  
73 After solidification, the plates were incubated at 37°C for 24 hours inverted ().

74 **Fungi isolation from termitarium**

75 A 1ml of dilution of choice from already prepared sample was pour plated into the Petri dish. The  
76 plates were swerved gently to allow proper mixture and were allowed to solidify. All the Petri dishes  
77 were stacked conveniently for storage in the incubator and were incubated at 25-27°C for 72 hours in  
78 an un-inverted position (Cheesebrough, 2006).

79

80

81 **Fungi Isolation from cashew tree bark**

82 A 1ml of the prepared sample was pour plated in sterile Petri dish aseptically. The plates were  
83 swerved gently to ensure even mixture and then allowed to gel. The plates were incubated after  
84 solidifying at 25-27<sup>0</sup>C for 72 hours inverted (Cheesebrough, 2006).

85 **Fungi Isolation from macerated dead termite**

86 A 1ml of the suspension was dispensed in to the Petri dish and the prepared media were poured on it.  
87 After solidification, the plates were incubated at 25-27<sup>0</sup>C for 72 hours inverted (Cheesebrough, 2006).

88 **Identification and characterization of isolated Bacteria and fungi**

89 Standard and conventional methods of cultural, morphological and biochemical characteristics were  
90 employed in the identification of the organisms following the method of Sarah.*et al* (2016).

91

92 **Sub culturing of isolates**

93 **Sub culturing of bacterial isolates**

94 From the nutrient agar plate, sterile inoculating loop previously flamed to red-hot and cooled was  
95 used to pick different isolates from mixed culture plates and then streaked on an already prepared  
96 nutrient agar. This inoculated medium was incubated at 37<sup>0</sup>C for 24 hours in an inverted position  
97 (Cheesbrough, 2006).

98 **Sub culturing of fungal isolates**

99 Sub culture of distinct colonies of fungal growth was done on fresh potato dextrose agar in other to  
100 get pure fungi isolates. Sterile inoculating needle flamed to red hot was used to pick a distinct  
101 mycelium from the previous plate and the placed at the centre of the fresh culture medium. The  
102 medium was incubated at 25-27<sup>0</sup>C for 48-72 hours inverted (Cheesbrough, 2006).

103

104

**Comment [R4]:** No need sub title.write it in a few lines

105    **Preservation of bacterial isolates**

106    A 10ml of already prepared double strength potato dextrose agar was measured into sterile McCartney  
107    bottles. After sterilization, it was allowed to cool to about 45<sup>0</sup>C and left to solidify in a slant position  
108    at an angle of 45<sup>0</sup>. On solidification, the inoculum was introduced into the bottle aseptically and  
109    incubated at 37<sup>0</sup>C for 24 hours. After 24 hours, growth was seen and was stored at 4<sup>0</sup>C in the  
110    refrigerator until further tests (Cheesbrough, 2006).

111    **Preservation of fungal isolates**

112    A 10ml of already prepared double strength potato dextrose agar was measured into sterile McCartney  
113    bottles. After sterilization, it was allowed to cool to about 45<sup>0</sup>C and left to solidify in a slant position  
114    at an angle of 45<sup>0</sup>. After solidification, the inoculum was introduced into the bottle aseptically and  
115    incubated at 25-27<sup>0</sup>C for 72 hours. After 72 hours, growth was seen and was stored at 4<sup>0</sup>C in the  
116    refrigerator until further tests (Cheesbrough, 2006).

117    **Antibiotic sensitivity screening of bacterial isolates**

118    This test was carried out to determine the resistance and susceptibility of the isolated bacteria to  
119    antibiotics. The various antibiotics impregnated in the gram-positive disc used are as follows:  
120    Erythromycin, Amoxicilin, Ofloxacin, Streptomycin, Chloramphenicol, Cefuroxime, Gentamycin,  
121    Pefloxacin, Co-trimoxazole, Ciprofloxacin. The antibiotic susceptibility testing was carried out using  
122    Kirby-Bauer method as described by (Cheesebrough, 2006). A loop full of a bacteria colony was  
123    picked and emulsified in a Bijou bottle containing 3.0ml of normal saline. A cotton swab was dipped  
124    into the suspension and the swab was pressed against the side of the bottle to remove excess fluid. The  
125    inoculated swab was then streaked across the surface of Mueller Hinton agar and allowed to dry for  
126    five minutes after which sterile forceps were used to carefully remove the disc from its pack and  
127    gently pressed onto the agar surface. The plates were incubated at 37<sup>0</sup>C for 24 hours. The zones of  
128    inhibition were measured in millimetres using a ruler.

129    The zones of inhibition were classified into susceptible (16mm and above), intermediate (11mm-  
130    15mm), and resistant (0-10mm) based on the specified standard of zone of inhibition as described by

131 Cheesebrough, 2006. Antibiotic sensitivity screening was also carried out on multiple drug resistant  
132 isolates already cured of their plasmids with broad spectrum antibiotics (CM128PR100).

### 133 **Antagonistic properties of isolates against selected pathogens**

#### 134 **Bacteria against bacteria**

135 This test was carried out on Mueller Hinton agar on Petri dishes using Fokkema method. Fresh culture  
136 (18-hour culture) was used for this test; bacteria isolates previously preserved on nutrient agar slant  
137 were sub cultured on freshly prepared nutrient agar medium and incubated for 18 hours before the  
138 antagonistic test was carried out. Selected bacteria pathogens such as *Staphylococcus aureus*,  
139 *Streptococcus sp* and *Shigella sp* were sourced as clinical samples from the Ondo State General  
140 Hospital, Akure, Nigeria and used against the isolates from the termitarium. It was by streaking the  
141 test organism on one side of the agar plate and the known pathogen on the other side of the agar plate.  
142 The paired cultures were incubated at 37°C for 24-48hrs and observed for zones of inhibition.

#### 143 **Fungi against bacteria**

144 This was carried out on Mueller Hinton agar too. The fungi isolates from a slant were sub cultured on  
145 a fresh Potato Dextrose Agar for 48-72hrs, until the growth is covering the entire plate. Known  
146 pathogen used for the bacteria above was also used. A cork borer was used to cut out that diameter  
147 from the fungal growth into the center of the fresh Mueller Hinton agar and the known bacterial  
148 pathogen was streaked on the side of the fungi about 5mm apart. The paired cultured plates were  
149 incubated at 25°C for duration of 7 days and the zones of inhibition observed

#### 150 **Plasmid Profile Analysis**

151  
152 An 18 hours old broth culture was used for this analysis. The procedure described by CLSI (2008) was  
153 adopted for this analysis.

#### 154 **Plasmid Curing**

155 The plasmid curing was done by exposing the overnight grown culture at 37 °C and 10mg/ml of  
156 Etidium bromide. After plasmid curing, isolates were subjected to antibiotic sensitivity test again  
157 using broad spectrum antibiotics (CM128PR100) (Brown, 2010).

## RESULT

*Corynebacterium sp*, *Bacillus sp*, *Streptococcus sp*, and *Micrococcus sp* were isolated from termitarium in this research, gram staining showed the microorganisms to be gram positive, glucose positive with variation in subsequent biochemical tests result obtained

UNDER PEER REVIEW

189

Table 1.0 Morphological and Biochemical characteristics of bacterial isolates

I	Gram reaction	Sugar Fermentation				COT	CAT	OX	SP.	MOT	VP/MR	N.I.
		Suc.	Lac.	Glu.	Mann							
C	+ve (short rods)	+ve	-ve	+ve	-ve	+ve	+ve	+ve	-ve	-ve	-ve/-ve	3
B	+ve(bacilli rods)	+ve	+ve	+ve	+ve	-ve	-ve	+ve	+ve	+ve	+ve/-ve	5
S	+ve (cocci in chains)	+ve	+ve	+ve	-ve	-ve	-ve	-ve	-ve	-ve	-ve/+ve	4
M	+ve (cocci)	+ve	-ve	+ve	-ve	-ve	+ve	-ve	-ve	-ve	+ve/-ve	3

190

191 Keys; I- Isolate , C- *Corynebacterium sp*, B- *Bacillus sp*, S- *Streptococcus sp*, M- *Micrococcus sp*, Glu- glucose, Lac- lactose, Suc- sucrose, Mann- Mannitol,

192 COT- coagulase, CAT- Catalase, OX- oxidase, SP- spore forming, MOT- motility, VP/MR- voges poskauer/methyl red, N.I- number of Isolate.

193



194 **Fungal Isolates Obtained from Cashew tree termitarium**

195 Three different fungi were isolated from the termitarium, their microscopic and macroscopic  
 196 characteristics vary greatly and were presented in table 2.

197

198 **Table 2.0: identification of Fungal Isolates**

Fungal isolates	Macroscopic characteristics	Microscopic characteristics	Probable Organism
Isolate 1	Colonies are black with a pale yellow reverse side	Hypha is septate.  Simple upright canidiophores that terminates in glucoseSwelling, bearing phialides at the apex orradiating form the entire surface. Conidiaare one-celled and globose.	<i>Aspergillus niger</i>
Isolate 2	White mycelia with areas of whitish yellow	Aerial mycelium.  They appeared as sickle-shaped.  Conidiophores arose singly from the mycelium and branched near the apex tip.	<i>Fusarium solani</i>
Isolate 3	A yellowish reversed side with black colonies	Hyaline or bright coloured mass that appeared one-celled, ovoid in dry basipetal chains.	<i>Penicillium notatum</i>

199

200

201

## 202 **ANTIMICROBIAL SENSITIVITY RESULT**

203 Test results shows that *Corynebacterium sp*, *Streptococcus sp* were sensitive to most of the antibiotics

204 used in this study compared with *Bacillus spp* which was resistance to about seven of the antibiotics.

205 *Micrococcus spp* was totally resistant to the antibiotics hence the need for a plasmid profile analysis

206 using electrophoresis. Table 3 and table 4 shows the antimicrobial characteristics.

207

UNDER PEER REVIEW

208

209

Table 3.0 Zones of inhibition (in mm) of isolates bacteria against antibiotics

I.C	Antibiotic used with zones of inhibition (mm)										N.I
	ERY	CPX	COT	AMX	OFL	STR	CHL	CEF	GEN	PEF	
C	17.33±0.5	12.22±0.7	16.86±0.3	10.02±0.1	16.23±0.5	17.25±0.9	12.33±1.2	18.13±0.8	00.00	13.37±1.4	3
B	13.10±1.6	00.00	8.15±1.2	00.00	00.00	00.00	00.00	16.23±0.9	00.00	11.13±0.6	4
S	14.05±0.7	00.00	18.05±1.4	17.33±0.8	16.75±0.5	17.05±0.7	16.23±0.3	16.03±0.2	00.00	15.45±0.3	5
M	00.00	00.00	00.00	00.00	00.00	00.00	00.00	00.00	00.00	00.00	3

210 Keys; I.C- isolate codes, ERY= Erythromycin, AMX=Amoxicilin, OFL=Ofloxacin, STR=Streptomycin, CHL=Chloramphenicol, CRO= Cefuroxime,  
 211 GEN=Gentamycin, PFX =Pefloxacin, COT = Co-trimoxazole, CPX=Ciprofloxacin, C- *Corynebacterium sp*, B- *Bacillus cereus*, S- *Streptococcus sp*, M-  
 212 *Micrococcus sp*, N.I- number of isolates, 0-10mm- Resistant, 11-16- Intermediate, 16-above- Susceptible. (Cheesebrough, 2006).

213

214

215

216

217

218

219

220

221

Table 4.0 Antibiotic sensitivity patterns of bacteria isolates

I.C	Antibiotics used										N.I
	ERY	CPX	COT	AMX	OFL	STR	CHL	CEF	GEN	PEF	
C	S	I	S	R	S	S	I	S	R	I	3
B	I	R	R	R	R	R	R	S	R	I	4
S	I	R	S	S	S	S	S	S	R	I	5
M	R	R	R	R	R	R	R	R	R	R	3

222 Keys; I.C- isolate codes, ERY= Erythromycin, AMX=Amoxicilin, OFL=Ofloxacin, STR=Streptomycin, CHL=Chloramphenicol, CRO= Cefuroxime,  
 223 GEN=Gentamycin, PFX =Pefloxacin, COT = Co-trimoxazole, CPX=Ciprofloxacin, C- *Corynebacterium sp*, B- *Bacillus cereus*, S- *Streptococcus sp*, M-  
 224 *Micrococcus sp*, N.I- number of isolates, 0-10mm- Resistant, 11-15- Intermediate, 16-above- Susceptible (Cheesebrough, 2006).

225

226

227

228


229

230

231

232

UNDER PEER REVIEW



## Antagonistic result of Fungi

Results indicate that only *Penicillium notatum* had positive antagonistic effect on *Staphylococcus aureus* and mild antagonistic effect on *Shigella sp* and *Salmonella sp*. Table 5 shows the antagonistic pattern.

Table 5.0: Antagonistic patterns of identified fungi against selected pathogens

I.C	Selected pathogens				N.I
	S.A	SH	S	B.S	
A.N	-ve	-ve	-ve	-ve	4
P.N	+ve	I	I	-ve	3
F.S	-ve	-ve	-ve	-ve	3

Keys: I.C- isolate codes, A.N- *Aspergillus niger*, P.N-*Penicillium nonatum*, F.S-*Fusarium solani* S.A- *Staphylococcus aureus*, SH-*Shigella sp*, S-*Salmonella sp*, B.S-*Bacillus subtilis*, N.I- number of isolates, 0-10mm- -ve (no antagonism), 11-16mm- I (mild antagonism), 16-above- +ve (strong antagonism), (Cheesebrough,2006).

252

253

254 **Antagonistic result for bacteria isolates from Cashew Trees**

255 Test results shows that none of the bacterial isolate had antagonistic effect on selected pathogenic test  
 256 organisms. Table 6 shows the antagonistic pattern of identified bacterial against selected pathogen.

257

258

259 Table 6.0: Antagonistic patterns of identified bacteria against selected pathogens

I.C	Selected pathogens				N.I
	S.A	SH	S	E.C	
C	-ve	-ve	-ve	-ve	3
B	-ve	-ve	-ve	-ve	4
S	-ve	-ve	-ve	-ve	5
M	-ve	-ve	-ve	-ve	3

260 Keys: I.C- isolate codes, C- *Corynebacterium sp*, BC- *Bacillus cereus*, S- *Streptococcus sp*, M-  
 261 *Micrococcus sp*, S.A- *Staphylococcus aureus*, SH-*Shigella sp*, S-*Salmonella sp*, E.C- *Escherichia*  
 262 *coli*, N.I- number of isolates, 0-10mm- Negative (no antagonism), 11-16mm- Intermediate (mild  
 263 antagonism), 16-above- Positive (strong antagonism), +ve- positive, -ve- negative (Cheesebrough,  
 264 2006).

265

266

267


268

269

270 **Plasmid profile of bacteria isolates from Cashew Trees**

271 The results obtained revealed the presence of plasmid bands of different molecular weights. The  
272 molecular weights of the plasmids were determined using DNA- Hind III molecular weight  
273 marker (fig- 1). It was observed that *Bacillus sp* and *Micrococcus sp* contains plasmid with an  
274 estimated molecular weight of 1000 bp and 980 bp respectively.

275



276

277

278 Fig 1.0: Electrophorogram of isolated bacteria plasmid DNA

279 KEY; L – Gene ladder, 1 - *Micrococcus sp*, 2- *Bacillus sp*



280

281 **Sensitivity result of bacteria isolates after plasmid curing**

282 Result shows that *Bacillus sp* and *Micrococcus sp* were both sensitive to the generally antibiotics.

283 This makes the initial resistance of this isolates to be plasmid mediated. Thus, resistivity is extra  
284 chromosomal in nature.

285

UNDER PEER REVIEW

286 Table 7:0 Antibiotic sensitivity patterns of bacterial isolates after plasmid curing

I.C	Antibiotic sensitivity patterns after plasmid curing								N.I
	ERY	CXC	OFL	AUG	CAZ	CRX	GEN	CTR	
M	S	I	S	S	S	I	S	I	3
B	S	S	I	S	S	S	S	S	4

287 Keys: I.C- isolate code, ERY: Erythromycin, CXC: Cloxacillin, OFL: Ofloxacin, AUG: Augumentin, CAZ: Ceftrazidine, CRX: Cefuroxime, GEN:  
288 Gentamicin, CTR: Ceftriaxone, B- *Bacillus sp*,M- *Micrococcus sp*, N.I- number of isolates, 0-10mm- Resistant, 11-16- Intermediate, 16-above-  
289 Susceptible (Cheesebrough, 2006). S- Susceptible, I- intermediate, R- resistant.

## 290 DISCUSSION

291 The microbial load obtained in this study shows the importance of termitarium sampled from cashew  
292 trees as suitable habitats for microorganisms. Relevant studies have opined the rich mineral and  
293 nutrient contents of cashew tree gum which is composed of polysaccharides such as glucose,  
294 mannose, galactose and cellulose; this affords termite nests, bark sheaths and termites inhabiting the  
295 tree environments enough growth factors for wide arrays of microorganisms (Nicoletti *et al.*, 2009  
296 and Adeigbe *et al.*, 2015). However, some fungi isolates obtained especially *Fusarium sp* have also  
297 been implicated in causing damping off disease in cashew plant hence, this justifies the presence of  
298 this fungi in the samples analyzed; this also bears similarities to the findings of Adeigbe *et al.*, (2015).

299 The antagonistic test carried out on the fungi isolates against selected pathogen showed mild  
300 antagonism in the fungus *Penicillium notatum* especially against *Salmonella sp*. Since species of  
301 *Penicillium* are ubiquitous as soil and air fungi, their presence in the termitarium indicates a positive  
302 mutualism of these fungi isolates with the termite guts or the termitarium microenvironment  
303 themselves considering the known potentials of *Penicillium notatum* in production of antimicrobials  
304 against pathogenic bacteria (Nicoletti *et al.*, 2009).

305 The bacteria isolates showed varying degrees of resistance to the antibiotics used against them. This  
306 could be as a result of the microorganisms being exposed to several chemicals used by the farmers on  
307 their crops. The termites on cashew trees may have also been exposed to some insecticides and their  
308 active ingredients which are similar analogues to many of the antibiotics used to evaluate their  
309 sensitivity patterns; resulting in possession of resistant (R-factor) plasmids as survival mechanisms  
310 against these antimicrobials (Adeigbe *et al.*, 2015).

311 Bacteria isolates such as *Bacillus sp*, and *Micrococcus sp*, were screened out to be multiple drug  
312 resistant isolates displaying stellar antibiotic resistance against antibiotics used. These isolates were  
313 analyzed via plasmid profiling to determine if they possess resistant gene encoding plasmids in their  
314 cell structures and if their genetic basis of antimicrobial resistance was extra-chromosomal or not.  
315 They were discovered to possess heavy chained resistant factor chromosomes that encode for

antibiotic resistance, after which they were cured of their plasmids and then subsequent exposure to broad spectrum antibiotic treatments again showed they were susceptible to antibiotic treatments, this also agrees with the findings described in Nicoletti *et al.*, (2009).

## CONCLUSION

This study has shown that the termitarium is a microbial habitat that is rich in many nutrients that enables optimum growth of many microbes, revealed the mild antagonistic potentials of isolated microorganisms obtained from test samples against known selected pathogens and shown that the possession of resistant factor plasmids is responsible for the antibiotic resistance patterns of isolated bacteria to antibiotic used.

## Bibliography

1. Adeigbe, O., Olasupo, F., Adewale, B., and Muyiwa, A. (2015) Review on cashew research and production in Nigeria in the last four decades. *Scientific Research and Essays* 10(5): 196-209.
2. Brown, T. (2010) Vectors for Gene Cloning: Plasmids and Bacteriophages. Gene Cloning and DNA Analysis: An Introduction. Academic Press, U.S.A; 64-76.
3. Chessbrough, M. (2006) District laboratory practice in tropical countries. Cambridge University press, United Kingdom, 124-176.
4. Clinical and Laboratory Standards Institute (CLSI). "Performance standards for antimicrobial disk and dilution susceptibility tests for bacteria isolated from animals Approved standard

- Third edition, CLSI document M31-A3, Clinical and Laboratory Standards Institute, 940 West Valley Road, Wayne Pennsylvania, USA, 28 (2008): 1-99.
5. Fall, S., Hamelin, J., Ndiaye, F., Assigbetse, K., Aragno, M., Chotte, J. and Brauman, A., (2007) Differences between bacterial communities in the gut of a soil-feeding termite (*Cubitermes niokoloensis*) and its mounds. *Appl. Environ. Microbiol.*, 73(4):5199–5208.
  6. Fawole, M. and Oso, B., (2007) Laboratory manual on microbiology. Spectrum books limited, Ibadan, Nigeria, 127.
  7. Hamad, F., and Mubofu, E., (2015) "Potential biological applications of bio-based anacardic acids and their derivatives". *Int J Mol Sci.* 16 (4): 85–90.
  8. Longair, R. (2004) Tusked males, male dimorphism and nesting behaviour in a sub-social Afro tropical wasp, *Synagris cornuta* and weapons and dimorphism in the genus (*Hymenophera, vespidea, enumenie*), *Journal of the Kansas Entomological Society*, 77(4), 528 – 557.
  9. Nicoletti, R., Buommino, E., De Filippis, A., Lopez-Gresa, M., Manzo, E., Carella, A., Petrazzuolo, M., and Tufano, M., (2009) "Bioprospecting for antagonistic *Penicillium* strains as a resource of new antitumor compounds". *World Journal of Microbiology*. 24 (2): 185–95.
  10. Sarah K., *et al.* "Descriptions of Medical Fungi". Department of Molecular and Cellular Biology, School of Biological Sciences, University of Adelaide, Australia. 3rd edition (2016).