

Plug Cell Volume, Growing Media Quality and Benzyl Aminopurine (BAP) Spray Effects for Nursery *Impatiens walleriana* Growth

Original Research Article

ABSTRACT

Plant grower's propagators must take two technological critical decisions: the plug-cell size and the growing medium, both of which have been mentioned as abiotic stress sources for bedding pot plants, including for *Impatiens walleriana*. However, only a few recent reports on bedding pot plants have simultaneously included limiting and non-limiting plug cell volumes and growing medium during nursery. The aim of this work was to assess the nursery performance of *I. walleriana* seedlings grown in four plug cell volumes and four growing media with significant differences in both physical and chemical properties, sprayed or not with an early and single BAP dose, aiming to understand how they interact on determining biomass accumulation at the pot transplant stage. The above sentence is not clear. It must be rewritten. The hypothesis tested was that the both plug cell volume and growing medium, must be seen as additive abiotic stress sources, which can be partially overridden/override by exogenous cytokinin supply. The main result was that, in *I. walleriana* seedlings, the abiotic stress imposed by the growing medium quality during nursery has a higher effect on biomass accumulation (on a fresh and dry base), leaf area expanded and photo assimilates partitioning than plug cell volume and constitute an interactive process associated with cytokinins synthesis. From a grower's point of view, one expensive option to avoid root restriction is to use high quality growing media and increase the plug cell volume. In contrast, a single 100 mg L⁻¹ BAP spray can partially override the root restriction symptoms related to both abiotic stresses. The novelty of this work is related to the fact that growing media quality would be a more limited factor than plug cell volume for *I. walleriana* seedlings during nursery.

Keywords: abiotic stress, bedding plants, cytokinins, plug trays.

1. INTRODUCTION

Technological advances in transplanting have contributed to the growth of the bedding plant industry. The plug system for seedling propagation has been adopted by bedding plant growers to minimize labor costs, increase crop uniformity, and reduce cropping duration. Many growers specialized in the young plant stage (plug) know that the time required for seedlings to reach transplantable size determines greenhouse utilization efficiency and greatly affects production costs. A fundamental objective of most commercial bedding plants growers, including propagators, is to produce a crop that meets the quality standards of the market in the shortest time.

An ornamental crop undergoes different abiotic stresses during commercial production cycle, which decrease both plant growth and quality. Previous reports from our laboratory have shown that plug cell volume [1,2] and growing media quality [3,4,5,6,7] would be considered as abiotic

stresses. Negative growth effects on nursery remain and increased during pot growth phase [3,8,9]. To override the root restriction syndrome related to plug cell volume and growing media quality [9] an early benzyl amino purine spray (BAP) has been suggested [10,11,12,13]. However, quantitative information on the combined effects of these main abiotic stresses during nursery is lacking.

Plant roots can sense adverse soil conditions and, via some internal signal, transmit the condition of the soil to extending leaves, a process named "root restriction syndrome". The typically net result is a decrease in biomass accumulation rates. The primary function of a container is to provide a discrete space for the growth medium, which also affects the physical conditions of the medium. Balancing the air and water content growing media is one of the biggest problems facing plug growers. The second corner of plug air and water content are container size. Plug cells have two basic problems: they are too short and too small.

Soilless substrates are used in horticulture for growing seedlings, plant propagation, and ornamental plant production due to its high physical and chemical stability and low degradation rate, but peat mixes are not necessarily the optimal growth substrates because of their water-logging tendency and low oxygen availability under watering. Alternative soilless growing media can be better than peat for several ornamental bedding plants [14,15,6,7], including *I. walleriana* [16,4,17,9].

It has been claimed that the close coordination between root and shoot growth is controlled by a hormonal signaling pathway located in the root system. Cytokinins are root-synthesized, which are transported via the xylem to the shoot where they exert a major regulatory influence on growth [18,19,20]. Increased root growth may lead to a corresponding increase in the synthesis of cytokinins [21,22,23]. On the other hand, close relationships between root dry weight and physiological processes determining plant growth has been found for ornamentals [6,7,9,24] and vegetables [13] when exogenous cytokinins was applied.

Impatiens walleriana (Hook.f.), also known as busy Lizzie (United Kingdom), balsam, sultan or simply impatiens, is a commercially important year-round garden crop for landscape, and the first best-selling bedding plants in both developed and undeveloped countries. In a previous report [9,24] showed the significant influence of plug cell volume and growing media quality on pot *I. walleriana* growth, but there is no report, until today, on the combined effects of them during nursery.

The aim of this work was to assess the nursery performance of *I. walleriana* seedlings grown in four plug cell volumes and four growing media with significant differences in both physical and chemical properties. Plants were sprayed or not with an early and single BAP dose, aiming to understand how they interact on determining biomass accumulation at the pot transplant stage. The hypothesis tested was that the both plug cell volume and growing medium, must be seen as additive abiotic stress sources, which can be partially override by exogenous cytokinin supply.

2. MATERIALS AND METHODS

2.1. Plant material

Experiment were carried out in a greenhouse at the Faculty of Agronomy, University of Buenos Aires, Argentina (34° 35' 59"S, 58° 22' 23"W) from October 10th 2012 to December 9th 2013 and repeated from October 16th 2013 to December 15th 2014.

I. walleriana 'Xtreme White' seeds (Goldsmith Inc., NY, USA) were germinated and grown in 50-, 128-, 288- and 512-cell plug tray⁻¹ (55.70, 17.37, 6.18 and 2.50 cm³ cell⁻¹ respectively) in four different growing media as follows:

- 1) Klasmann 411[®] medium (Klasmann-Deilmann, GmbH, Germany): Canadian *Sphagnum* peat moss-perlite-vermiculite (70/20/10 v/v/v) (K)

- 2) *Sphagnum maguellanicum*-perlite (80/20 v/v) (S)
- 3) River waste-perlite (80-20 v/v) (R)
- 4) *Sphagnum maguellanicum*-river waste-perlite (40-40-20, v/v/v) (SR).

Leaves were sprayed at sunset with 0 (control plants) or 100 mg L⁻¹ BAP solutions when the first true leaf pair was developed. The BAP concentration was chosen from previous experiments [25].

Comment [N1]: How did authors decide to use this BAB dose (single dose)?

2.2. Cultivation and meteorological data

Formatted: Highlight

Plants were irrigated as needed with high quality tap water using intermittent overhead mist. Growing media were weekly fertilized with 1.0: 0.5: 1.0: 0.5 (v/v/v/v) N: P: K: Ca through the overhead irrigation water (Stage 2: 50 mg L⁻¹ N; Stage 3-4: 100 mg L⁻¹ N; pot: 150 mg L⁻¹ N).

Daily mean temperatures (22.26 to 25.06°C) and daily photosynthetic active radiation (4.24 to 5.03 mol photons m⁻² day⁻¹) for the two experiments were recorded with a HOBO sensor (H08-004-02) (Onset Computer Corporation, MA, USA) connected to a HOBO H8 data logger. The plants were arranged at a density of 25 plants m⁻², which avoided mutual shading.

2.3. Sampling and growth evaluations

Samples of each substrate were collected at the beginning of the experiments (before transplant) and total porosity, air-filled porosity, bulk density and container capacity were determined according to Fonteno [26]. Data are indicated in Table 1 and show significant physical properties differences in of the growing media tested.

Table 1: Physical properties for the growing media tested. K: [Canadian *Sphagnum* peat (70%) + Perlite (20%) + Vermiculite (10%)], S: [*Sphagnum maguellanicum* (80%) + Perlite (20%)], R: [River waste (80%) + Perlite (20%)], SR: [*Sphagnum maguellanicum* (40%) + River waste (40%) + perlite (20%)].

Growing media	Total porosity (%)	Air-filled porosity (%)	Bulk density (g dm ⁻³)	Container capacity (%)
F	60.00 ± 0.55	12.93 ± 0.98	0.21 ± 0.04	36.89 ± 1.46
S	70.67 ± 0.67	29.67 ± 2.15	0.15 ± 0.01	48.00 ± 0.38
R	72.67 ± 0.18	44.60 ± 0.95	0.18 ± 0.01	50.22 ± 0.44
SR	67.53 ± 0.64	23.27 ± 2.43	0.21 ± 0.01	42.67 ± 0.38

Plants were harvested at the transplant stage. Roots were washed and root, stem and leaf fresh weights (FW) were recorded. Dry weights (DW) were obtained after drying roots, stems and leaves to constant weight at 80°C for 96 h. The number of leaves was recorded, and each leaf area was determined using the ImageJ® (Image Processing and Analysis in Java) software.

The relative rate of leaf area expansion (RLAE) was calculated as the slope of the regression of the natural logarithm (ln) of total leaf area versus time (in days). The rate of leaf appearance (RLA) was calculated as the slope of the number of fully expanded leaves versus time (in weeks). The relative growth rate (RGR) was calculated as the slope of the regression of the ln of whole plant DW versus time (in days).

The mean net assimilation rate (NAR) and leaf area ratio (LAR) were calculated as follows:

$$NAR = \frac{k_w W_0 e^{k_w t}}{A_0 e^{k_a t}}$$

$$LAR = \frac{k_w}{NAR}$$

where W_0 : extrapolated value of total DW (g) at time zero; k_w : RGR (day^{-1}); A_0 : extrapolated value of leaf area (cm^2) at time zero; k_a : RLAE (day^{-1}); t : time (days) at the midpoint of the experimental period and e : base of the ln.

The allometric coefficients between root and shoot and between leaf blades and the stem fraction were calculated as the slope (β) of the straight-line regression of the ln of the root DW versus the ln of the shoot DW and between the ln of the leaf blade DW and the ln of the stem DW respectively.


2.4. Statistical analysis

The experimental design was a randomised factorial with three blocks of five single-pot replications of each treatment combination (plug cell volume \times growing medium \times BAP concentration). Since there were no significant differences between the two experiments, they were considered together ($n = 30$). Data were subjected to three-way analysis of variance (ANOVA). STATISTICA 8 (StatSoft) software was used and the assumptions of the ANOVA were checked. Means were separated by Tukey's tests ($P \leq 0.05$). Slopes from straight-line regressions of RLA, RLAE, RGR and allometric values were tested using the SMATR package [27].

3. Results

3.1. Fresh and Dry Weight Accumulation

Although the lower plug cell volume the lower FW at the end of the nursery for *I. walleriana* plants, growing media quality showed a significant effect. The lower FW values were found in plants grown at K-growing media for most treatments. A single BAP spray when the first true leaf pair was developed significantly increased FW in some of the plug cell volume tested (Fig. 1A). When the mean aerial FW was plotted against the mean root FW (Fig. 1B), a positive correlation was found ($r^2 = 0.910$; $P < .001$).






Figure 1. Total FW ($n = 30$) at the end of the experiments (near 60 days from sowing) for *Impatiens walleriana* seedlings grown in four plug cell volumes (50-, 128-, 288-, and 512-cell tray⁻¹) and four growing media, sprayed with zero (control plants) or 100 mg L⁻¹ BAP. Bars indicate standard errors and vertical line indicate least significant differences (LSD). Panel B. Relationships between shoot and root FW according to four plug cell volumes (50-, 128-, 288-, and 512-cell tray⁻¹) and sprayed with zero or 100 mg L⁻¹ BAP at the pre-transplant stage. For substrate abbreviations see Table 1. F: ♦; R: ●; S: ▲; SR: ■. The straight-line regression was: Shoot FW = 5.29 Root FW - 0.006 ($r^2 = 0.910$). The probability of the slope being zero was $P < 0.001$.




Figure 2. Dry weight for roots, stems and leaves ($n = 30$) at the end of the experiments (near 60 days from seeding) for *Impatiens walleriana* seedlings grown in four plug cell volumes (50-, 128-, 288-, and 512-cell tray⁻¹) and four growing media, sprayed with zero (control plants) or 100 mg L⁻¹ BAP. Bars indicate standard errors and vertical line indicate least significant differences (LSD). For substrate abbreviations see Table 1. Panel A showed control plants results while panel B showed BAP-sprayed ones.

When biomass accumulation was plotted on a DW base of different plant organs, lower values were found in control plants (Fig. 2A). The higher DW was found in plants in **S**-, **R**- and **SR**-growing media grown in 50-plug cell tray⁻¹. An early BAP spray increased DW differences in each plant organ (Figure 2B). Leaves showed the higher DW biomass accumulation in all treatments tested and organs.

3.2. Leaf Growth and Plant Growth Analysis

Total leaf areas at the transplant stage (60 days from the beginning of the experiment) showed similar results than FW although a BAP spray in 50-cells tray⁻¹ failed to increase leaf areas (Fig. 3).




Figure 3. Total leaf area ($n = 30$) at the end of the experiments (near 60 days from seeding) for *Impatiens walleriana* seedlings grown in four plug cell volumes (50-, 128-, 288-, and 512-cell tray⁻¹) and four growing media, sprayed with zero (control plants) or 100 mg L⁻¹ BAP. Bars indicate standard errors and vertical line indicate least significant differences (LSD). For substrate abbreviations see Table 1.

R- and **SR**-growing media showed, during nursery, the higher RLA values as control plants as BAP-sprayed ones. The higher plug cell volume the higher RLA in most growing media tested. RLAE differences were quite lower in **K**-growing media but showed the same response than RLA in the rest of growing media tested. An early BAP spray increased both RLA and RLAE. Plants grown in 50-plug cell tray⁻¹ showed the higher RGR values in all growing media tested. BAP-sprayed plants increased RGR only in **S**-, **R**- and **SR**-growing media. Changes in RGR were related to higher NAR values and lower LAR ones (Table 2).

Table 2. Changes in RLA, RLAE, RGR, NAR and LAR for *Impatiens walleriana* seedlings grown in four plug cell volumes (50-, 128-, 288-, and 512-cell tray⁻¹) and four growing media, sprayed with zero (control plants) or 100 mg L⁻¹ BAP at the pre-transplant stage. For substrate abbreviations see Table 1. Different lower-case letters indicate significant differences (P < 0.05) between control and BAP-sprayed plants, while different capital letters indicate significant differences (P < 0.05) between different substrates for each plant cell size. The probability of the slope being zero for RLA, RLAE and RGR was P < 0.001.

	RLA (leaves week ⁻¹ plant ⁻¹)		RLAE (cm ² cm ⁻² day ⁻¹)		RGR (g g ⁻¹ day ⁻¹)		NAR (g cm ⁻² day ⁻¹) (x 10 ⁻⁵)		LAR (cm ² g ⁻¹)	
	Control	BAP	Control	BAP	Control	BAP	Control	BAP	Control	BAP
K										
50	0.087 ^{aD}	0.100 ^{aD}	0.046 ^{aB}	0.063 ^{aB}	0.0632 ^{aC}	0.0627 ^{aD}	10.57 ^{bA}	17.57 ^{aA}	56.22 ^{aB}	43.07 ^{bB}
128	0.087 ^{aC}	0.113 ^{aC}	0.040 ^{aB}	0.048 ^{aB}	0.0489 ^{aC}	0.0468 ^{aD}	8.39 ^{bA}	14.60 ^{aA}	70.52 ^{aC}	46.80 ^{bB}
288	0.080 ^{aA}	0.093 ^{aB}	0.034 ^{aB}	0.043 ^{aB}	0.0469 ^{aC}	0.0403 ^{bD}	6.12 ^{bA}	12.30 ^{aA}	95.88 ^{aB}	7.25 ^{bC}
512	0.067 ^{aB}	0.083 ^{aB}	0.030 ^{aA}	0.032 ^{aB}	0.0428 ^{aB}	0.0406 ^{aC}	3.02 ^{bB}	7.20 ^{aA}	172.91 ^{aA}	85.53 ^{bB}
S										
50	0.139 ^{aC}	0.139 ^{aC}	0.079 ^{aA}	0.080 ^{aA}	0.0724 ^{aB}	0.0751 ^{aB}	8.45 ^{aB}	6.39 ^{bD}	69.83 ^{bA}	92.50 ^{aA}
128	0.067 ^{aC}	0.080 ^{aD}	0.042 ^{aB}	0.057 ^{aB}	0.0394 ^{bD}	0.0563 ^{aC}	8.09 ^{aA}	7.46 ^{aB}	64.39 ^{aC}	73.59 ^{aA}
288	0.063 ^{aB}	0.070 ^{aC}	0.031 ^{aB}	0.049 ^{aB}	0.0363 ^{bD}	0.0516 ^{aC}	6.66 ^{aA}	5.94 ^{aB}	70.87 ^{bC}	103.76 ^{aA}
512	0.033 ^{bC}	0.077 ^{aC}	0.019 ^{bB}	0.044 ^{aA}	0.0290 ^{bC}	0.0508 ^{aB}	5.70 ^{aA}	5.81 ^{aB}	78.45 ^{aB}	80.92 ^{aB}
R										
50	0.210 ^{bA}	0.278 ^{aB}	0.081 ^{aA}	0.090 ^{aA}	0.0781 ^{bA}	0.0852 ^{aA}	9.89 ^{bA}	15.06 ^{aB}	51.66 ^{aB}	34.46 ^{bB}
128	0.133 ^{bA}	0.153 ^{aB}	0.065 ^{aA}	0.076 ^{aA}	0.0618 ^{bB}	0.0714 ^{aB}	2.53 ^{bC}	8.45 ^{aB}	212.60 ^{aA}	62.85 ^{bA}
288	0.094 ^{bA}	0.197 ^{aA}	0.056 ^{aA}	0.071 ^{aA}	0.0607 ^{bA}	0.0706 ^{aA}	3.45 ^{bB}	6.02 ^{aB}	134.08 ^{aA}	84.94 ^{bB}
512	0.087 ^{bA}	0.117 ^{aA}	0.044 ^{aA}	0.056 ^{aA}	0.0545 ^{bA}	0.0600 ^{aA}	1.10 ^{bC}	7.02 ^{aA}	42.79 ^{bC}	89.95 ^{aB}
SR										
50	0.183 ^{bB}	0.390 ^{aA}	0.072 ^{bA}	0.092 ^{aA}	0.0764 ^{aA}	0.0679 ^{aC}	6.92 ^{bC}	11.96 ^{aC}	75.43 ^{aA}	49.07 ^{bB}
128	0.117 ^{bB}	0.200 ^{aA}	0.068 ^{aA}	0.077 ^{aA}	0.0684 ^{bA}	0.0839 ^{aA}	5.40 ^{bB}	13.17 ^{aA}	97.46 ^{aB}	42.14 ^{bB}
288	0.087 ^{aA}	0.096 ^{aB}	0.058 ^{aA}	0.060 ^{aA}	0.0588 ^{bB}	0.0651 ^{aB}	5.31 ^{bB}	7.63 ^{aB}	89.47 ^{aB}	67.20 ^{bB}
512	0.070 ^{aB}	0.083 ^{aB}	0.037 ^{bA}	0.053 ^{aA}	0.0428 ^{bB}	0.0577 ^{aA}	3.00 ^{aB}	2.52 ^{aC}	175.18 ^{aA}	206.44 ^{aA}

3.3 Photo Assimilates Partitioning


Allometries from roots: shoots and from leaves: stems showed that the lower plug cell volume the higher β coefficients (photo assimilates would be partitioned preferably to roots and leaves respectively) for all growing media tested. A BAP spray increased photo assimilates partitioning to shoots and stems respectively (lowered β coefficients) (Table 3).

Table 3. Changes in allometric relationships between roots and shoots and between leaves and stems for *Impatiens walleriana* seedlings grown in four plug cell volumes (50-, 128-, 288-, and 512-cell tray⁻¹) and four growing media, sprayed with zero (control plants) or 100 mg L⁻¹ BAP at the pre-transplant stage. For substrate abbreviations see Table 1. Different lower case letters indicate significant differences (P < 0.05) between control and BAP-sprayed plants, while different capital letters indicate significant differences (P < 0.05) between different substrates for each plant cell size. The probability of the slope being zero for RLA, RLAE and RGR was P < 0.001.

	Roots: Shoots		Leaves: Stems	
	β		β	
	Control	BAP	Control	BAP
K				
50	0.889 ^{aD}	0.834 ^{bB}	0.941 ^{aC}	0.803 ^{bC}
128	0.911 ^{aC}	0.906 ^{aA}	0.995 ^{aB}	0.828 ^{bB}
288	0.987 ^{aA}	0.812 ^{bC}	1.104 ^{aB}	0.823 ^{bB}
512	0.965 ^{aB}	0.803 ^{bC}	1.121 ^{aA}	0.921 ^{bA}
S				
50	1.157 ^{aA}	0.966 ^{bA}	1.006 ^{aD}	1.022 ^{aA}
128	0.900 ^{aB}	0.865 ^{bB}	1.024 ^{aC}	0.822 ^{bC}
288	0.841 ^{aC}	0.739 ^{bC}	1.068 ^{aB}	0.782 ^{bD}
512	0.773 ^{aD}	0.682 ^{bD}	1.117 ^{aA}	0.974 ^{bB}
R				
50	0.905 ^{aA}	0.908 ^{aA}	1.058 ^{aC}	0.676 ^{bC}
128	0.871 ^{aB}	0.790 ^{bB}	1.125 ^{aB}	0.789 ^{bA}
288	0.817 ^{aC}	0.795 ^{bB}	1.151 ^{aA}	0.773 ^{bB}
512	0.923 ^{aA}	0.626 ^{bC}	1.162 ^{aA}	0.786 ^{bA}
SR				
50	0.961 ^{aB}	0.826 ^{bC}	1.115 ^{aD}	0.852 ^{bA}
128	0.981 ^{aB}	0.957 ^{bA}	1.157 ^{aC}	0.632 ^{bC}
288	0.853 ^{aC}	0.852 ^{aB}	1.297 ^{aB}	0.667 ^{bB}
512	1.002 ^{aA}	0.729 ^{bD}	1.386 ^{aA}	0.855 ^{bA}

3.4 Relationships between Growth Parameters and Root Dry Weight

Positive relationships between RLAE ($r^2 = 0.653$ $P < 0.001$) (Fig. 4A), RLA ($r^2 = 0.775$ $P < 0.001$) (Fig. 4B), RGR ($r^2 = 0.637$ $P < 0.001$) (Fig. 4C), NAR ($r^2 = 0.730$ $P < 0.001$) (Fig. 4D) and root DW were found. The highest values were found from plants sprayed with BAP, with a significant influence of the growing medium used.






Fig. 4. Relationship between RLAE (A), RLA (B), RGR (C), NAR (D) and root dry weight (RDW) ($n = 30$) in plants of *Impatiens walleriana* grown in four plug cell volumes (50-, 128-, 288-, and 512-cell tray⁻¹) and four growing media sprayed with zero (control plants, empty symbols) or 100 mg L⁻¹ BAP (full symbols) at the pre-transplant stage. The straight-line regressions were RLAE = 0.02 RDW + 0.045 ($r^2 = 0.653$ P < 0.001), RLA = 0.539 RDW + 0.52 ($r^2 = 0.775$ P < 0.001), RGR = 0.02 RDW + 0.042 ($r^2 = 0.637$ P < 0.001), NAR = 11.94 RDW + 0.95 ($r^2 = 0.730$ P < 0.001). F: \blacklozenge - \circ ; R: \bullet - \circ ; S: \blacktriangle - Δ ; SR: \blacksquare - \square .

4. Discussion

For bedding ornamental plants, quality at the transplant stage is related to a squat, compact seedling and profusely branching with large growing points. A well-developed rooted system with white roots and without damage is needed as well. A decrease in plug cell volume increases plant density and lowered labor costs, however, it gives a vertical root restriction, which leads a pronounced decrease in both root and shoot growth of the early vegetative stages [3]. A non-settle question until today is the impossibility to choose a growing media without regarding for the container size but at same time it is not possible to choose a plug cell size without regarding the growing media used. Decisions such as the choice of container cell size and growing media quality before transplanting are frequently made on the basis of economics [8]. When seedlings are grown in typical transplant containers, growth tends to be proportional to the volume of the container cell. The more the space available to the plant from the nursery, the larger it becomes after transplant and the more quickly it attains particular growth stages. Both a fresh (Fig. 1) and dry (Fig. 2) weight analysis are in agreement with this previous statements. However, in contrast with the previous reports, which indicated the plug cell size as the main abiotic stress during nursery [1,2], growing media quality would be a more restrictive abiotic stress at that growth stage that previously supposed.

Abiotic stress is defined as environmental conditions that reduce growth and yield below optimum levels [28]. Abiotic stress responses in plants occur to various organ levels among which the root specific processes are of particular importance [29]. The growth dynamics of nursery plants is critical because seedlings complete their life cycle in a short time and normally do not have enough time to adjust to unfavorable environmental conditions. On the other hand, Pagani et al. [17] have previously suggested the growing medium as an abiotic stress source would allow changing the media-based paradigm to optimize both the growth and productivity of bedding pot plants. Our previous and present results are in agreement with this previous statements.

A general statement showed that plants increased biomass production through both shoots appearance and leaves expansion; photo assimilates from leaves are the force for the roots, shoots and leaves initiation and growth. The size of the different plant sinks determines the partition of photo assimilates between the plant organs.

A meta-analysis on the effects of pot size showed that root growth responds directly to impedance and decrease root growth and leaf area [30,31,32]. On the other hand, Puig et al. [33] and Chen et al. [34] have concluded that plants can sense the volume of the rooting space available and respond accordingly, having in mind that roots are a major source of cytokinins in plants, which are synthesized in roots and transported to shoots [35]. In this way, the significantly higher shoot FW related to plug cell volume of *I. walleriana* seedlings from Figs. 1 and 3 would not be an unexpected result, although shoot FW increase in plug-grown vegetables exogenously BAP-sprayed has been previously indicated [35,11,12,37]. However, the most significant effect of a BAP spray on both FW and leaf area by growing media than plug cell volume is clearly a novelty result.

To characterize leaf area development, two growth parameters can be analyzed: first, the rate of leaf appearance (RLA) as an estimator of plastochron length, which involve leaf initiation rates. Second, the rate of leaf area expansion (RLAE), which quantify leaf growth. Our results showed that the higher plug cell volume the higher RLA and RLAE. It has been showed that plastochron may be altered in transgenic plants with reduced cytokinin levels [38], which explain the effect of a single BAP spray to override root restrictions. In the same way, Shani et al. [39] showed the effect of cytokinin on leaf growth rates. Both a decrease in the plastochron length and an increase in leaf expansion are accompanied by the SAM size increase through the synthesis of high-molecular weight substances essential for cell growth. Plant tissues and organs rich in cytokinins are known to attract the assimilate translocation and increased the sink capacity of the benzyl adenine-treated leaves [40]. Our results showed that limited both plug cell volume and growing media quality decreased RLA more than RLAE (Table 2), which would conditioned post-transplant growth rates during the pot-grown stages.

Variation in RGR has the result of two key traits: the 'physiological component' NAR and the 'morphological component' LAR [41]. Although leaf area determined plant capacity of light interception, RGR, which ultimate quantify biomass accumulation, is greatly influenced by photosynthetic efficiency [42]. Although the higher plug cell volume the higher RGR and NAR, growing media quality and exogenous BAP increase both growth parameters (Table 2). Shipley [43] indicated that, in general, NAR was the best general predictor of variation in RGR, in agreement with our results from Table 3. On the other hand, Shi et al. [30] showed that root restriction often depresses photosynthetic capacity but indicated that the mechanism for this reduction remains unclear. However, cytokinin is known to stimulate the expression of photosynthetic enzymes like Rubisco [44,45]. Plants synthesize different cytokinin-ribosides but not all have biological activity [46], although the higher root system, the higher the zeatin ribosides [21,22,23]. In any case, we can show positive relationships between RLAE (Fig. 4A), RLA (Fig. 4B), RGR (Fig. 4C) and NAR (Fig. 4D) and root DW.

Poorter and Sack [47] indicated that sink organs can potentially stimulate sugar supply by activating their consumption rate, thereby increasing their sink strength. The relative carbon allocation to a particular organ must be regarded as a function of source and sink activities of all parts of the plant regulate by the relative photo-assimilates allocation [48]. Benincasa et al. [49] indicated that crop management and environmental factors modify the source-sink relationships, including abiotic and biotic stresses. Changes in the allometric slopes (β) reflect variations in carbon partitioning in response to biotic and abiotic environment interactions [50]. Carbohydrate partitioning between competing sites is explained by the fact that plants are capable of modifying their resource allocation to favor the growth of their growing parts [51,52]. Differences in plant allometries shown in Table 3 are in agreement with these assumptions and indicated that a root restriction by plug cell volume or low growing media quality partitioned photo assimilates to roots and leaves while non limited plug cell volume and high growing media quality or exogenous BAP applications changes photo assimilates partitioning to shoots and stems respectively. The lower root systems would not be a limited factor due to the constant water supply under commercial facilities. In agreement with Zwack and Rashotte [53], who indicate that multiple factors influence how cytokinin treatment affects stress signaling and that the spatial, temporal, and developmental context may be important factors in the downstream stress response. The response to an exogenous BAP supply from our results was quantitatively different if plug cell volume or growing media quality abiotic stress sources was analyzed.

5. CONCLUSIONS

The only way to be economically competitive in the future during nursery is to decrease labor costs and increase business efficiency. However our results indicated that it does not mean necessarily to use exclusively more cells per tray and a decrease in growing media quality for bedding pot plant propagation in plug trays. Nursery optimization needs for higher cell tray volume and high growing media quality relationships as much as to identify the different ornamental bedding plant environmental requirements, which is the subject for future research.

REFERENCES

1. Di Benedetto A, Klasman R. The effect of plug cell volume on the post-transplant growth for *Impatiens walleriana* pot plant. Eur. J. Hort. Sci. 2004;69(2): 82-86.
2. Di Benedetto A, Klasman R. The effect of plug cell volume, paclobutrazol height control and the transplant stage on the post-transplant growth for *Impatiens wallerana* pot plant. Eur. J. Hort. Sci. 2007;72(5): 193-197.
3. Di Benedetto A. 2011. Root restriction and post-transplant effects for bedding pot plants. In: Ornamental Plants: Types, Cultivation and Nutrition. (J.C. Aquino, ed.), Nova Science Publishers, Inc. NY, USA. 47-79.
4. Thibaud J, Mc. Loughlin T, Pagani A, Lavado R; Di Benedetto A. Alternative substrates and fertilization routine relationships for bedding pot plants: *Impatiens wallerana*. Eur. J. Hort. Sci. 2012;77(4): 182-191.
5. Pagani A, Molinari J, Lavado RS, Di Benedetto A. Behavior of *Impatiens wallerana* Hook. F in alternative pot substrates: mechanisms involved and research perspectives. J. Plant Nutr. 2015;38: 2185-2203.
6. Gandolfo E, Hakim E, Geraci J, Feuring V, Giardina E, Di Benedetto A. Responses of pansy (*Viola wittrockiana* Gams.) to the quality of the growing media. Amer. J. Exp. Agric. 2016;12(3): 1-10.
7. Hakim G, Gandolfo E, Giardina E, Di Benedetto A. The effect of the pre-transplant pot media quality on pansy garden performance. Int. J. Plant Soil Sci. 2017;19(4): 1-12.
8. Di Benedetto A, Pagani A. Difficulties and possibilities of alternative substrates for ornamental bedding plants: An ecophysiological approach. In: Draguhn C, Ciarimboli N, editors. Peat: Formation, Uses and Biological Effects, Nova Science Publishers, Inc. NY, USA; 2012.
9. DeLojo, J., Gandolfo, E., Gómez, D., Feuring, V., Monti, S., Giardina, E., Boschi, C., and Di Benedetto, A. Root restriction effects on the bedding pot plant *Impatiens walleriana*. J. Exp. Agric. Int. 2017;15(4):1-16.
10. Gandolfo E, De Lojo E, Gómez E, Pagani A, Molinari J, Di Benedetto A. Anatomical changes involved in the response of *Impatiens wallerana* to different pre-transplant plug cell volumes and BAP sprays. Eur. J. Hort. Sci. 2014;79(4): 226-232.
11. Di Matteo J, Rattin J, Di Benedetto A. Increase of spinach growth through the use of larger plug cell volume and an exogenous BAP spray. Amer. J. Exp. Agric. 2015;6(6):372-383.
12. Rattin J, Pico Estrada O, Giardina E, Di Benedetto A. Nursery pre- and post-transplant effects on tomato (*Solanum lycopersicum* L.) growth and yield. J. Exp. Agric. Int. 2017;18(5):1-14.
13. Geraci J, Di Matteo J, Feuring V, Giardina E, Di Benedetto A. Exogenous BAP spray on biomass accumulation in spinach. J. Exp. Agric. Int. 2018;25(6): 1- 17.
14. Di Benedetto A, Klasman R, Boschi C. Use of river waste in growing media for growing ornamental herbaceous perennials. J. Hort. Sci. Biotechnol. 2004;79(1):119-124.
15. Marchese N, Di Benedetto A, Lavado R. The possibilities of river waste and Argentinean peat as a plug growing media for *Verbena x hybrida*. Int. J. Agric. Res. 2006;1(2): 142-150.
16. Di Benedetto A, Petracchi JC, Marcella G, Montaron P, Chavez P. Evaluation of alternative substrates for bedding plants. Int. J. Agric. Res. 2006;1(6): 545-554.
17. Pagani A, Molinari J, Lavado RS, Di Benedetto A. Behavior of *Impatiens wallerana*

- Hook. F in alternative pot substrates: mechanisms involved and research perspectives. *Journal of Plant Nutr.* 2015; 38: 2185-2203
18. Ghanem ME, Albacete A, Smigocki AC, Frebort I, Pospilova H, Martinez-Andujar C, Acosta M, Sanchez-Bravo J, Lutts S, Dodd IC, Perez-Alfocea F. Root synthesized cytokinins improve shoot growth and fruit yield in salinized tomato (*Solanum lycopersicum* L.) plants. *J. Exp. Bot.* 2011;6(1): 125-140.
 19. Hwang I, Sheen J, Müller B. Cytokinin signaling networks. *Annual Rev. Plant Biol.* 2012; 63, 353-80.
 20. Brenner WG, Schmölling T. Summarizing and exploring data of a decade of cytokinin-related transcriptomics. *Front. Plant Sci.* 2015;6, 29.
 21. O'Hare TJ, Turnbull CGN. Root growth, cytokinin and shoot dormancy in lychee (*Litchi chinensis* Sonn.). *Scientia Hort.* 2004;102(2):257-266.
 22. Aloni R, Langhans M, Aloni E, Dreieicher E, Ullrich CI. Root-synthesized cytokinin in *Arabidopsis* is distributed in the shoot by the transpiration stream. *J. Exp. Bot.* 2005; 56(416): 1535-1544.
 23. Kotov AA, Kotova LM. Role of acropetal water transport in regulation of cytokinin levels in stems of pea seedlings. *Russian J. Plant Physiol.* 2015;62(3):390-400.
 24. Williams T, Sala A, Feuring V, Giardina E, Di Benedetto A. Relationships between plug cell size and quality substrate in the bedding pot plant *Impatiens wallerana* (Hook.f.). *Int. J. Plant Soil Sci.* 2016;12(2): 1-12.
 25. Di Benedetto A, Pagani A. Dry weight accumulation in the *Impatiens walleriana* pot plant in responses to different pretransplant plug cell volume. *Europ. J. Hort. Sci.* 2013;78(2):76-85.
 26. Fonteno WC. Growing Media Types and Physical/Chemical Properties. In: *Water, Media and Nutrition for Greenhouse Crops. A Grower's Guide.* (Reed, D.W., Ed.), Ball Publishing, Batavia, Illinois, U.S.A.; 1996.
 27. Warton DI, Duursma RA, Falster DS, Taskinen S. SMATR 3-an R package for estimation and inference about allometric lines. *Methods Ecol. Evol.* 2012;3(2):257-259.
 28. Cramer GR, Urano K, Delrot S, Pezzotti M, Shinozaki K. Effects of abiotic stress on plants: A systems biology perspective. *BMC Plant Biol.* 2011;11(1):163-176.
 29. Ghosh D, Xu J. Abiotic stress responses in plant roots: a proteomics perspective. *Front. Plant Sci.* 2014; 5: 6.
 30. Shi K, Ding XT, Dong DK, Zhou YH, Yu JQ. Root restriction-induced limitation to photosynthesis in tomato (*Lycopersicon esculentum* Mill.) leaves. *Scientia Hort.* 2008;117(3):197-202.
 31. Shi K, Fu LJ, Dong DK, Zhou YH, Yu JQ. Decreased energy synthesis is partially compensated by a switch to sucrose synthase pathway of sucrose degradation in restricted root of tomato plants. *Plant Physiol. Biochem.* 2008;46(12):1040-1044.
 32. Poorter H, Bühler J, van Dusschoten D, Climent J, Postma JA. Pot size matters: a meta-analysis of the effects of rooting volume on plant growth. *Funct. Plant Biol.* 2012;39(11):839-850.
 33. Puig J, Pauluzzi G, Guiderdoni E, Gantet P. Regulation of shoot and root development through mutual signaling. *Mol. Plant.* 2012;5(5):974-983.
 34. Chen BJ, During HJ, Vermeulen PJ, Kroon H, Poorter H, Anten NP. Corrections for rooting volume and plant size reveal negative effects of neighbour presence on root allocation in pea. *Functional Ecol.* 2015;29(11):1383-1391.
 35. Coro M, Araki A, Rattin J, Miravé P, Di Benedetto A. Lettuce and celery responses to both BAP and PBZ related to the plug cell volume. *Amer. J. Exp. Agric.* 2014;4(10):1103-1119.
 36. Yruela I. Plant development regulation: Overview and perspectives. *J. Plant Physiol.* 2015;182(5): 62-78.
 37. Rattin J, Wagner P, Ferreyro D, Riverti D, Giardina E, Di Benedetto A. Roots partially drive super sweet maize yield. *J Exp. Agric. Int.* 2017;16(6):1-17.
 38. Lee BH, Johnston R, Yang Y, Gallavotti A, Kojima M, Travencolo BAN, Costa LF, Sakakibara H, Jackson D. Studies of aberrant phyllotaxy1 mutants of maize indicate complex interactions between auxin and cytokinin signaling in the shoot apical meristem. *Plant Physiol.* 2009;150(1): 205-216.
 39. Shani E, Ben-Gera H, Shleizer-Burko S, Burko Y, Weiss D, Ori N. Cytokinin regulates compound leaf development in tomato. *Plant Cell.* 2010;22(10):3206-3217.
 40. Francis D, Halford NG. Nutrient sensing in plant meristems. *Plant Mol. Biol.* 2006;60(6):981-993.

41. Enquist BJ, Kerkhoff AJ, Stark SC, Swenson NG, McCarthy MC, Price CA. A general integrative model for scaling plant growth, carbon flux, and functional trait spectra. *Nature*. 2007;449(7159):218-222.
42. Demura T, Ye ZH. Regulation of plant biomass production. *Curr. Op. Plant Biol.* 2010;13(3):298-303.
43. Shipley B. Net assimilation rate, specific leaf area and leaf mass ratio: Which is most closely correlated with relative growth rate? A meta-analysis. *Funct. Ecol.* 2006; 20(4):565-574.
44. Boonman A, Prinsen E, Gilmer F, Schurr U, Peeters AJ, Voesenek LA, Pons TL. Cytokinin import rate as a signal for photosynthetic acclimation to canopy light gradients. *Plant Physiol.* 2007;143(4):1841-1852.
45. Wu X, Zhu Z, Li X, Zha D. Effects of cytokinin on photosynthetic gas exchange, chlorophyll fluorescence parameters and antioxidative system in seedlings of eggplant (*Solanum melongena* L.) under salinity stress. *Acta Physiol. Plant.* 2012;34(416):2105-2114.
46. Van Staden J, Zazimalova E, George EF. Plant growth regulators II: Cytokinins, their analogues and antagonists. In: George EF, Hall MA, De Klerk GJ, editors. *Plant Propagation by Tissue Culture* Springer. The Netherlands; 2008.
47. Poorter H, Sack L. Pitfalls and possibilities in the analysis of biomass allocation patterns in plants. *Front. Plant Sci.* 2012;3:259.
48. Feller C, Favre P, Janka A, Zeeman SC, Gabriel JP, Reinhardt D. Mathematical modeling of the dynamics of shoot-root interactions and resource partitioning in plant growth. *PLoS One.* 2015;10: e0127905.
49. Benincasa P, Guiducci M, Tei F. The nitrogen use efficiency: meaning and sources of variation-case studies on three vegetable crops in central Italy. *Horttechnol.* 2011;21(3):266-273.
50. Kobe RK, Iyer M, Walters MB. Optimal partitioning theory revisited: Nonstructural carbohydrates dominate root mass responses to nitrogen. *Ecology.* 2010;91(1):166-179.
51. Yang Z, Midmore DJ. Modelling plant resource allocation and growth partitioning in response to environmental heterogeneity. *Ecol. Model.* 2005;181(1): 59-77.
52. Price CA, Weitz JS. Zero-sum allocational strategies determine the allometry of specific leaf area. *Am. J. Bot.* 2010;97(11):1808-1815.
53. Zwack PJ, Rashotte AM. Interactions between cytokinin signalling and abiotic stress responses. *J. Exp. Bot.* 2015;66(16):4863-4871.