

# Ms\_IJPSS\_49082.doc

*by*

---

**Submission date:** 06-May-2019 10:27 AM (UTC+0700)

**Submission ID:** 1125458709

**File name:** Ms\_IJPSS\_49082.doc (123K)

**Word count:** 3805

**Character count:** 20787

## Original Research Article

# Effect of municipal solid waste compost on soil chemical properties and growth performance of cocoa (*Theobroma cacao* L.) seedlings at the nursery in Ghana

### ABSTRACT

**Aims:** Availability of nutrients-rich topsoil for nursing cocoa seedlings is becoming limited and poor growth of cocoa seedlings in the nurseries has been ascribed to the use of unsuitable potting media. Experiments were conducted to investigate the suitability of compost in improving soil chemical properties and boost the growth of cocoa seedlings at the nursery.

**Study design:** The experiment was laid out in a Completely Randomized Design (CRD) with four replications.

**Place and Duration of Study:** The experiment was carried out at the main nursery of Cocoa Research Institute of Ghana, New Tafo-Akim, between September, 2014 and June 2015.

**Methodology:** Polybags were filled with soil obtained from an old cocoa plot (K6O2) at Cocoa Research Institute of Ghana. The soil has been classified as Rhodic-Lixic Ferrasol. Three soil: compost mixtures treatments, that is, 90:10, 80:20 and 70:30 % w/w were tested. A Standard foliar fertilizer and unamended soil were included as treated and untreated controls. Seedlings were raised from mixed hybrid cocoa and assessed at bi-monthly intervals for six months for growth. Pre and post treatments soil analyses were carried out using standard laboratory procedures.

**Results:** Initial soil analyses showed that OC (1.18 %), Ca (5.60 cmol kg<sup>-1</sup>), P (14.23 mg kg<sup>-1</sup>) and pH (5.63) were below the critical values required for good cocoa growth. The 70:30 soil: compost treatment produced significantly ( $P=0.05$ ) tallest plant (41.9 cm) with the unamended control the shortest (30.7 cm) at the end of the study. Residual pH (6.98), OC (2.30 %), P (14.23 mg kg<sup>-1</sup>) and Ca (13.02 cmol kg<sup>-1</sup>) were significantly ( $P=0.05$ ) higher under the same treatment compared to the unamended control; pH (5.36), OC (1.04 %), P (11.65 mg kg<sup>-1</sup>) and Ca (5.60 cmol kg<sup>-1</sup>).

**Conclusion:** Less fertile soils could be improved with the addition of Municipal Solid Waste (MSW) compost for raising good quality cocoa seedlings at the nursery in Ghana.

**Keywords:** [cocoa, compost, topsoil, seedling growth]

18

## 19 **1. INTRODUCTION**

20 The seed and seedling supply system of the Seed Production Division (SPD) and Cocoa  
21 Health and Extension Division (CHED) of COCOBOD is based on raising cocoa seedlings in  
22 nurseries using topsoil usually procured from contractors but the fertility of such topsoil could  
23 be problematic. Topsoil is becoming limited and poor growth of cocoa seedlings in the  
24 nurseries has been ascribed to the use of unsuitable potting media. This problem is  
25 compounded by inadequate quantities of fertile topsoil for potting cocoa [1; 2]. With the  
26 introduction of the cocoa rehabilitation programme in Ghana, different types of soils are used  
27 for filling the polybags. These soils differ in their fertility status, with the less fertile soils  
28 impacting negatively on the growth of cocoa seedlings [3; 4]. There is therefore, the need to  
29 search for suitable materials/soil additives with the view to improving the fertility status of the  
30 soils for raising healthy cocoa seedlings. A potting medium is a composition of organic  
31 materials formulated to achieve desirable chemical and physical needs required by the crop  
32 to attain its potential growth and development. According to [5], good potting media  
33 management is essential to the production of quality fruit tree seedlings, since vigorous  
34 growth is needed to face the seasonal hazards encountered on the field. The work by [6]  
35 indicated that cocoa seedlings need nitrogen, phosphorus, potassium and metabolites such  
36 as proteins, lipids, carbohydrates for their growth. Thus it is important that young nursery  
37 seedlings and transplanted seedlings are in optimal condition as far as their nutrient and  
38 energy status are concerned. Cocoa pod husk-based compost was used to raise cocoa  
39 seedlings at the nursery in Ghana [7]. However, soil amended with inorganic NPK fertilizer  
40 resulted in significantly poor seedling performance. The use of organic materials in potting  
41 media will not only improve the growth performance of cocoa seedlings but also improve the  
42 quality of soil used for raising the seedlings [8]. This will ensure adequate plant nutrients for  
43 the seedlings to boost its survival and establishment rates during field transplanting.  
44 However, the use of Municipal Solid Waste (MSW) compost in improving the fertility status of  
45 nutrient poor topsoil in the context of producing good quality cocoa seedlings at the nursery

46 in Ghana has not been studied. The objective of this study was therefore, to determine the  
47 effect of soil: compost mixtures on soil chemical properties and growth performance of cocoa  
48 seedlings in the nursery.

## 49 **2. MATERIALS AND METHODS**

### 50 **2.1 Study site**

51 The experiment was conducted at the main nursery of the Cocoa Research Institute of  
52 Ghana, New Tafo (latitude 6°13' N, longitude 0°22' W, altitude 222 m above sea level) in  
53 between September, 2014 and June, 2015.

### 54 **2.2 Soil sampling and analyses**

55 Topsoil (0-15 cm depth) collected from an old cocoa plot (K6O2) was used for the  
56 experiment. The soil at the site has been classified as Rhodic-Lixic Ferrasol [9] and belongs  
57 to the Wacri series according to the Ghanaian system of classification [10]. A sample of the  
58 soil was air-dried and sieved through a 2 mm mesh and stored for analysis of its physico-  
59 chemical properties. Soil samples were analyzed before and at the end of the experiment to  
60 find out the changes in soil properties following treatments application. Soil pH was  
61 determined using the glass electrode at soil: water ratio of 1:2.5 [11], organic carbon by the  
62 Walkley and Black wet oxidation method [12] and total N by the Kjeldahl digestion and  
63 distillation method [13]. Available phosphorus was measured by the Troug method [14].  
64 Exchangeable basic cations (K, Ca and Mg) were extracted with 1N ammonium acetate  
65 solution and the leachate analyzed by the Atomic Absorption Spectrophotometer [15].

### 66 **2.3 Nursery studies**

67 The experiment was conducted in the nursery at the Cocoa Research Institute of Ghana,  
68 Tafo. Standard nursery polybags measuring 18 cm x 25 cm were used for raising the cocoa  
69 seedlings [2; 16]. Different proportions of soil and Municipal Solid Waste compost (NPK 2-1-

0.5+0.85Ca+0.17Mg+0.65Fe+0.02Zn+18%OM) were mixed and used to fill the polybags. The following treatments were tested (i) Soil alone (ii) Standard foliar fertilizer (10 ml/11 liters (iii) Soil: compost (90:10 w/w) (iv) Soil: compost (80:20 w/w) and (v) Soil: compost (70:30 w/w). The polybags with the soil: compost mixtures were subsequently watered and allowed to settle for two weeks. Mixed hybrid cocoa seeds were sown at a seeding rate of two per polybag which were thinned to one seedling per polybags one month after sowing. The Standard foliar fertilizer was sprayed on the seedlings at monthly interval using pneumatic knapsack sprayer. Each treatment had thirty seedlings and the experiment was laid out in a Completely Randomized Design (CRD) with four replications. Watering was done as and when necessary. Seedling girth, height, number of leaves and dry matter production were measured at bi-monthly intervals for six months. Residual soil analyses were carried at six months after sowing.

82

## 83 **2.4 Data analysis**

84 Data collected were subjected to analysis of variance (ANOVA). Treatment means were compared using the least significant difference (LSD) method at  $P=.05$ . All statistics were performed using GenStat Statistical Package [17].

87

## 88 **3. RESULTS AND DISCUSSION**

### 89 **3.1 Initial soil fertility status**

90 Initial analysis showed that organic carbon, available P and exchangeable Ca contents of the soil were below the critical levels considered adequate for good cocoa growth [Table 1]. A good cocoa soil is reported to have organic carbon of above  $> 3\%$ ,  $20 \text{ mg kg}^{-1}$  available P and  $7.5 \text{ cmol kg}^{-1}$  exchangeable Ca respectively [18]. Total N, exchangeable K and Mg were above the critical values of  $0.09\%$ ,  $0.25 \text{ cmol kg}^{-1}$  and  $1.33 \text{ cmol kg}^{-1}$  respectively, considered adequate to support the growth of cocoa. The soil was moderately acidic with pH value below the soil critical limit required for cocoa.

97 **Table 1: Some chemical properties of the soil used in the experiment (0-15 cm)**

Parameter	Measured value
pH (soil: water, 1:2.5)	5.63
Organic carbon (%)	1.18
Total N (%)	0.16
Available P (mg kg <sup>-1</sup> )	14.23
Exchangeable cations (cmol kg <sup>-1</sup> )	
K	0.38
Ca	5.60
Mg	1.79

98

### 99 **3.2 Effect of the different treatments on soil chemical properties**

100 Soil chemical properties as affected by the application of MSW compost and foliar fertilizer at  
 101 the end of the study are presented in Table 2. Municipal Solid Waste (MSW) compost  
 102 application increased soil chemical composition compared to the untreated control and  
 103 Standard foliar fertilizer treatments. Soil nutrient content tended to increase with the level of  
 104 compost applied except exchangeable K. Residual soil properties decreased under the  
 105 Standard foliar fertilizer and unamended control treatments below the threshold values  
 106 required for good cocoa growth.

107

108

109

110

111

112

113

114

115 **Table 2: Effect of soil: compost mixture and foliar fertilizer on soil chemical**  
 116 **composition at 6 months of application**

117

Treatments	pH	% OC	%N	P (mg kg <sup>-1</sup> )	K (cmol kg <sup>-1</sup> )	Ca (cmol kg <sup>-1</sup> )	Mg (cmol kg <sup>-1</sup> )
T1 Control (soil alone)	5.36 <sup>c</sup>	1.04 <sup>c</sup>	0.15 <sup>c</sup>	11.65 <sup>d</sup>	0.20 <sup>c</sup>	5.60 <sup>d</sup>	1.22 <sup>d</sup>
T2 Standard foliar fertilizer	5.38 <sup>c</sup>	1.17 <sup>c</sup>	0.13 <sup>c</sup>	11.29 <sup>d</sup>	0.20 <sup>c</sup>	5.92 <sup>c</sup>	1.23 <sup>d</sup>
T3 Soil: compost (90:10)	6.43 <sup>b</sup>	2.03 <sup>b</sup>	0.21 <sup>b</sup>	23.64 <sup>c</sup>	0.30 <sup>b</sup>	8.01 <sup>b</sup>	1.49 <sup>c</sup>
T4 Soil: compost (80:20)	6.76 <sup>a</sup>	2.20 <sup>a</sup>	0.25 <sup>a</sup>	32.48 <sup>b</sup>	0.27 <sup>c</sup>	12.11 <sup>a</sup>	2.29 <sup>b</sup>
T5 Soil: compost (70:30)	6.98 <sup>a</sup>	2.30 <sup>a</sup>	0.27 <sup>a</sup>	40.17 <sup>a</sup>	0.30 <sup>a</sup>	13.02 <sup>a</sup>	2.55 <sup>a</sup>
Lsd ( <i>P</i> =.05)	0.27	0.22	0.03	2.45	0.02	0.28	0.15

118 Means in a column followed by the same superscript alphabets are not significantly different *P*>.05

### 119 3.3 Growth parameters

#### 120 3.3.1 Seedling Girth

121 Girth increments of cocoa seedlings due to the various treatments are presented in Table 3.

122 There was a significant (*P*=.05) seedling girth increment between the treatments at 2 months  
 123 after sowing. The values recorded ranged between 2.0 and 3.5 mm. Similar trends were  
 124 observed at 4 and 6 months after sowing. Seedling girth was generally bigger in the soil:  
 125 compost (70:30) treatments compared to the unamended soil and Standard foliar fertilizer  
 126 treatments.

127

128

129

130

131

132

133

134

135

**Table 3: Girth increments of cocoa seedlings grown in different soil: compost mixtures or sprayed with foliar fertilizer at 2, 4 and 6 months after sowing.**

Treatments	Seedling girth increments (mm)		
	2 months	4 months	6 months
T1 Control (soil alone)	2.0 <sup>c</sup>	3.8 <sup>c</sup>	5.1 <sup>c</sup>
T2 Standard foliar fertilizer	2.3 <sup>b</sup>	4.0 <sup>bc</sup>	5.7 <sup>bc</sup>
T3 Soil: compost (90:10)	2.4 <sup>b</sup>	4.1 <sup>b</sup>	5.9 <sup>b</sup>
T4 Soil: compost (80:20)	3.3 <sup>a</sup>	4.3 <sup>b</sup>	5.8 <sup>b</sup>
T5 Soil: compost (70:30)	3.5 <sup>a</sup>	5.3 <sup>a</sup>	7.3 <sup>a</sup>
Lsd ( $P=.05$ )	0.2	0.3	0.6

Means in a column followed by the same superscript are not significantly different ( $P=.05$ ).

### 3.3.2 Seedling Height

Table 4 shows height increments of cocoa seedlings under the different treatments. All the treatments except 90:10 soil: compost were significantly ( $P=.05$ ) taller than the untreated control 2 months after sowing. The 70:30 soil: compost treatment produced significantly ( $P=.05$ ) taller plants than the other treatments at 4 and 6 months after sowing. Height increments were linearly related to the quantity of compost applied.

**Table 4: Height increments of cocoa seedlings grown in different soil: compost mixtures or sprayed with foliar fertilizer at 2, 4 and 6 months after sowing.**

Treatments	Seedling height increments (cm)		
	2 months	4 months	6 months
T1 Control (soil alone)	23.1 <sup>c</sup>	24.6 <sup>c</sup>	30.7 <sup>c</sup>
T2 Standard foliar fertilizer	25.6 <sup>ab</sup>	28.2 <sup>b</sup>	37.9 <sup>ab</sup>
T3 Soil: compost (90:10)	24.8 <sup>ac</sup>	26.6 <sup>bc</sup>	36.4 <sup>b</sup>



T4 Soil: compost (80:20)	25.4 <sup>b</sup>	28.7 <sup>b</sup>	38.9 <sup>b</sup>
T5 Soil: compost (70:30)	27.3 <sup>a</sup>	32.8 <sup>a</sup>	41.9 <sup>a</sup>
Lsd ( $P=.05$ )	1.9	2.1	2.2

Means in a column followed by the same superscript are not significantly different ( $P=.05$ ).

### 3.3.3 Number of leaves

Seedlings grown on soil: compost mixture treatments developed significantly ( $P=.05$ ) more leaves relative to unamended soil and the Standard foliar fertilizer treatments at 2 months after sowing (Table 5). Seedlings grown on 70:30 soil: compost produced significantly ( $P=.05$ ) more leaves compared to the other treatments at 4 and 6 months after sowing.

**Table 5: Number of leaves of cocoa seedlings grown in different soil: compost mixtures or sprayed with foliar fertilizer at 2, 4 and 6 months after sowing**

Treatments	Number of leaves		
	2 months	4 months	6 months
T1 Control (soil alone)	4.3 (2.08) <sup>b</sup>	6.7 (2.58) <sup>c</sup>	10.0 (3.16) <sup>c</sup>
T2 Standard foliar fertilizer	5.1 (2.27) <sup>ab</sup>	8.7 (2.94) <sup>b</sup>	12.0 (3.46) <sup>b</sup>
T3 Soil: compost (90:10)	6.2 (2.48) <sup>a</sup>	8.3 (2.88) <sup>b</sup>	11.7 (3.42) <sup>b</sup>
T4 Soil: compost (80:20)	6.0 (2.45) <sup>a</sup>	9.2 (3.03) <sup>ab</sup>	12.8 (3.58) <sup>b</sup>
T5 Soil: compost (70:30)	6.3 (2.51) <sup>a</sup>	10.3 (3.21) <sup>a</sup>	16.3 (4.04) <sup>a</sup>
Lsd ( $P=.05$ )	1.5	1.4	1.8

Means in a column followed by the same superscript are not significantly different ( $P=.05$ ).

Values in bracket are square root transformation of actual values.

### 3.3.4 Dry matter production

Dry matter yield of the cocoa seedlings was found to increase with time for all the treatments [Table 6]. The variations in dry matter production between the fertilizer treatments at the different sampling periods were significant ( $P=.05$ ). Similar to the growth measurements (seedling height and girth increments and number of leaves), highest compost level (70:30

171 w/w) produced significantly ( $P=.05$ ) higher dry matter yield than the other treatments.

172

173 **Table 6: Dry matter yield of cocoa seedlings grown in different soil: compost mixtures**  
174 **or sprayed with foliar fertilizer at 2, 4 and 6 months after sowing**

Treatments	Dry matter (g plant <sup>-1</sup> )		
	2 months	4 months	6 months
T1 Control (soil alone)	0.87 <sup>b</sup>	1.96 <sup>c</sup>	3.35 <sup>c</sup>
T2 Standard foliar fertilizer	1.03 <sup>b</sup>	2.65 <sup>b</sup>	6.96 <sup>ab</sup>
T3 Soil: compost (90:10)	0.96 <sup>b</sup>	2.02 <sup>c</sup>	4.35 <sup>bc</sup>
T4 Soil: compost (80:20)	0.98 <sup>b</sup>	2.60 <sup>b</sup>	4.60 <sup>b</sup>
T5 Soil: compost (70:30)	1.43 <sup>a</sup>	4.69 <sup>a</sup>	8.23 <sup>a</sup>
Lsd ( $P=.05$ )	0.20	0.49	1.13

175 Means in a column followed by the same superscript are not significantly different ( $P>.05$ )

176

#### 177 4. DISCUSSIONS

178 The observed significant effects of soil: compost mixture on the growth (girth and height) and  
179 total biomass production of cocoa seedlings in this present study could be attributed to the  
180 bioavailability of vital nutrients in the compost. Similar effect of compost on the growth of  
181 cocoa seedlings has been reported by [19]. This result indicates that the seedlings that were  
182 raised in soil: compost mixture are of higher growth performance and would have higher  
183 survival and establishment rates after field transplanting than those raised in sole soil and  
184 seedlings treated with foliar fertilizer. The observations made in this present study confirm  
185 earlier findings on growth of cocoa seedlings [7; 20].

186

187 It is observed in this study that number of leaves of cocoa seedlings and total dry plant  
188 biomass increased with increasing compost rate. This emphasizes the importance of the  
189 compost in providing nutrient for the growth of the cocoa seedlings as noted by [20]. Better  
190 cocoa seedlings growth performance with the use of cocoa pod husk-based compost as

191 potting medium was reported by [7]. However, on the contrary, the authors observed that  
192 potting media amended with inorganic NPK fertilizer significantly produced narrower leaves.  
193 The high total dry plant biomass production of cocoa seedlings in soil: compost mixture  
194 (70:30 w/w) could be attributed to the chemical composition of the compost and ability to  
195 release nutrients through mineralization for plant uptake.

196  
197 Application of MSW compost to the less fertile topsoil used in this experiment improved the  
198 soil chemical properties at the end of the study compared to the unamended control and the  
199 foliar fertilizer treatments. The use of organic fertilizers has been associated with desirable  
200 soil characteristics including higher plant available nutrients, water holding capacity, CEC  
201 and lower bulk density, and can foster beneficial microorganisms [(21; 22]. Benefits of  
202 organic soil amendments also include pH stabilization and increased water infiltration rate  
203 due to enhanced soil aggregation. Since they are the ultimate slow-release fertilizers, it's  
204 very difficult to over fertilize (and harm) plants. There's little to no risk of toxic buildups of  
205 chemicals and salts that can be deadly to the cocoa seedlings.

206  
207 The residual soil nutrients were high in the soil: compost mixtures compared to the  
208 unamended control and Standard foliar fertilizer treatments. The soil: compost mixture  
209 treatments increased the levels of carbon in the soil, which leads to an increase in fertility  
210 because of an increase in microorganism activity using carbon as energy source [23].  
211 Similarly, high residual total N and available P were recorded by soil: compost mixtures.  
212 According to [24], the application of organic soil amendments increased the N content in soil  
213 because of the greater N and organic C concentration in the amendment. Organic fertilizers  
214 have been suggested to increase the availability of P because as the organic component  
215 decomposes it releases CO<sub>2</sub>, and higher CO<sub>2</sub> concentrations would increase the rate of  
216 decomposition of phosphate minerals and thereby increase soil available P [25]. These  
217 minerals synthesize phospho-humic complexes that are available to the plant and allow for

218 the exchange of organic radicals by phosphates. The increase in pH, K, Ca and Mg recorded  
219 for the Soil: compost mixtures treatments compared to the unamended control and Standard  
220 foliar fertilizer treatments could be attributed to the increased availability of organic matter  
221 and release of some cations from the decomposed organic amendment as noted by [26].  
222 The low residual soil nutrients value recorded under the unamended control could be  
223 ascribed to nutrient mining by the cocoa seedlings without soil amendment. This observation  
224 is consistent with the fact that organic fertilizers are a natural source of macro and  
225 micronutrients. The above findings corroborate the need for application of organic  
226 amendments to less fertile topsoil in raising healthy cocoa seedlings to ensure balanced  
227 nutrition.

228

229 The high residual soil nutrients observed under soil: compost mixtures treatments meant that  
230 application of compost to less fertile topsoil could improve its fertility status and also ensure  
231 that adequate nutrients are available to the cocoa seedlings during field transplanting.

232

## 233 **5. CONCLUSION**

234 The results showed that MSW compost increases the level of soil organic carbon, total N,  
235 available P, soil pH and exchangeable cations of nutrients poor topsoil. Subsequently, the  
236 compost increased the growth of the cocoa seedlings compared to the unamended control  
237 and Standard foliar fertilizer. The compost is found to be a suitable source of nutrients for  
238 improving the fertility status of less fertile topsoil for raising good quality cocoa seedlings at  
239 the nursery in Ghana.

240 **COMPETING INTERESTS**

241 Authors have declared that no competing interests exist.

242 **REFERENCES**

- 243 1. Donkor MA, Henderson CP, Jones AP. Survey to quantify adoption of CRIG  
244 recommendations. FSU Research Paper 3. Tafo. Cocoa Research Institute of  
245 Ghana; 1991.
- 246 2. Ofori-Frimpong K, Afrifa AA, Appiah MR. Improving the growth of cocoa seedlings in  
247 nursery by the use of fertilizers. Journal of Ghana Science Association 2006;  
248 8(2):85-91
- 249 3. Oyewole OS, Ajayi IM, Rotimi RI. Growth of cocoa (*Theobroma cacao* L.) seedlings  
250 on old cocoa soils amended with organic and inorganic fertilizers. African Journal of  
251 Agricultural Research. 2012; 7(24):3604-3608.
- 252 4. Quaye AK, Konlan S, Arthur A, Pobee P, Dogbatse JA. Media type and compost  
253 mixtures effect on growth and nutrient uptake of cocoa seedling at the nursery in  
254 Ghana. Proceedings, 2017 International Symposium on Cocoa Research (ISCR),  
255 Lima, Peru; 2017.
- 256 5. Khan MM, Khan MA, Mazhar A, Muhammad J, Ali JMA, Abbas H. Evaluation of  
257 potting media for the production of rough lemon nursery stock. Pakistan Journal of  
258 Botany. 2006;38(3) 623-629.
- 259 6. Gockowski J, Weise SF, Sonwa D, Tchata M, Ngobo M. Conservation because it  
260 pays: shaded cocoa agroforests in West Africa. Paper presented at National  
261 Academy of Sciences conference on the science behind cocoa's benefits,  
262 Washington, D.C; 2004.
- 263 7. Ofori-Frimpong K, Afrifa AA, Acquaye S. Relative efficacy of cocoa pod husk-based  
264 compost on growth and nutrient uptake of cocoa seedlings in the nursery. Ghana  
265 Journal of Agricultural Science. 2010; 43: 1

- 266 8. Adejobi KB, Akanbi OS, Ugioro O, Adeosun SA, Mohammed I, Nduka BA. et al.  
267 Comparative effects of NPK fertilizer, cowpea pod husk and some tree crops wastes  
268 on soil, leaf chemical properties and growth performance of cocoa (*Theobroma*  
269 *cacao* L.). African Journal of Plant Science. 2013; 8(2): 103-107.
- 270 9. World Reference Base for Soil Resources. IUSS Working Group WRB, International  
271 soil classification system for naming soils and creating legends for soil maps. World  
272 Soil Resources Reports No. 106. FAO, Rome; 2014.
- 273 10. Dwomo O, Dedzoe CD. Oxisol (Ferralsol) Development in Two Agro-Ecological  
274 Zones of Ghana: A Preliminary Evaluation of Some Profiles. Journal of Science and  
275 Technology. 2010;30(2);1-11.
- 276 11. McLean EO. Soil pH and lime requirement. In: Page AL, Miller RH, Keeney DR,  
277 editors. Methods of soil analysis. Part 2. Chemical and microbiological properties.  
278 Second edition. American Society of Agronomy and Soil Science Society of  
279 America, Madison, Wisconsin USA; 1982.
- 280 12. Nelson DW, Sommers LW. Total carbon, organic carbon and organic matter. In:  
281 Page AL, Miller RH, Keeney DR, editors. Methods of soil analysis. Part 2. Second  
282 edition. Chemical and microbiological properties. American Society of Agronomy  
283 and Soil Science Society of America. Madison, Wisconsin USA; 1982.
- 284 13. Bremner JM, Mulvaney CS. Total nitrogen. In: Page AL, Miller RH, Keeney DR,  
285 editors. Methods of soil analysis. Part 2. Chemical and microbiological properties.  
286 American Society of Agronomy and Soil Science Society of America, Madison  
287 Wisconsin USA; 1982.
- 288 14. Truog E. The determination of the readily available phosphorus in soils. Journal of  
289 American Society of Agronomy. 1930;22: 874-882.

- 290 15. Thomas GW. (1982). Exchangeable cations -In Page AL, Miller RH, Keeney DR,  
291 editors. Methods of soil analysis, part 2; Chemical and Microbiological properties.  
292 Madison, WL; Soil Science Society of America, Madison Wisconsin USA; 1982.
- 293 16. Oppong FK, Ofori-Frimpong K, Fiakporku R. Effect of polybags size and foliar  
294 application of urea on cocoa seedling growth. Ghana Journal of Agricultural Science.  
295 2008;40:207-213.
- 296 17. GenStat. GenStat Release 11.1. Eleventh edition. Lawes Agricultural Trust  
297 (Rothamsted Experimental Station). <http://www.vsni.co.uk>. 2008
- 298 18. Ahenkorah Y. The influence of environment on growth and production of the cacao  
299 tree: soils and nutrition. Proceedings of the 7<sup>th</sup> International Cocoa Research  
300 Conference, Douala, Cameroon; 1981.
- 301 19. Adejobo KB, Famaye AO, Akanbi OS, Adeosun SA, Nduka AB, Adeniyi DO.  
302 Potentials of cocoa pod husk ash as fertilizer and liming materials on nutrient uptake  
303 and growth performance of cocoa. Research Journal of Agriculture and Environment  
304 Management. 2013;2 (2): 243-251.
- 305 20. Sosu G. Growth of cocoa seedlings as affected by different growth media and  
306 different polybag sizes. A Thesis Submitted to the University of Ghana, Legon in  
307 Partial Fulfillment of the Requirements for the Award of Mphil (Crop Science)  
308 Degree. 2014.
- 309 21. Doran J. Building soil quality. In: Proceedings of the 1995 Conservation Workshop  
310 on Opportunities and Challenges in Sustainable Agriculture. Red Deer, Alta.,  
311 Canada, Alberta Conservation Tillage Society and Alberta Agriculture Conservation,  
312 Development Branch, 1995.
- 313 22. Drinkwater LE, Letourneau DK, Workneh F, van Bruggen AHC, Shennan C.  
314 Fundamental differences between conventional and organic tomato agroecosystems  
315 in California. Applied Ecology. 1995(5): 1098–1112.

- 316 23. Vargas GMC, Suárez F. Effect of application of compost on soil biological  
317 properties. In: Moreno CJ, Moral HR, editors. Composting. Mundi-press Madrid;  
318 2007.
- 319 24. Wu SC, Cao ZH, Li ZG, Cheung MKC, Wong WH. Effects of biofertilizer containing  
320 N fixer, P and K solubilizers and AM fungi on maize growth: a greenhouse trial.  
321 Geoderma. 2005;125:155-166.
- 322 25. Núñez ER. The soil as a natural medium in the nutrition of crops. In: Alcántar, G.G  
323 Trejo-Téllez LI, editors. Crop nutrition, Mundi Press. College of Postgraduates,  
324 México; 2007.
- 325 26. Agbede TM, Adekiya AO. Effect of Cocoa Pod Ash and Poultry Manure on soil  
326 properties and cocoyam productivity of nutrient-depleted Alfisol. International  
327 Journal of Agricultural and Biosystems Engineering. 2016;10(3);172-179.
- 328  
329  
330  
331  
332  
333



ORIGINALITY REPORT

0%

SIMILARITY INDEX

0%

INTERNET SOURCES

0%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

Exclude quotes On

Exclude bibliography On

Exclude matches < 3%