

**IDENTIFICATION AND MANAGEMENT OF PESTS AND DISEASES
OF GARDEN CROPS IN SANTA, CAMEROON**

ABSTRACT

Agriculture is one of the pillars of the economy of Cameroon although in most instances it is still small scaled and depends largely on house hold labour, with about 70% of the active population engaged in agricultural activities. The Western Highlands of Cameroon is noted for its high involvement in agriculture especially the cultivation of vegetable crops such as Cabbage (*Brassica oleracea var capitata* L.), carrots (*Daucus carota* L.), leeks (*Allium porrum* L.), tomatoes (*Lycopersicon esculentum* Mill.), celery (*Apium graveolens* L.) and onions (*Allium cepa* L.) (Abang *et al.*, 2013). Generally this work sought to identify the pests and diseases that hinder successful gardening and how they are managed in the Santa community of Mezam Division. This work was carried out in Santa, a Sub-division in Mezam Division of the North West Region of Cameroon. On each of the farms, an area of 20 x 20 m was mapped out and the plants in that area observed. The crops were observed for pests and nature of damage inflicted, at each growth stage. The main diseases identified here were clubroot disease (*Plasmodiophora brassicae*) and late blight (*Phytophthora infestans*) while Aphids (*Myzus persicae* S.), whiteflies (*Bemisia tabaci*) fruit worms (*Helicoverpa amigera*), Cutworms, black garden ants (*Lasius niger* L.) and grasshoppers (*Zonocerus variegatus*) were the prominent pests. The most applied pesticides were Cypercal, Parastar, Banko plus, Manozane, Mancozan, Pencozeb, Gramoxon and Action80. From this study the most prominent pest of cabbage here was the black cutworm (*Agrotis ipsilon*) which affected the early growing stage. The main disease that affected tomato was blight. This was seen in both seasons but the severity of attack was greater in the rainy season. Fontem, (1993) in a study on the severity of tomato diseases in Cameroon found that blight is the most severe disease in the wet season in Cameroon and is widely distributed on foliage and fruits. From this study it is seen that the most prominent insect pest is the cutworm (*Agrotis ipsilon*). The main disease of cabbage was clubroot disease. It is also noted from the research that insect pests were a main problem in the dry season causing high economic losses while there was reduced infestation in the wet season. In this regard, there is an urgent need to educate the Santa gardeners on good agricultural practices through Integrated Crop and

Comment [A1]: Division of

35 Pest Management (ICPM) practices which will include both cultural, physical or
36 mechanical, biological and chemical pests control methods.

37 **Key Words:** Pests, Diseases, Pesticides, Santa

Comment [A2]: Keywords should be minimum five meaningful and alphabetically arranged.

38 INTRODUCTION

39 Agriculture is one of the pillars of the economy of Cameroon though in most instances it
40 is practiced at small scale and depends largely on house hold labour, with about 70% of
41 the active population of this country engaged in it. Also this sector is responsible for
42 providing food security to both the rural and urban populations of this country via local
43 production (Wilfred *et al.*, 2016). The Western Highlands of Cameroon is noted for its
44 high involvement in agriculture especially the cultivation of vegetable crops such as
45 cabbage (*Brassica oleraceavar capitata* L.), carrots (*Daucus carota* L.), leeks (*Allium*
46 *porrum* L.), tomatoes (*Lycopersicon esculentum* Mill.), celery (*Apium graveolens* L.) and
47 onions (*Allium cepa* L.) (Abang *et al.*, 2013). The main areas noted for this production of
48 these garden crops in Cameroon are Santa in the North West and Foumbot in the West
49 Regions. Their cultivation has brought about an increase in agricultural production that is
50 used to feed the nation. Among various economic and social benefits, market gardening
51 has a vital and multifaceted role in providing food security, meeting the demands of
52 consumer markets, utilising labour and generating income. It can provide both personal
53 satisfaction and supplementary or even full-time income. The income generated from
54 market gardening also provides indirect socio-economic benefits for market gardeners,
55 such as greater access to household items (televisions, chairs) and greater mobility from
56 the purchase of motor vehicles, motorbikes or bicycles (Porter *et al.*, 2003). As urban
57 centres expand, the demand for fresh garden produce increases and the land devoted to
58 market gardening also expands, usually in the periphery (Friesen, 1998). This is
59 particularly true in developing countries where rapid urbanisation is prevalent.

60 Yield and quality are central to sustainable vegetable production. If not properly
61 managed, pests and diseases can dramatically reduce crop yield quality and subsequent
62 returns. At this economic injury level, there is the need to employ control measures,
63 which may have a great negative effect on the practice of market gardening if not properly
64 managed. Today, pests and diseases are better managed using an integrated approach and
65 this approach brings together the best mixture of chemical, biological and cultural
66 methods to manage pests and diseases. To successfully apply any management strategy

against pests or diseases, the first step is to identify them correctly for appropriate action to be taken and this gave reason for this work to be carried out to identify the pests and diseases that hinder successful gardening and how they are managed in the Santa community of Mezam Division.

MATERIALS AND MEHODS

Study area

This study was carried out in Santa which is one of the Sub-divisions in Mezam Division of the North West Region of Cameroon. It is located between latitudes 5° 42' and 5° 53' north of the equator and longitudes 9° 58' and 10° 18' east of the Greenwich Meridian (Santa Rural Council Monographic Study, 2003). The population of this area estimated in 2008 was 99851(Fogwe, 2014) and 90% of this population are engaged in farming and grazing. It covers a surface area of about 532.67 km². It is bordered to the North by Bamenda Sub Division, to the West by Bali and Batibo Sub-Divisions, to the South by Wabane, Babadjou and Mbouda and to the East by Galim (Sonchieu *et al.*, 2017).

The mean annual temperature of the area varies from 21.8 to 30.8°C. The annual rainfall is between 2000 -3000 mm mostly from March to September and the dry season from October to February. The soils in this area are fertile and support a large human population. The altitudinal range is from 600 to 2600 m making this highland favourable for animal rearing, crop and vegetable cultivation aptly qualifying this area as an agricultural production basin in the Western Highlands of Cameroon.




Figure 1. Map of Santa Sub Division showing the different villages

Identification of pests and diseases and their mitigation

Identification of pests and diseases was carried out three times on six farms for each (two at the upper, two at the middle and two at the lower Santa) during different growth stages of tomatoes, cabbage, potatoes, leeks and celery. They were observed from seedling through flowering to maturity.

On each of the farms, an area of 20 x 20 m was mapped out and the plants in that area observed. The crops were observed for pests and nature of damage inflicted, at each growth stage. The parts of the crops observed were stems, leaves, flowers and fruits. The crops were also observed for diseases. The observation for pests and diseases was done for the dry and rainy seasons. Also the types of pesticides, and their frequency of application these two seasons to combat pests and diseases by farmers on the different crops were noted.

Comment [A3]: X replace by x

Comment [A4]: pesticides

DATA COLLECTION

Pre-designed data recording forms were used in gathering information on the following variables: type of pesticide, frequency of application, insect pests and nature of damage, diseases and nature of damage.

STATISTICAL ANALYSIS

Data was entered into Microsoft excel. Descriptive (frequency and percentage) statistics was used to represent the results.

RESULTS

Pests of Selected Garden Crops and Management Practices

Cabbage (*Brassica oleracea*)

The insect pests common with cabbage at transplant stage were cutworms (*Agrotis ipsilon*), which eat through the stems of the crop at the ground level and made the crop to fall, whiteflies (*Bemisia tabaci* L), aphids (*Myzus persicae* L) and fruitworm (*Helicoverpa armigera* L) . The farmers used cypercal and parastar for their control during dry season. During this transplant stage, there was no disease affecting cabbage (Table 1). Whiteflies, aphids and fruit worms affected the crop mostly in the dry seasons while the only pest of economic importance was the rainy season was cutworms.

At the flowering and maturation stages, whiteflies, aphids, fruitworms and grasshoppers were seen and same chemicals used for their control as during transplant. At the flowering and maturation stages clubroot was the only disease affecting cabbage and no pesticide was applied for its control (Table 1).

Comment [A5]: fruit worms

Comment [A6]: club root

152 Table 1: Pests and diseases of cabbage, pesticides and their frequency of application used for their control

	Growth stage					
	Transplant		Flowering		Maturity	
	Pests	Disease(s)	Pests	Disease(s)	pests	Disease(s)
Pest/ disease	Cutworms, whiteflies, aphids and fruit worms	-	whiteflies, aphids, fruit worms, and grasshoppers	Clubroot	whiteflies, aphids, fruit worms, and grasshoppers	Clubroot
Pesticide	***Cypercal ***Parastar	-	***Cypercal ***Parastar	-	***Cypercal ***Parastar	-
Frequency during dry season	Thrice	-	Thrice	-	Thrice	-
Frequency during rainy season	twice	-	Twice	-	Twice	-

Comment [A8]: club root

Comment [A9]: club root

Comment [A7]: worms, and

153 ***=Dry season, **=Rainy season

154

155 **Tomato** (*Lycopersicon esculentum*)

Comment [A10]: *Lycopersicon esculentum*

156 The major insect pest that damaged tomato at transplant stage was cutworm. The
157 other insects at this stage were crickets (*Gryllus sp.*) and spider although their
158 damage was not severe. Cypercal, Parastar and Cypercot were used against the
159 cutworms. Blight was observed at this stage but did not cause severe damage in the
160 dry season and was only sprayed in the rainy season with Mancozeb and Mancozane
161 (Table 2).

162 The flowering stage of this crop suffered from a new set of pests. These were fruit
163 worms (*Helicoverpa amigera* L), fruitflies (*Dacus puntatifrons* L), aphids, leaf
164 miners and to a lesser extent the cutworms. The fruit worms ate through the fruits,
165 fruitflies stung the fruits creating black spots on them, the leaf miners mined the
166 leaves and cutworms present at this stage did not have major effects because the
167 stems of the plant were already hardened. The insecticides used at this stage were
168 Cypercal, Parastar, Cypercot (Table 2).

Comment [A11]: Fruit flies

169 In the third stage of growth when the crop had reached maturity, the pests were fruit
170 worms, aphids and whiteflies. Blight was also present and caused damage such as
171 fruit rot, irregular ripening of fruits, some dropping to the ground and leaves
172 yellowing and dry off. The chemicals used to spray were still those used at the
173 flowering stage with insecticides being sprayed at higher frequencies per month
174 (Table 2).

175 Table 2: Pests and diseases of tomato, pesticides and their frequency of application used for their control

	Growth stage					
	Transplant		Flowering		Maturity	
	Pests	Disease(s)	Pests	Disease(s)	Pests	Disease(s)
Pest/ disease	Cutworms, crickets, spiders	Blight	Cutworms, whiteflies, aphids, fruit worms, fruit flies.	Blight	Fruit worms, Aphids, Whiteflies, Fruit flies,	Blight
Pesticide	***Cypercal ***Parastar ***Cypercot	**Pencozeb	***Cypercal ***Parastar	**Banko plus, **Manozane, **Mancozan, **Pencozeb	***Cypercal ***Parastar	**Banko plus, **Manozane, **Mancozan, **Pencozeb,
Frequency during dry season	Twice	Thrice	Four	Eight	Thrice	Four
Frequency during rainy season	Thrice	Four	Thrice	Twelve	Four	Sixteen

176 ***=Dry season, **=Rainy season

Celery (*Apium graveolens* L.)

The most prominent pest of celery at transplantation during dry and rainy seasons was the cutworm which feed on the stem of the celery plant. Another insect seen at this growth stage was cricket that ate through the leaves creating holes on them. The insecticides used for pests control were Cypercal, Parastar and Cypermax which were pyrethroids. Blight was the lone disease during all stages and was managed using pencozeb and balear at transplant stage. These different pesticides were either sprayed once or twice a month (Table 3).

In the second growth stage, cutworms were still seen and whiteflies and aphids were mostly seen in the dry season. Pests were managed using cypermax, cypercal or parastar. Blight was controlled with Balear, Banko plus, Mancozeb, Manozane or Pencozeb. They were used only once at this stage (Table 3).

At maturity, leafminers were the only insects seen during dry season were sprayed two times with Parastar, and Cypercal, at this stage for insect pests. Blight had its damaging effects at this stage mostly in the rainy season and Pencozeb, Mancozeb, Manozane, Banko plus and Balear were used for its control (Table 3).

Leeks (*Allium porrum*L.)

The main pest of leek at transplant was cutworm that fed on the stems of the plant cutting through and was managed using cypercal, Parastar and Fastac as the main insecticides to kill these cutworms in the farms (Table 4).

At flowering and maturation stages, the main insect pest was aphid which was controlled with Parastar and Callidim during flowering and cypercal during maturation. Blight affected the crops causing the leaves to turn yellow at the flowering and maturation stages mostly during rainy season. It was controlled with Manozane, Moncozeb or Pencozeb during flowering and Pencozeb, Manozane and Moncozan at maturation. In the rainy season blight was sprayed 7 to 8 times in a month. In the dry season the effect of blight was very minimal that some farmers did not spray their farms with the fungicides (Table 4).

Comment [A12]: *Allium porrum* L.

208 Table 3: Pests and diseases of celery, pesticides and their frequency of application used for their control

	Growth stage					
	Transplant		Flowering		Maturity	
	Pest	Disease(s)	Pests	Disease(s)	Pests	Disease(s)
Pest/ disease	Cutworms	Blight	Cutworms, Whiteflies, Aphids	Blight	Leaf miners,	Blight
Pesticide	***Parastar ***Cypercal, ***Cypermax	-	**Parastar, **Cypercal	**Pencozeb **Balear **Banko **Mancozeb, **Manozane, Plus	***Parastar, ***Cypercal,	**Pencozeb, **Mancozeb, **Balear,
Frequency during dry season	Thrice	-	Four	Four	Five	Five
Frequency during rainy season	Thrice	-	Thrice	Sixteen	Five	Twenty

209 ***=Dry season, **=Rainy season,

210 Table 4: Pests and diseases of leek, pesticides and their frequency of application used for their control

	Growth stage					
	Transplant		Flowering		Maturity	
	Pest	Disease(s)	Pest	Disease	pest	Disease
Pest/ disease	Cutworms	-	Aphids	Blight	Aphids,	Blight
Pesticide	***Cypercal, ***Parastar *** Fastac	-	***Parastar, ***Callidim	**Manozane, **Mancozeb	***Cypercal,	**Pencozeb, **Mancozeb, **Manozane
Frequency during dry season	Twice	-	Twice	Seven	Twice	Seven
Frequency during rainy season	Twice	-	Once	Seven	Once	Eight

211 ***=Dry season, **=Rainy season

212

213 **Potato (*Solanum tuberosum*)**

214 During sprouting, the pest of potato was cutworms and the severity was greater in the
215 dry than rainy season with Parastar, Cypercal and Fastac used to control it. At this
216 early growth stage blight was also observed. This was more of economic importance
217 in the rainy than dry season. Most farmers did not bother about blight at this stage,
218 but the few who did used Ridomil for its control (Table 5).

219 At the flowering stage, the insect pests were fruitworms and aphids. The pesticides
220 used for their control were Parastar, Cypercot and Fastac. Blight was persistent at this
221 stage causing leaves to turn yellow and eventually drying off. Pencozeb, Manozane
222 and Mancozeb were the main fungicides used to tackle blight at this stage. Bacteria
223 wilt was also noticed at this stage. Crops affected by bacteria became yellowish in
224 nature and withered. When uprooted the potato tuber inspected was watery and soft
225 in texture.

226 At maturity, aphids, fruitworms and blight were still persistent. The insects were
227 sprayed with Parastar. Plantineb, Pencozeb and Balear were the main fungicides used
228 against blight at this stage (Table 5).

229

230

231

232

233

234

235 Table 5: Pests and diseases of potato, pesticides and their frequency of application used for their control

	Growth stage					
	Transplant		Flowering		Maturity	
	Pest	Disease	Pests	Diseases	pests	Disease
Pest/ disease	Cutworms	Blight	Fruitworms, Aphids, Leafminers,	Blight Bacteria wilt	Aphids, Fruitworms	Blight
Pesticide	***Parastar, ***Cypercal, ***Fastac,	Ridomil	***Parastar, ***Fastac,	**Pencozeb, ***Monozane, ** Mancozeb	***Parastar, ***Fastac,	**Pencozeb, ***Monozane ** Mancozeb
Application frequency for dry season	Thrice	-	Thrice	Seven	Twice	Seven
Application frequency for rainy season	-	Twice	Twice	Eight	Twice	Seven

236

237 ***=Dry season, **=Rainy season

238

239 **DISCUSSION**

240 Six cabbage farms observed through the growth stages revealed insect pests as the main
241 problem to proper cabbage growth. The pests were cutworms, fruit worms, aphids and
242 whiteflies. This is in line with the findings of Dzomekuet *al.* (2011). The most prominent
243 pest of cabbage here was the black cutworm (*Agrotis ipsilon*) which affected the early
244 growing stage. Norida and John (2005), in Malaysia found *A. ipsilon* to be recognized by
245 80% of the farmers during the early growing period. As opposed to the findings of
246 Talekar and Shelton, (1993) who found diamondback moth (*Plutella xylostella*) to be the
247 most prominent pest of cabbage worldwide, diamondback moth was not noticed in this
248 area. This might be due to climatic factors that do not favour its survival in the Santa
249 area or the farmers sprayed with the insecticides and controlled its population. A range
250 of insecticides were used to kill insect pests by the farmers, at different spraying
251 frequencies. The insects caused much damage in the dry season than in the rainy season.
252 This is because the populations of insects were higher during dry season compared with
253 rainy season. This is in conformity with studies by Nsobinenyui *et al.* (2017) who
254 indicated insects are more abundant in dry season than rainy season. This might be due
255 to increase temperatures. Increase temperature is known to speed up the life cycle of
256 insects leading to faster increase in pest population. It has been estimated that a 2°C
257 increase in temperature has the potential to increase the number of insect life cycles by
258 one to five times (Bale et al., 2002; Petzoldt and Seaman, 2010). The main disease of
259 cabbage in this area was the clubroot disease (*Plasmodiophora brassicae*), commonly
260 called 'Ginger' in this area, that affected the roots of the cabbage plant. Here the disease
261 did not respond to any pesticide. The only method farmers had to use was to practice
262 crop rotation to disrupt the life cycle of the fungus.

263 The different tomato farms observed experienced the same kind of pests at the same
264 level of the plant growth. In the dry season the effects of insect pests were more than that
265 of fungi on the crop. Many more farmers spray against insects than diseases in the dry
266 season and this could suggest that insect pests are more serious in the dry season. The
267 main insect pests of tomatoes in the dry season that caused economic damage were
268 cutworms, aphids, fruit flies, leaf miners, whiteflies and fruit worms. This is also

269 reported by Sait (2003). The main disease that affected tomatoes was blight. This was
270 seen in both seasons but the severity of attack was greater in the rainy season.
271 Fontem,(1993) in a study on the severity of tomato diseases in Cameroon found that
272 blight is the most severe disease in the wet season in Cameroon and is widely distributed
273 on foliage and fruits. The plant was affected at all the three growth stages examined by
274 this blight.

Comment [A13]: Sait, (2003).

275 A wide range of pests affected the celery crops in the fields observed. Key insect pests
276 here included cutworms, whiteflies, aphids, crickets and fruit worms. Blight was also
277 seen affecting the crops. Farmers relied heavily on the use of pesticides to control these
278 pests (Ntonifor *et al.*, 2013). Producers used a wide range of pesticides, as many farmers
279 believe that the only way to tackle pest problems was to use pesticides.

Comment [A14]: Fontem, (1993)

280 Insects affected the potatoes in the field at its different growth stages. Some affected the
281 foliage, some tubers and some transmitted diseases as seen in the findings of Radcliffe
282 and Ragsdale, (2002). The effect was greater in the dry season. Blight was less important
283 in the dry season such that some farmers did not spray their farms against this disease
284 during this season. Blight caused the greatest damage in the rainy season (Fontem,
285 1993). Bacteria disease was also a problem in the farms as crops were affected by this
286 disease leading to low yields. This is in line with the findings of Kaguongo *et al.*, (2008)
287 who pointed out bacteria as an important disease contributing to yield reduction and
288 considered it more problematic than blight since it has no known chemical control
289 procedures and many farmers do not know how to control it.

290 CONCLUSION

291 From this study it is seen that the most prominent insect pest is the cutworm. This insect
292 pest is seen to attack all the crops that were used in this study. They attack primarily at
293 the stage when the crop has just been transplanted due to the fact that the stems of the
294 crops are still very tender and so they can chew through during feeding with their
295 mandibles. Other insect pests noted in this study were aphids, crickets, whiteflies, fruit
296 flies, leaf miners and black ants. These insects were all treated with insecticides.

297 The main insecticides that the farmers here used were Parastar and Cypercal with the
298 prominent fungicides being Pencozeb and Mancozeb. Also these farmers used herbicides

of which the popular ones were Gramoxone, Tromissil and Action 80. Each group of these pesticides had almost the same active ingredients

All these crops suffered from fungal attack except the cabbage plant that was affected mainly by insect pests. This fungus that attacked the crops was *Pythophthora infestans* commonly known as blight. It caused the leaves of Tomato, potato, celery and leeks to become yellow and eventually dry off. Bacterial wilt was also reported in the potato farms that were observed.

The findings of this study also present another pest which affects only cabbage called clubroot disease and it affects the roots of the crop such that the roots do not extend into the soil, and thus the crop would wither and die as a result of no water being drawn up by the roots as they were damaged.

It is also noted from the research that insect pests were a main problem in the dry season causing high economic losses while there was reduced infestation in the wet season. Blight affected crops more in the rainy season than in the dry season.

REFERENCES

- Abang, A.F., Kouame, C.M., Abang, M., Hannah, R. and Fotso, A.K. (2013). Vegetable growers perception of pesticide use practices, cost, and health effects in the tropical region of Cameroon. *International journal of Agronomy and Plant Production*. **4(5)**: 873- 883
- Bale, J.S., Masters, G.J., and Hodkinson, I.D., Awmack, C., Bezemer, T.M., Brown, V.K. *et al.* (2002). Herbivory in global climate change research: Direct effects of rising temperatures on insect herbivores. *Global Change Biology*, **8**: 1-16.
- Dzomeku, K., Abudulai, M. and Abukari, M. (2011). Influence of weeding regime and neem seed extract on the population of insect pests and yield of cabbage in the Guinea savannah zone. *Agriculture and Biology Journal of North America*. **(6)**: 921-928

Comment [A15]: Bold font

328 FOGWE Z. N and Bonglam C. Z. (2016): Perception and Adaptation Adjustments to
 329 Climate Variability Within the Santa Agrarian Basin in the Western
 330 Highlands of Cameroon. *Journal Of Humanities And Social Science*. Volume
 331 21, Issue 12, e-ISSN: 2279-0837, p-ISSN: 2279-0845.
 332 Fogwe, Z. N. (2014) "Montane resources exploitation and the emergence of gender
 333 issues in Santa economy of the Western Bamboutos Highlands, Cameroon"
 334 in *International Journal of Geography and Régional Planning Research*,
 335 *European Centre for Research Training and Development UK*
 336 (www.eajournals.org) Vol. 1, No. 1, pp. 1-12,
 337 Fontem, D.A. (1993). Severity of tomato diseases in Cameroon. *Tropicultura*,
 338 11(3):87-90.
 339 Friesen, L.G. (1998). Toward a market economy: fruit and vegetable production by
 340 the peasants of New Russia, 1850-1900', *Canadian Slavonic Papers*.
 341 40(1/2):27- 4218,
 342 Kaguongo, W. P., Gildemacher, P., Demo, P., Wagoire, W., Kinyae, P., Andra, de
 343 J., Forbes, G., Fuglie, K. and Thiele, G (2008). Farmer practices
 344 and adoption of improved potato varieties in Kenya and
 345 Uganda. International Potato Center (CIP) Lima, Peru. Social
 346 Sciences Working Paper. Pp.5-85.
 347 Norida, M. and John, M. (2005). Insecticide use in cabbage pest management in the
 348 Cameron Highlands, Malaysia. *Crop Protection*. 24: 31-39
 349 Nsobinyui, D., Ntonifor, N.N. and Fokam, B.E.(2017). Seasonal changes in field-
 350 to-store insect pests of maize and implications for their control in South
 351 Western Cameroon. *Journal of Agriculture and Ecology Research*
 352 International, 12(3)1-17.
 353 Ntonifor, N. N., Divine, N. S. N., Eric B. F. and Lum A. F. (2013). Developing an
 354 Integrated Management Approach for the Fruit Fly *Dacus*
 355 *punctatifrons* on Tomatoes. *American Journal of Experimental Agriculture*.
 356 3(3): 470-481,

Comment [A16]: (2016). Perception

Comment [A17]: No italic

Comment [A18]: No italic

357 Petzoldt, C. and Seaman, A. (2010). Climate change effects on insects and pathogens.
 358 Climate change and agriculture: Promoting practical and profitable
 359 responses.

360 Radcliffe, E.B., and Ragsdale, D.W. (2002). Aphid-transmitted potato viruses: the
 361 importance of understanding vector biology. *American Journal of Potato*
 362 *Research*. **79**: 353–386

363 Sait, E. (2003). Economic analysis of pesticide use on processing tomato growing:
 364 A case study for Turkey. *Crop Protection*. **25**: 534–541

365 Sonchieu J, Ngassoum M. B, Nantia A. E, Laxman P. S. (2017): Pesticide
 366 Applications on Some Vegetables Cultivated and Health Implications in
 367 Santa, North West-Cameroon. *International Journal of Agriculture &*
 368 *Environmental Science (SSRG – IJAES) – Volume 4 Issue 2*

369 Talekar, N. S. and Shelton, A. M. (1993). Biology, ecology and management of the
 370 Diamondback moth. *Annual Review of Entomology* **38**: 275-301.

371 Wilfred A. Abia, Conalius E. Shum, Richard N. Fomboh, Epole N. Ntungwe and
 372 Markjovert T. Ageh (2016): Agriculture in Cameroon: Proposed
 373 Strategies to Sustain Productivity. *International Journal for Research in*
 374 *Agricultural Research*