

# Asymptomatic Bacteriuria and Candida colonization among Pregnant Women in a District Hospital in Eastern Uganda.

## Abstract

**Background:** Urinary tract infection (UTI) is the most frequently encountered infection worldwide besides those that are of intestinal origin. It is the most common reason for which antimicrobials are prescribed in pregnancy Worldwide. A greater proportion of females is affected, to as high as fourteen times more than their male counterparts. This study aimed to determine the prevalence of asymptomatic bacteriuria, Candida colonization and antimicrobial susceptibility patterns among pregnant women attending antenatal in a District Hospital in Eastern Uganda.

**Materials and methods:** A descriptive cross sectional study was conducted in which pregnant mothers who had come for routine antenatal care were counseled and their consents obtained before taking off urine samples for laboratory diagnosis. For those samples found to have pus cells, culture and sensitivity test was done in Busitema University Microbiology Laboratory to identify the organisms and determine susceptibility to particular antibiotics and antifungal agents.

**Results:** Both gram positive and gram negative bacteria were isolated with gram positives (41.7%) showing higher frequency than the gram negatives (29.6%) while the remaining percentage was accounted for by *Candida spp* (28.8%). Gram negative isolates were more sensitive to meropenem (100%), and ciprofloxacin (93.8%) but less sensitive to trimethoprim/sulphurmethoxazole (20%), Cefazidime (7%), and Cefepem (6%). Gram positive isolates were more sensitive to vancomycin (100%), meropenem (87%) and linezolid (88.1%) but less sensitive to Cefotaxime (31%) and Trimethoprim/sulphurmethoxazole (14%). All bacteria isolated in this study were resistant to three or more classes of antibiotics therefore referred to as multi-drug resistant (MDR). All Candida isolates were susceptible to Econazole and Nystatin whereas all isolates were resistant to Griseofulvin. High susceptibility to Clotrimazole (95%-100%) and Ketoconazole (81%-100%) was noted among all isolates whereas high resistance to Amphotericin B, Fluconazole and Voriconazole was noted.

**Conclusion:** Asymptomatic bacteriuria is highly prevalent in Butaleja district with many of the bacteria isolated exhibiting resistance to the commonly used antibiotics. Antifungal resistance was common in this study. *In vitro* antifungal susceptibility testing is important in

**Comment [u1]:** The abstract is generally too long. Fine a way to cut off not too relevant information

**Comment [u2]:** Too much information in one abstract.

**Comment [u3]:** Delete. Instead mention the methods used in carrying out the culture and sensitivity

**Comment [u4]:** Italicize. Put a 'period' after "spp"

**Comment [u5]:** space

**Comment [u6]:** This statement is hanging. Recast please.

35 guiding therapeutic decision making, as an aid in drug development studies, and as a means  
36 of tracking the development of antifungal resistance in epidemiological studies.

Comment [u7]: You may consider taking this to your review of literature.

## 37 Introduction

### 38 Background

39 Urinary tract infection (UTI) is the most frequently encountered infection worldwide besides  
40 those that are of intestinal origin (1). Globally, it has been estimated that about 150 million  
41 people are diagnosed with a UTI per year (2). Urinary Tract infections can be classified on  
42 the basis of presentation that can be lower UTI (urethra and urinary bladder affected) or  
43 upper urinary tract (kidneys affected) or whether a pregnant woman presents with  
44 (symptomatic) or without symptoms (asymptomatic). Although UTIs affect individuals of all  
45 ages, the females are fourteen times likely to be affected than men and 50-60% will suffer an  
46 episode of UTI once in their life time since incidence increases by 10% for every decade of  
47 life above 20yrs of age and this is due to their shorter urethra and close proximity of the anus  
48 to the genital area. Amongst women too, some groups are more susceptible than others such  
49 as the sexually active, elderly and pregnant women. These infections affect individuals of all  
50 age groups but show greater occurrence in particular groups like women that are sexually  
51 active, the pregnant and the elderly. A greater proportion of females is affected, to as high as  
52 fourteen times more than their male counterparts (3) and 50-60% of women will suffer from a  
53 UTI at least once in their lifetime (4).

Comment [u8]: years

54 UTIs are more prevalent in pregnancy due to the physiological changes of pregnancy. An  
55 estimated 25% of the pregnant women develop UTI in developing countries and it is the most  
56 common cause of admission in obstetric wards. This figure is much lower in developed  
57 countries (2-10%) (5).

58 The prevalence of UTI among pregnant women in African countries revolves around 14% as  
59 shown by researches carried out in Sudan (14.0%), Tanzania (14.6%), and Ethiopia (11.6%).  
60 These figures do not regard the women's age, parity and gestational age. However, studies  
61 point out *E. coli* as the commonest isolated organism with multi resistance toward different  
62 antibiotics (6).

63 Asymptomatic bacteriuria is common among ante-natal mothers in Uganda (7).  
64 Asymptomatic bacteriuria in pregnancy is more likely to cause adverse effects that could lead  
65 to maternal and perinatal morbidity and mortality. Since screening and treatment has been  
66 shown to be beneficial for both maternal and fetal wellbeing especially where prevalence

67 exceeds 2%, treatment reduces the prevalence of pyelonephritis by 75%, it is important to  
68 know the dominant uropathogens and the sensitivity patterns. This study aimed to determine  
69 the prevalence of asymptomatic Bacteriuria, Candida colonization and antimicrobial  
70 susceptibility patterns among Pregnant Women in a District Hospital in Eastern Uganda

Comment [u9]: the prevalence

Comment [u10]: Italicize

## 71 MATERIALS AND METHODS

### 72 Study design.

73 A descriptive cross sectional study was conducted in which asymptomatic pregnant mothers  
74 who had come for routine antenatal care were counseled and their consents obtained before  
75 taking off urine samples for laboratory diagnosis. For those samples found to have pus cells,  
76 culture and sensitivity test was done in the microbiology laboratory of Busitema University  
77 Faculty of Health Sciences to identify the organisms and determine susceptibility to particular  
78 antibiotics.

### 79 Study area

80 This study was conducted at Busolwe Hospital; the district hospital of Butaleja district, which  
81 serves the districts of Butaleja, Namutumba, Budaka and some people from Tororo.  
82 Busolwe hospital is one of the Community Based Education, Research and Services  
83 (COBERS) sites of Busitema University Faculty of Health Sciences.


84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99

100

101

**Fig. 1: Map of Uganda showing the distribution of COBERS sites in Eastern Uganda**

**Comment [u11]:** Figure titles are usually placed below the figure itself.



102

103

### Study population

The study involved testing asymptomatic mothers attending antenatal services in Busolwe Hospital maternity department during the month of April, 2017. The population in this area is predominantly Banyole (85%), a Bantu tribe whose language is called Lunyole. Other tribes in the district (15%) include the Japadhola, Bagisu, Basoga, Iteso, Karimojong and Bagwere; making a total population of 245,873 as per the census report 2014.

109

110

## 111 Sampling techniques and sample collection

112 The mothers were sampled consecutively as they came to the antenatal clinic to give equal  
113 chance for participation in the study. The sample collection procedure was explained to all  
114 mothers and written informed consent was obtained from the mothers who accepted to  
115 participate in the study. The mother was availed with a labeled urine sample container, clean  
116 cotton, and water after explaining the procedure for sample collection to collect mid-stream  
117 urine.

## 118 Laboratory procedures

119 Urine dipsticks were used to screen for presence of pus cells in urine indicated by leucocyte  
120 positivity. Samples that were positive for leucocytes were transported to Busitema University  
121 microbiology laboratory for culture and sensitivity testing. Quantitative urine cultures were  
122 done in the laboratory, that is; Urine was gently shaken, tipped to slant and with a sterile  
123 pipette, 100µml of urine was transferred onto the CLED, MacConkey agar and chocolate  
124 agar. The urine was spread evenly across the plates with a sterile glass rod and allowed to  
125 soak in. The plates were incubated at 37°C for 18-24hrs and read for growth. Plates which  
126 showed no growth at 24hrs were incubated for another 24hrs to allow for detection of slow  
127 growers. The plates that were negative after 48 hours were reported as no significant growth.  
128 Growth of more than three colony types typically indicated contamination. Bacterial counts  
129 of  $\geq 10^4$  Cfu/ml were indicative of an infection and counts below  $10^4$  Cfu/ml was taken to  
130 indicate contamination and further tests would not be performed unless the organisms were  
131 Enterobacteriaceae.

## 132 Morphological and biochemical identification of the bacteria

133 The bacteria was identified by colony morphology on culture plates and the microscopic  
134 appearance on Gram stain. Biochemical identification of bacterial isolates was done using  
135 standard methods (8-10) Briefly, the tests employed were catalase, free and bound coagulase,  
136 DNAase, Mannitol Salt Agar (Oxoid), oxidase, motility test using motility indole urea  
137 medium, reactions on triple sugar iron agar (TSI), urease, nitrate reduction, indole, methyl red  
138 (MR), Voges Proskauer (VP), citrate utilization, lysine decarboxylase, and sugar fermentation  
139 tests.

140

141 Drug susceptibility testing (DST) was performed using the Kirby-Bauer disc diffusion method  
142 on Mueller Hinton Agar (MHA) (Oxoid, Hampshire, United Kingdom) plates as

Comment [u12]: 'Isolation of Bacteria from Samples' would be a better title.

Comment [u13]: µl or?

Comment [u14]: Space: 24 hrs

Comment [u15]: Observed or examined

Comment [u16]: Please be consistent: hrs or hours

Comment [u17]: Contamination from what source?

Comment [u18]: CFU/mL

Comment [u19]: Double space

Comment [u20]: space

Comment [u21]: space

Comment [u22]: spelling

143 recommended by the clinical laboratory standards institute (11). Bacterial colonies were  
144 emulsified into sterile saline and the turbidity of the suspension adjusted to the 0.5 McFarland  
145 standard.

146  
147 *Candida* species were identified by Gram stain, colony morphology on Sabouraud Dextrose  
148 Agar (SDA) and color changes on Candida chromogenic Agar. Susceptibility to the  
149 antifungals was done on SDA and interpreted using the CLSI guidelines (11).

Comment [u23]: Italicize

Comment [u24]: Anti-fungal

Comment [u25]: What is the full meaning of CLSI

## 150 Data management and analysis.

151 Findings were entered in excel, exported to STATA v14 for analysis and presented in form of  
152 tables and figures.

Comment [u26]: Were entered

Comment [u27]: Tables and Figures

## 153 Ethical Considerations.

154 Ethical clearance was obtained from Mbale Regional Referral Hospital Research and Ethics  
155 committee, the District Health Office, Medical superintendent and the administration of  
156 Busolwe Hospital gave administrative clearance before commencement of the study.  
157 Participating mothers consented to participate in the study and research information was  
158 accessible to only the research team. Serial numbers were used instead of names to maintain  
159 confidentiality.

## 160 RESULTS


### 161 Demographics characteristics and the prevalence of asymptomatic bacteriuria

162 The prevalence of asymptomatic bacteriuria among pregnant women in Butaleja district was  
163 32.8%. A majority of the pregnant women with asymptomatic bacteriuria were between the  
164 ages of 18-30 (77.1%) followed by those above 35 years (16.7%) and the least were those  
165 below 18 years (6.2%). Asymptomatic bacteriuria was most prevalent among the prime  
166 gravidae (44.8%) and least prevalent among the grand multi-gravidae (14.6%). Busolwe  
167 accounted for the largest number of participants in this study with 67 participants, 50 of  
168 whom showed growths on culture. The rest of the participants were from a number other  
169 parishes (Fig.2).

Comment [u28]: recast

170  
171  
172  
173  
174

**Fig.2: Distribution of asymptomatic bacteriuria by place of residence**



**Comment [u29]:** figures are labeled below the actual figure

### Microbial etiology of asymptomatic bacteriuria

Both gram positive and gram negative bacteria were isolated with gram positives (41.7%) showing higher frequency than the gram negatives (29.6%) while the remaining percentage was accounted for by *Candida* spp (28.8%). *Staphylococcus aureus* (35.6%), was the most commonly isolated among the gram positive cocci whereas *Escherichia coli* (12.9%) was the most isolated gram negative bacilli.

**Comment [u30]:** italicize

**Comment [u31]:** spp.

**Comment [u32]:** space

### Sensitivity to antibiotics

Antibiotic susceptibility to the commonly used antibiotics used in our setting was carried out using the Kirby-Bauer disc diffusion method on Mueller-Hinton agar. The antibiotic discs used included; Amoxicillin/clavulanic acid (20/10µg), Ciprofloxacin (5µg), Gentamicin (10µg), Meropenem (10µg), Chloramphenicol (30µg), Trimethoprim/Sulphurmethoxazole (1.25/23.75µg), Cefotaxime (30µg), Cefepime (30µg), Ceftazidime (30µg), Clindamycin (2µg), Vancomycin (30µg), Nitrofurantoin (300µg) and Linezolid (30µg).

**Comment [u33]:** Antibiotic Sensitivity Pattern of Isolates

**Comment [u34]:** spelling

Gram negative isolates were more sensitive to meropenem (100%), and ciprofloxacin (93.8%) but less sensitive to trimethoprim/sulphurmethoxazole (20%), Ceftazidime (7%), and Cefepem (6%). Gram positive isolates were more sensitive to vancomycin 100%, meropenem 87% and linezolid 88.1% but less sensitive to Cefotaxime (31%) and Trimethoprim/sulphurmethoxazole (14%). All bacteria isolated in this study were resistant to

**Comment [u35]:** This is better mentioned in the materials and methods section of the article.

**Comment [u36]:** space

196 three or more classes of antibiotics therefore referred to as multi-drug resistant (MDR) (Table  
197 1).

### 198 Sensitivity to the antifungals

199 Commonly used antifungal agents in our setting were used to determine the antifungal  
200 susceptibility pattern using the Kirby-Bour disc diffusion method on Sabourauds Dextrose  
201 Agar (SDA). The antifungal agents used were Voriconazole (1µg), Ketoconazole (10µg),  
202 Amphotericin B (20u), Griseofulvin (10µg), Itraconazole (10µg), Fluconazole (25µg), Clotrimazole  
203 (10µg), Econazole (50µg), and Nystatin (100µg).

204  
205 *Candida albicans*, *Candida parasilosis* and *Candida glabrata* isolated were subjected to  
206 antifungal susceptibility to the commonly used antifungals. All *Candida* isolates were  
207 susceptible to Econazole and Nystatin whereas all isolates were resistant to Griseofulvin.  
208 High susceptibility to Clotrimazole (95%-100%) and Ketoconazole (81%-100%) was noted  
209 among all isolates whereas high resistance to Amphotericin B, Fluconazole and Voriconazole  
210 was noted.

Comment [u37]: spelling



Type of isolate	Antimicrobial resistance profile of the isolates, frequency (%)												
Gram negative rods	Profile	AMC	CIP	C	CN	MEM	SXT	CTX	FEP	CAZ	MDR		
<i>E. coli</i> (n=17)	S	0(0)	14(82)	5(29)	8(47)	16(94)	3(18)	12(71)	0(0)	1(6)			
	R	17(100)	3(18)	12(71)	9(52)	1(6)	14(82)	5(29)	17(100)	16(94)	17(100)		
<i>Klebsiella spp</i> (n=10)	S	0(0)	8(80)	1(10)	6(60)	10(100)	2(20)	5(50)	1(10)	0(0.0)			
	R	10(100)	2(20)	9(90)	4(40)	0(0)	8(80)	5(50)	9(90)	10(100)	10(100)		
<i>Enterobacter spp</i> (n=12)	S	5(42)	9(75)	8(67)	5(42)	12(100)	4(33)	5(42)	1(8)	2(17)			
	R	7(58)	3(25)	4(33)	7(58)	0(0)	6(50)	7(58)	11(92)	10(83)	12(100)		
Type of isolate	Antimicrobial resistance profile of the isolates, frequency (%)												
Gram Positive cocci	Profile	CIP	C	CN	DA	MEM	SXT	CTX	P	VA	F	LNZ	MDR
<i>S. aureus</i> (n=47)	S	15(32)	29(62)	39(83)	40(85)	45(96)	13(28)	17(36)	25(53)	47(100)	45(96)	45(96)	
	R	32(68)	18(38)	8(17)	7(15)	2(4)	34(72)	30(64)	22(47)	0(0)	2(4)	2(4)	7(100)
<i>Enterococcus spp</i> (n=8)	S	7(88)	3(38)	2(25)	3(38)	8(100)	0(0)	2(25)	7(88)	8(100)	7(88)	7(88)	
	R	1(13)	5(63)	6(75)	5(63)	0(0)	8(100)	6(75)	1(13)	0(0)	1(13)	1(13)	8(100)

Comment [u38]: line numbering missing from here down

**Table 1: Antimicrobial susceptibility pattern of bacterial isolates from asymptomatic bacteriuria among pregnant women in Butaleja District.**  
**Key:**S- Sensitive, R- Resistant, AMC- Amoxicillin/Clavulanic acid, CIP-Ciprofloxacin, C-Chloramphenical, CN-Gentamycin, DA-Clindamycin, MEP-Meropenem, SXT- Sulfamethoxazole/Trimethoprim, CTX-Cefotaxime, VA-Vancomycin, F-Nitrofurantoin, LNZ- Linezolid, P-Penicillin, FEP-Cefepem, CAZ-Ceftazidime.

Comment [u39]: Tables have their labels at the top of the respective tables

Isolate	Antifungal susceptibility pattern, frequency (%)									
	Profile	VOR	KTC	AMB	GRS	ITR	FLU	CLT	ECO	NY
C. albicans	S	13(62)	17(81)	13(63)	0(0)	15(71)	13(62)	20(95)	21(100)	21(100)
	R	8(38)	4(19)	8(38)	21(100)	6(29)	8(38)	1(5)	0(0)	0(0)
C. Parasilosis	S	4(100)	4(100)	1(25)	0(0)	1(25)	2(50)	4(100)	4(100)	4(100)
	R	0(0)	0(0)	3(75)	4(100)	3(75)	2(50)	0(0)	0(0)	0(0)
C. glabrata	S	0(0)	1(100)	1(100)	0(0)	1(100)	1(100)	1(100)	1(100)	1(100)
	R	1(100)	0(0)	0(0)	1(100)	0(0)	0(0)	0(0)	0(0)	0(0)

**Table. 2: Antifungal susceptibility patterns of Candida Species Isolated**

**Key:** VOR-Voriconazole, KTC-Ketoconazole, AMB-Amphotericin B, GRS-Griseofulvin, ITR-Itraconazole, FLU-Fluconazole, CLT-clotrimazole, ECO-Econazole, NY-Nystatin

## DISCUSSION

### Prevalence of asymptomatic bacteriuria in Butaleja District

The prevalence of asymptomatic bacteriuria among pregnant women in Butaleja district was 32.8%. This is higher than the 13.3% reported in a similar study in Uganda (7). It's also higher than the specified split prevalence for symptomatic and asymptomatic bacteriuria recorded in Mwanza, Tanzania at 17.9% and 13.0% respectively (12), 21.5% in Nairobi, Kenya (13) and elsewhere (14, 15). There has generally been a varying prevalence of asymptomatic bacteriuria across Africa though the high prevalence in this study may be attributed to the observed low socioeconomic status in Busolwe district. Other studies elsewhere have indicated that low social economic status is associated with high prevalence of asymptomatic bacteriuria (16). Some studies in Africa have similarly documented a high prevalence of asymptomatic bacteriuria among pregnant women (17) with Benin City recording a prevalence of over 50% (18).

Comment [u40]: Not formal for an article

Comment [u41]: Full stop after the sentence

A majority of the pregnant women with asymptomatic bacteriuria were between the ages of 18-30 (77.1%) and the least were those below 18 years (6.2%). Similar studies have associated age group 18-34 with a high prevalence of asymptomatic bacteriuria (14). The association of the above age range with asymptomatic bacteriuria may be due to high rate of sexual activity expected in this age range. Some studies have however not associated the prevalence of asymptomatic bacteriuria with age (13, 19).

Asymptomatic bacteriuria was most prevalent among the prime gravidae (44.8%) and least prevalent among the grand multi-gravidae (14.6%), though there was no association between gravidity and asymptomatic bacteriuria. Another study in Nigeria did not find a significant association between gravidity and asymptomatic bacteriuria (19). Busolwe accounted for the largest number of participants in this study with 67 participants, 50 of whom showed growths on culture. The rest of the participants were from a number other parishes. The large number of participant from Busolwe sub-county could be due to the location of Busolwe hospital and this paints a picture that there is a lot of undiagnosed asymptomatic bacteriuria from communities far away from health facilities.

Comment [u42]: See line 166-169. Repetition

Comment [u43]: Recheck this sentence and fix it were necessary

Comment [u44]:

### Microbial etiology of symptomatic bacteriuria in Butaleja District.

*Staphylococcus aureus* (35.6%), was the most commonly isolated among the gram positive cocci whereas *Escherichia coli* (12.9%) was the most isolated gram negative bacilli. Whereas many studies in Uganda and elsewhere agree with the predominance of *E. coli* in Urinary

Comment [u45]: Too common a word in this article.

tract infections, (7, 15, 16, 18, 20-23). Other studies have demonstrated different bacterial species as dominant uropathogen like *Klebsiella* (24). Similar studies have demonstrated the rise in numbers of gram positive cocci in urinary tract infections for example, a study in North East Ethiopia, Gram-positive isolates were more prevalent (n=37/58: 63.8%) than Gram-negative bacteria (n=21/58; 36.2%) and the most commonly isolated bacteria were *S. aureus* (n=18; 31%) and *E. coli* (n=18; 31%) (25). Other studies have also demonstrated *S. aureus* as the most commonly isolated uropathogen ((19, 26). *Enterococcus spp* was the commonly isolated organism in a study in Ghana, followed by *Proteus mirabilis* and then *Escherichia coli* (14).

Comment [u46]: italicize

Comment [u47]: recast this sentence

Comment [u48]: double bracket

Comment [u49]: don't italicize

Co-infection with *Candida* was found in 28.8% of the asymptomatic bacteriuria cases with *Candida albicans* being the dominantly isolated yeast. The other *Candida* species isolated were *Candida parasilosis* and *Candida glabrata*. Candiduria has also been documented in coexistence with bacteriuria in other studies (16, 27). The co-existence of bacteriuria and candiduria may lead to works effects in pregnant women.

Comment [u50]: This is not clear

## Sensitivity of the antimicrobials

### Antibacterial susceptibility

While the range of bacteria causing asymptomatic bacteriuria is relatively constant, the susceptibility of the same to the antibiotics varies a lot by geographical location. Gram negative isolates in this study were more sensitive to meropenem, and ciprofloxacin with sensitivity patterns ranging from 75%-100%. There was no carbapenem resistance detected among the gram negative isolates. Similar patterns of sensitivity to the carbapenems have been noted by studies in neighboring Kenya (12, 13, 28, 29).

Comment [u51]: See comment U49

Comment [u52]: You should also state studies where their reported outcome differs from yours so as to drive home your point above.

Gram positive isolates were highly susceptible to Gentamicin, Vancomycin, Meropenem Nitrofurantoin and linezolid. Similar studies have reported high susceptibility to Nitrofurantoin which indicates that Nitrofurantoin is still a useful drug for management of bacteriuria (14, 18). Its high susceptibility may though be due to its less frequent use as compared to other antibiotics in the management of urinary tract infections in Uganda. High susceptibility to Gentamicin is worth celebrating because it is relatively cheap and can be afforded by the rural poor. On the other hand, drugs like Vancomycin, Meropenem and Linezolid which show high susceptibility in this study are very expensive and with invasive modes of administration.

Sulfamethoxazole-trimethoprim (SXT), Cefotaxime, Ceftazidime and Cefepem had the lowest sensitivities to both gram positive and gram negative isolates in our study. This correlates with findings that have documented high resistance to the first line antimicrobial drugs such as cotrimoxazole (13, 30). Our findings are also in agreement with reports from other regions of Uganda which showed that the commonly used antibiotics are non-effective (7). Cefotaxime is second line antimicrobial agent in the third generation of cephalosporins. A high resistance to cefotaxime, ceftazidime and cefepem in this study calls for an urgent need for surveillance of antimicrobial resistance among pregnant women with asymptomatic bacteriuria. The resistance to cotrimoxazole in our study may be attributed to its widespread over-the-counter use in our locality and its use for prophylaxis against opportunistic infections among people living with HIV. All bacteria isolated in this study were resistant to three or more classes of antibiotics therefore referred to as multi-drug resistant (MDR)

#### Antifungal susceptibility

The range of *Candida* species associated with candidiuria is relatively constant. In this study, *Candida albicans*, *Candida parasilosis* and *Candida glabrata* were isolated and subjected to antifungal susceptibility to the commonly used antifungals. A similar study in Mbarara isolated similar species of *Candida* (31). *In vitro* antifungal susceptibility testing now plays an increasingly important role in guiding therapeutic decision making, as an aid in drug development studies, and as a means of tracking the development of antifungal resistance in epidemiological studies (32). All *Candida* isolated in this study were susceptible to Econazole and Nystatin whereas all isolates were resistant to Griseofulvin. This compares with another study done in Mbarara, South-Western Uganda which showed 100% susceptibility of *Candida* isolates to Nystatin (31) and similar results were obtained in Argentina (33). High susceptibility to Clotrimazole (95%-100%) and Ketoconazole (81%-100%) was noted among all isolates whereas high resistance to Amphotericin B, Fluconazole and Voriconazole was noted. Our findings differ from the study in South-Western Uganda which showed good susceptibility to fluconazole.

Comment [u53]: Italicize

Comment [u54]: isolate

Comment [u55]: italicize

#### Conclusion

Asymptomatic bacteriuria is highly prevalent among pregnant women in Butaleja District with a number of bacteria exhibiting multi-drug resistance. The common etiological agents of asymptomatic bacteriuria in Butaleja district were *S. aureus*, and *E. coli* whereas *Enterococcus spp* was the list commonly isolated. The susceptibility patterns shown in this

Comment [u56]: italicize

study highlight the need for sensitivity studies before initiating treatment for a UTI, a challenge that is prevalent in the local health facilities like Busolwe Hospital. The prevalence of *Candida* colonization was also high with notable resistance to the commonly used antifungals.

Comment [u57]: Italicize

## Recommendations.

We recommend regular screening of pregnant women for urinary tract infections whether asymptomatic or not so that those with bacteriuria get timely treatment. We also recommend continuous surveillance for antimicrobial resistance in this community. Antifungal susceptibility for *Candida* isolated should be encouraged to promote rational use of antifungal agents.

## REFERENCES

1. August SL, and De Rosa, M.J., 2012. Evaluation of the prevalence of urinary tract infection in rural Panamanian women. *PloS one*. 2012;7(10):e47752. .
2. Prakash DS, R.S Distribution and antimicrobial susceptibility pattern of bacterial pathogens causing urinary tract infection in urban community of meerut city, India *ISRN microbiology*. 2013:749629.
3. Dielubanza EJ, & Schaeffer, A.J. Urinary Tract Infections in Women. *Medical Clinics of North America*. 2011;95(1):27–41.
4. Al-Shaikh AA-BG. Urinary Tract Infections in Women. *Sultan Qaboo University Medical Journal*. 2013;13(3):359–67.
5. Gilbert NM, O'Brien VP, Hultgren S, Macones G, Lewis WG, Lewis AL. Urinary tract infection as a preventable cause of pregnancy complications: opportunities, challenges, and a global call to action. *Global advances in health and medicine*. 2013;2(5):59-69.
6. Hamdan HZ, Ziad, A. H., Ali, S. K., & Adam, I. Epidemiology of urinary tract infections and antibiotics sensitivity among pregnant women at Khartoum North Hospital. *Annals of clinical microbiology and antimicrobials*. 2011;10(2).
7. Andabati GB, J. Microbial aetiology and sensitivity of asymptomatic bacteriuria among ante-natal mothers in Mulago hospital, Uganda. *African health sciences*. 2010;10(4):349–52.
8. Najjuka CF, Kateete, D.P, Kajumbula H.M, Joloba, M.L, Essack, S.Y. Antimicrobial susceptibility profiles of *Escherichia coli* and *Klebsiella pneumoniae* isolated from outpatients in urban and rural districts of Uganda. *BMC Res Notes*. 2016;9(235).
9. Iramiot J. Stanley. Freddie Bwanga, Herbert Itabangi, Martha Nakaye, Mwambi Bashir and Joel Bazira. Prevalence and Antibiotic Susceptibility Patterns of Clinical Isolates of Methicillin-Resistant *Staphylococcus aureus* in a Tertiary Care Hospital in Western Uganda. *British Microbiology Research Journal*. 2014;4(10):1168-77.
10. Iramiot Jacob Stanley. Henry Kajumbula., Joel Bazira., Catherine Kansiime., Innocent B. Rwego., Benon B. Asimwe. Multidrug resistance among *Escherichia coli* and *Klebsiella pneumoniae* carried in the gut of out-patients from pastoralist communities of Kasese district, Uganda. *PLoS ONE*. 2018;13(7).
11. Clinical and Laboratory Standards Institute. M100-S25 Performance Standards for Antimicrobial Susceptibility Testing. 2014.

12. Masinde. A, B. Gumodok, A Kilonzo, SE Mshana. Prevalence of urinary tract infection among pregnant women at Bugando Medical Centre, Mwanza, Tanzania. *Tanzania Journal of Health Research*. 2009;11(3).
13. Ayoyi AO, Kikuvu, G., Bii, C., & Kariuki, S. (2017). . Prevalence, aetiology and antibiotic sensitivity profile of asymptomatic bacteriuria isolates from pregnant women in selected antenatal clinic from Nairobi, Kenya. *The Pan African medical journal*. 2017;26(41).
14. Labi AK, Yawson, A. E., Ganyaglo, G. Y., & Newman, M. J. Prevalence and Associated Risk Factors of Asymptomatic Bacteriuria in Ante-Natal Clients in a Large Teaching Hospital in Ghana. *Ghana medical journal*. 2015; 49(3):154–8.
15. Mokube MN, Atashili, J., Halle-Ekane, G. E., Ikomey, G. M., & Ndumbe, P. M. Bacteriuria amongst pregnant women in the Buea Health District, Cameroon: prevalence, predictors, antibiotic susceptibility patterns and diagnosis. *PloS one*. 2013;8(8).
16. Oli A. Okafor C, Ibezim E., Akujiobi C., Onwunzp M. The prevalence and bacteriology of asymptomatic bacteriuria among antenatal patients in Nnamdi Azikiwe University Teaching Hospital Nnewi; South Eastern Nigeria. *Niger J Clin Pr*. 2010;13(4):409–12.
17. Imade PE IP, Eghafona NO, Enabulele OI, Ophori E. Asymptomatic bacteriuria among pregnant women. *North Am J Med Sci*. 2010;2:263-66.
18. Okonko IO, Donbraye-Emmanuel, L.A. Ijandipe, A. A. Ogun, A.O. Adedeji and A.O. Udeze Antibiotics Sensitivity and Resistance Patterns of Uropathogens to Nitrofurantoin and Nalidixic Acid in Pregnant Women with Urinary Tract Infections in Ibadan, Nigeria. *Middle-East Journal of Scientific Research*. 2009;4(2):105–9.
19. Onu FA AL, Ezeonu PO, Umeora OIJ, Ibekwe PC, Ajah MI. Profile and microbiological isolates of asymptomatic bacteriuria among pregnant women in Abakaliki, Nigeria. *Infect Drug Resist* 2015;8:231–5.
20. Beyene G, & Tsegaye, W. Bacterial Uropathogens in Urinary Tract Infection and Antibiotic Susceptibility Pattern in Jimma University Specialized Hospital, Southwest Ethiopia. . *Ethiopian Journal of Health Sciences*. 2011;21(2):141–6.
21. Çelen Ş. OA, Karayalçın R, Saygan S, Ünlü S, Polat B, et al. Asymptomatic Bacteriuria and Antibacterial Susceptibility Patterns in an Obstetric Population. *Isrn Obstet Gynecol*. 2010;1(4).
22. Hamdan ZH AH, Salah KA, Ishag A. Epidemiology of urinary tract infections antibiotic sensitivity among pregnant women at Khartoum North Hospital. *Ann Clin Microbiol Antimicrob* 2011;102.
23. Oli A. OC, Ibezim E., Akujiobi C., Onwunzp M. The prevalence and bacteriology of asymptomatic bacteriuria among antenatal patients in Nnamdi Azikiwe University Teaching Hospital Nnewi. South Eastern Nigeria *Niger J Clin Pr*. 2010;13(4):409–12.
24. Akoachere JFTK YS, Akum NH, Seraphine EN. Etiologic profile and antimicrobial susceptibility of community-acquired urinary tract infection in two Cameroonian towns. *BMC Res Notes*. 2012.
25. Ali IE, Gebrecherkos, T., Gizachew, M., & Menberu, M. A. Asymptomatic bacteriuria and antimicrobial susceptibility pattern of the isolates among pregnant women attending Dessie referral hospital, Northeast Ethiopia: A hospital-based cross-sectional study. *Turkish journal of urology*. 2018;44(3):251–60.
26. Amadi ES EO, Uneke CJ, Nwosu OK, Onyeagba RA, Ugbogu OC. Asymptomatic bacteriuria among pregnant women in Abakaliki, Ebonyi State Nigeria. *J Med Sci* 2007;7:698–700. 2007;7:698–700.
27. Alemu A, Moges F, Shiferaw Y, Tafess K, Kassu A, Anagaw B, et al. Bacterial profile and drug susceptibility pattern of urinary tract infection in pregnant women at

University of Gondar Teaching Hospital, Northwest Ethiopia. BMC Research Notes. 2012;5(1):197.

28. Rizvi Mea. Rising prevalence of antimicrobial resistance in urinary tract infections during pregnancy: necessity for exploring newer treatment options. Journal of laboratory physicians. 2011;3(2):98–103.

29. Kashef N, Djavid, G.E. & Shahbazi, S Antimicrobial susceptibility patterns of community-acquired uropathogens in Tehran, Iran. . The Journal of Infection in Developing Countries,. 2010; 4( 4):202–6.

30. Moyo SJ AS, Kasubi M, Maselle SY. Bacterial isolates and drug susceptibility patterns of urinary tract infection among pregnant women at Muhimbili National Hospital in Tanzania. Tanzan J Health Res. 2010;12(4):236–40.

31. Mukasa KJ, Herbert, I., Daniel, A., Sserunkuma, K. L., Joel, B., & Frederick, B Antifungal Susceptibility Patterns of Vulvovaginal Candida species among Women Attending Antenatal Clinic at Mbarara Regional Referral Hospital, South Western Uganda. British microbiology research journal. 2014;5 (4):322–31.

32. Pfaller MA, & Diekema, D. J Progress in antifungal susceptibility testing of Candida spp. by use of Clinical and Laboratory Standards Institute broth microdilution methods, 2010 to 2012. Journal of clinical microbiology. 2012;50 (9):2846–56.

33. García HM GS, Copolillo EF, Cora EM, Barata AD, Vay CA, et al. Prevalence of vaginal candidiasis in pregnant women. Identification of yeasts and susceptibility to antifungal agents. Revista Argentina de Microbiología. 2006;38(1):9-12.