


## Original Research Article

### **Maize response to leguminous biomass composted with phosphate rocks in the Northern zone of Tanzania**

#### **Abstract**

A field study was conducted in Babati District of Northern Tanzania to evaluate maize response to leguminous biomass composted with phosphate rocks (PRs). Three leguminous (*Crotalaria juncea*, *Lablab purpureus* and *Mucuna pruriens*) strips were cultivated in 2013/14 to produce a biomass which was harvested at flowering to early podding stage and air dried. Air-dry biomass was composted with PRs from Minjingu (medium reactive PR) and Panda Hill (low reactive PR). In 2014/2015 the field strips were used to evaluate maize response to different treatments in a split plot design. The strips served as main plots and each strip was divided into seven subplots which received different treatments at random. A medium term maize variety SC. 627 was used as a test crop. Average maize grain yields obtained from *crotalaria*, *lablab* and *mucuna* strips reached 5.3, 4.5 and 4.0 t ha<sup>-1</sup>, respectively and were statistically different (P=0.05). Application of Minjingu or Panda Hill PR alone failed to increase maize grain yield above the control while Minjingu PR applied with urea or composted with biomass increased maize grain yield by 2.40 and 1.58 t ha<sup>-1</sup>, respectively above the control. Application of Panda Hill PR with urea or composted with biomass increased grain yield by 1.20 and 1.06 t ha<sup>-1</sup>, respectively above the control. Average maize grain yield produced when cover crop biomass was removed but Minjingu/Panda Hill PR + urea was applied reached 5.62 t ha<sup>-1</sup> as compared with 5.15 t ha<sup>-1</sup> obtained following application of biomass-PR composts. The observed difference (0.47 t ha<sup>-1</sup>) was not statistically significant indicating that biomass composted with PR was as effective as the PR applied with urea.

26


27 Key words: *Crotalaria*, *Lablab*  *ucuna*, phosphate rocks, compost, maize yield

28

29 The present address of Mawazo Shitindi is Sokoine University of Agriculture, Department  
30 of Soil and Geological Sciences, P.O. Box 3008 Morogoro Tanzania.

31

## 32 1. Introduction

33 Maize (*Zea mays*) is Tanzania's most important staple food with an estimated annual  
34 per capita consumption of 113 kg, contributing about 60% of dietary calories [ and [2].  
35 According to [3], the crop also contributes about 50% of Tanzania's rural cash income.  
36 However, current production of maize in Tanzania is far below the national average yield  
37 potential of 4.8 t ha<sup>-1</sup>, fluctuating between 1.0 and 1.5 t ha<sup>-1</sup> [4]. Continuous maize production  
38 without or with limited fertilizer application coupled with crop residue removal have been  
39 reported as major factors for soil fertility decline and low crop yields [5; 6; 7]. Limited  
40 fertilizer use in most developing countries has been attributed to their high costs and limited  
41 availability [8; 9].

42 While food production per unit land is declining because of soil fertility deterioration,  
43 the population of Tanzania has more than tripled from 12.3 million to 44.9 million between  
44 1967 and 2012. Based on 2012 census projections, the population is expected to reach 47.42  
45 million people by the year 2016 [10]. This increase in the population will cause additional  
46 pressure on arable land because more than 70% of Tanzanians depend entirely on agriculture  
47 for their food and income. This calls for integrated soil fertility management programs based  
48 on locally available resources so as to improve soil fertility and reduce smallholders'  
49 dependence on imported industrial fertilizers.

50 Phosphate rock (PR) deposits located in Tanzania could serve as alternative source of  
51 phosphorus (P) for smallholders but (P) contained in the rocks is not readily available for  
52 plant uptake. Upon decomposition, plant biomass releases low- molecular-weight organic  
53 acids that may complex calcium and other metals in the rock to free P for plant uptake [11].  
54 Thus, composting the rocks with leguminous biomass may improve the availability of  
55 nitrogen (N) and P for plant uptake. The objective of the field experiment was to investigate  
56 carbon (C), N and P content of three common leguminous plants (*Crotalaria juncea*, *Lablab*  
57 *purpureus*, and *Mucuna pruriens*) used in Tanzania and their effects when composted with  
58 PRs on maize yield. The PRs used were those of Mijingu (a PR of medium reactivity) and  
59 Panda Hill (a PR of low reactivity).

60

## 61 **2. Materials and Methods**

### 62 **2.1 Site Description, Soil Characterization and Fertility Assessment**


63 This study was conducted at Wang'waray Farmers Training Center (F.T.C) located in  
64 Babati District of Manyara region in the Northern zone of Tanzania. The site is about 167 km  
65 from Arusha and 4.5 km to the South East of Babati town along the road to Mamire Ward.  
66 The center is at 1410 m above sea level on the foot hills of mount Kwaraa, and receives a  
67 bimodal rainfall with average precipitation around 700-900 mm year<sup>-1</sup>. However, as with  
68 other areas in Tanzania, rainfall distribution at Wang'waray F.T.C and Babati District as a  
69 whole has been altered by climate change to such an extent that the two seasons are now not  
70 very distinct and average precipitation is less than 700 mm year<sup>-1</sup>. Crop production is a major  
71 land use activity at Wang'waray F.T.C. dominated by maize-legume intercropping and  
72 rotation systems. Because soils at Wang'waray FTC were not characterized before, a profile  
73 was opened and described according to FAO guidelines [12]. Representative profile and  
74 surface (0-15 cm) soil samples were collected and shipped to the Soil and Geological

75 Sciences (SGS) laboratory at Sokoine University of Agriculture (SUA) in Morogoro for  
76 physical and chemical analyses (Table1). Based on morphological description of the site, and  
77 laboratory analyses performed on the profile samples, the soil was classified down to sub  
78 group as *Rhodic Eutrotox* using the USDA-NRCS Keys to soil taxonomy [13]. Analyses of  
79 representative surface (1-15cm) soil samples collected from the rest of the field were used for  
80 assessment of general fertility status of soils.

## 81 2.2 Leguminous Biomass Production

82 Following soil characterization, two portions of the field separated by a contour band were  
83 ploughed and harrowed. On each portion of the field, three strips of 5 m x 76 m each were  
84 established and randomly assigned to one of the three legume crops (two strips for each cover  
85 crop) as shown in Figure 1. *Mucuna pruriens* and *Lablab purpureus* were planted at 50 cm x  
86 30 cm spacing, while *Crotalaria juncea* was drilled at 50 cm inter row spacing. The first  
87 weeding was done two weeks after germination and there after weeding was repeated  
88 whenever weeds emerged to keep the competition for moisture and nutrients to a minimum.

89






Figure 1. Layout of the field for leguminous crop biomass production at Wang'waray F.T.C

### 2.3 Carbon, nitrogen and phosphorus contents of the biomass

At flowering - podding initiation stage, the biomass was cut above the ground, air dried by the species for later composting with Minjingu or Panda Hill PR. Before composting, the air-dry biomass was chopped into small pieces to increase surface area and thoroughly mixed. Subsamples were collected, oven dried at 55°C for 72 hours, and finely ground to < 0.5mm using a CT 193 Cyclotec™ Sample Mill [Foss Allé 1 Post box 260 DK-3400 Hillerød Denmark] for chemical analyses. Organic carbon (OC) was determined following the Walkely & Black procedure (Nelson and Sommers [14], while total N was determined following Kjeldahl procedures [15]. For the determination of P and sulfur (S) in the biomass, a 0.5 g sample < 0.5 mm was digested following the HNO<sub>3</sub> - H<sub>2</sub>O<sub>2</sub> wet digestion procedure using a 40 space Foss Tecator block digester. Phosphorus content of the digest was determined by a colorimetric procedure using ascorbic acid method [16], while S content was determined by a turbidity method [17].

## 109 **2.4 Phosphate Rock Collection, Processing, and Chemical Analysis**

110 Minjingu PR was collected from Minjingu Mines and Fertilizers Company in  
111 Manyara region while Panda Hill PR was obtained from a storage facility at SUA. Both PRs  
112 were ground to pass a 100-mesh sieve at the Geological Survey of Tanzania (GST) laboratory  
113 in Dodoma region. A representative sample was collected from each PR and shipped to the  
114 Southern and Eastern Africa Mineral Center (SEAMIC) laboratory in Dar es Salaam for X-  
115 ray fluorescence (XRF) analysis.

## 116 **2.5 Production of Biomass-PR Composts**

117 Previously chopped leguminous biomass (< 2 cm) and ground PRs (< 100 mesh) were  
118 composted by the pit method [18] with some modifications. In the modifications, the size of  
119 an individual pit was 2 m x 2 m x 1m; floor and walls of each pit were lined with a  
120 polyethylene plastic sheet to avoid leaching losses during decomposition. The biomass was  
121 composted with a PR in alternating layers (i.e. PR was applied over every layer of biomass)  
122 followed by a small amount of dried cattle manure to inoculate the biomass. The biomass:PR  
123 ratio varied from 12:1 to 18:1 based on the biomass size and N contents. Following  
124 inoculation, water was applied to bring the moisture content of the compost mixture to about  
125 60%.

126

127 Three PVC aeration pipes were inserted into each compost mix at regular intervals  
128 and the material was covered with polyethylene plastic sheets to protect it from rain water  
129 and undesirable/ foreign materials. The compost material in each pit was turned into a  
130 different pit every 30 days for 120 days to allow optimum decomposition and water was  
131 sprinkled at every turn to maintain the moisture at 60%. After the last turn, representative  
132 samples were collected from each pit for laboratory analysis and all composts were air dried  
133 to around 20% moisture content and stored for later use as source of N and P for maize.

134 Representative samples taken from each pit were shipped to the SUA-SGS laboratory for  
135 chemical analysis. In the laboratory, representative compost samples were dried and ground  
136 to pass through 0.5 mm for total N, P and SO<sub>4</sub>-S analysis as previously described.

137

## 138 **2.6 Evaluation of Maize Response to Treatments**

139 The field strips previously used for cover crop biomass production were used in the  
140 next season to evaluate maize response to newly imposed treatments. The experiment was  
141 designed as a split plot arranged in a randomized complete block design (RCBD). The field  
142 was divided into four blocks where half of each strip initially used to produce the crop  
143 biomass was used as a main plot within a block and each main plot was divided into seven  
144 sub plots (16 m<sup>2</sup>) which received randomly assigned treatments (Figure 2).

145

146

147

148

149

150

151

152

153


154

155

156

157

158



159  
160

Figure 2. Layout of maize field experiment at Wang'waray FTC

161 Letters represent leguminous species (M = Mucuna, L = Lablab, C = Crotalaria) while  
 162 numbers  
 163 (1 - 7) represent treatments imposed on experimental units.  
 164 Seven treatments were evaluated on each main plot. These include a common control  
 165 where maize was grown without external inputs after removal of the crop biomass (1),



166 Minjingu PR alone applied (2), Minjingu PR + urea (3), composted Minjingu PR + biomass  
167 (4). Panda Hill PR alone (5), Panda Hill PR + urea (6), and composted Panda hill PR +  
168 biomass (7). Thus, treatment combinations were identified as C1 to C7, L1 to L7, and M1 to  
169 M7 where C, L, and M stand for crotalaria, lablab, and mucuna strip, respectively.

170  
171 The composts were applied at a rate corresponding to 112 kg N ha<sup>-1</sup> recommended for  
172 maize in the Northern Zone [19]. The PRs were applied at 45kg P ha<sup>-1</sup> with or without urea  
173 while urea was applied at 112kg N ha<sup>-1</sup> (split application at planting and two weeks following  
174 germination) on selected plots based on treatment scheme. A medium term hybrid maize  
175 variety (SC.627) was planted at 90 x 30 cm spacing (five rows per plot). At tasselling stage,  
176 nine representative ear leaf samples were collected from each plot for nutrient analysis. At  
177 maturity stage, maize ears of the three inner rows in each plot were harvested for yield  
178 determination. Maize grain yield was reported at 13% moisture content, while maize stover  
179 yield from the three inner rows of each plot was reported on oven-dry basis. The data  
180 collected were subjected to analysis of variance (ANOVA) using a mixed procedure of SAS  
181 software version 9.4 (SAS Instit. Inc. Cary, NC) and the means were separated at P = .05 by  
182 Tuckey-Kramer procedure.

183

### 184 **3. Results and Discussion**

#### 185 **3.1 Fertility status of soil at Wang'waray FTC**

186 Selected physical-chemical analyses of soil at Wang'waray FTC were as presented in  
187 table 1. The soil had a medium pH value suitable for production of most crops with a very  
188 low electrical conductivity indicating that there were no limitations for crop production due  
189 to salt accumulation.

190

191 Table 1. Selected chemical properties of surface (0 -15 cm) soil samples at Wang'waray  
192 F.T.C

Soil property	Mean <sup>†</sup>	Rating	Reference
pH – H <sub>2</sub> O	6.88	Medium	[20]
EC (MScm <sup>-1</sup> )	0.05	Very low	[20]
Organic Carbon (g kg <sup>-1</sup> )	14.3	Low	[20]
Total N (g kg <sup>-1</sup> )	1.03	Low	[21]
Bray 1 P (mg kg <sup>-1</sup> )	5.54	Low	[20]
SO <sub>4</sub> – S (mg kg <sup>-1</sup> )	9.38	High	[22]
Exch. Ca (Cmol kg <sup>-1</sup> )	7.40	High	[20]
Exch. Mg (Cmol kg <sup>-1</sup> )	2.96	High	[20]
Exch. K (Cmol kg <sup>-1</sup> )	3.28	High	[20]
Exch. Na (Cmol kg <sup>-1</sup> )	0.27	Low	[20]
PBS (%)	70.9	High	[20]
DTPA Extract. Cu (mg kg <sup>-1</sup> )	3.6	High	[20]
DTPA Extract. Zn (mg kg <sup>-1</sup> )	0.5	Low/medium	[20]
DTPA Extract. Mn (mg kg <sup>-1</sup> )	116.5	High	[20]
DTPA Extract. Fe (mg kg <sup>-1</sup> )	22.0	High	[21]
Sand (g kg <sup>-1</sup> )	643		
Silt (g kg <sup>-1</sup> )	87		
Clay (g kg <sup>-1</sup> )	270		
Textural class	Sandy Clay Loam		[23]

193

194 †, Each reading is an average of six representative surface soil samples

195

196 Levels of extractable S, exchangeable bases and DTPA extractable Fe, Cu and Mn were all  
 197 high but the levels of organic carbon, total N, Bray-1 extractable P were low, and therefore  
 198 limiting. The low levels of organic carbon, N, and P have been reported in highly weathered  
 199 tropical soils like those of Babati [24].

### 200 3.2 Carbon, Nitrogen, and Phosphorus Content of Leguminous Biomass Used

201 Carbon contents of the leguminous biomass used varied significantly (P=.05) while  
 202 there were no significant differences in the P contents (P = .05).

203

204

205 Table 2. Chemical composition of the leguminous biomass used  
 206

Crop species	C	N	P	C:N	C:P	N:P
	-----%-----					
<i>Crotalaria juncea</i>	48.7 a	2.44 a	0.37 a	20.1	136	6.74
<i>Lablab purpureus</i>	41.6 b	2.30 a	0.34 a	18.1	124	6.76
<i>Mucuna pruriens</i>	44.5 b	2.00 b	0.36 a	22.3	122	5.75
LSD	6.30	0.14	0.05	-	-	-

207  
 208 †, Values in the same column followed by the same letter are similar (P = .05)  
 209

210 Chemical composition of plant species grown for compost production is an important  
 211 factor to take into account because it has effect on the rate at which plant material is acted  
 212 upon by decomposers to release nutrients in plant available forms. On average, OC contents  
 213 of the biomass used were 48.7%, 41.6% and 44.5% for crotalaria, lablab, and mucuna  
 214 biomass, respectively. On the other hand, total N content of the biomass used were 2.4%,  
 215 2.3% and 2.0%, while the C:N ratios were 20.1, 18.1, and 22.3 for crotalaria, lablab and  
 216 mucuna biomass, respectively. The total N values determined in all leguminous crop biomass  
 217 were below 3.0 % which is considered as critical value for sufficiency in most legume plants.  
 218 However, the tropical soil biology and fertility program data base cited by [25] specified total  
 219 N in the range of 1.6-5.7%, 1.7-6.3% and 1.4-6.5%, as normal for of crotalaria, lablab and  
 220 mucuna biomass respectively when harvested at flowering stage depending on soil properties  
 221 and environmental condition of a given area. The data base also specified the C:N ratios in  
 222 the range of 8.0-32.1, 7.4-29.1, and 9.8-30.8 for crotalaria, lablab and mucuna biomass,  
 223 respectively when harvested at flowering stage. Based on these specifications, the OC, total N  
 224 and C:N ratios were all within the normal range for the crop species used. Furthermore, the

225 C:N ratios of the biomass used were below 30:1 which is the recommended highest value  
226 acceptable for an effective decomposition and mineralization of plant biomass [26].

227


### 228 **3.4 Selected Chemical Properties of PRs Used**

229 Selected chemical properties of PRs are as presented in Figures 3, 4, and 5. Solubility  
230 of PR depends largely on soil moisture status, soil pH, exchangeable Ca, available P and P  
231 adsorption capacity of a soil [27]. Composition of PRs also affects relationships between  
232 concentrations of their dissolution products and their sinks in the soil hence affecting  
233 dissolution reactions in equilibrium. Apart from affecting the nature and rates of dissolution  
234 reactions, chemical constituents of the PRs also play different roles in plant nutrition hence  
235 contributing to variations in crop responses following application of PRs of different  
236 chemical compositions [28].

237

238 Minjingu PR as shown in figure 3, has higher concentrations of  $P_2O_5$ , CaO,  $MgO_2$ ,  
239  $K_2O$  and NaO than Panda Hill PR. These differences are functional of geological origin of the  
240 rocks and dictate the relative availability of P, Ca, Mg, K and Na from the two PRs. Apart  
241 from Na which is only essential in some plants where it has been reported to take over the  
242 function of K when the latter is not readily available; P, Ca, K and S are essential elements  
243 for all plants and therefore contribute to the fertilizer value of Minjingu PR. Furthermore,  
244 with the exception of Ca, most of the elements found in higher concentrations in Minjingu PR  
245 have low affinity for P. This explains the reason for higher reactivity and therefore positive  
246 crop response reported following applications of Minjingu PR than that of Panda Hill PR [29;  
247 30; 31; 32].


248



249

250 Figure 3. Concentrations of P<sub>2</sub>O<sub>5</sub>, K<sub>2</sub>O, MgO, NaO, and SO<sub>3</sub> in Minjingu and Panda Hill PRs.  
251

252 High concentration of Ca in Minjingu PR is also in agreement with the liming effects  
253 reported following application of Minjingu PR on acid soils [20; 33]. Apart from creating a  
254 more favorable environment for plant root growth, the liming effect of Minjingu PR on acid  
255 soils can also correct imbalance of exchangeable cations in the soil system. A combination of  
256 these effects explains the reason for higher crop response reported following application of  
257 Minjingu PR than Panda Hill PR. Figure 4 indicates that Panda Hill PR has higher  
258 concentrations of FeO<sub>3</sub>, SiO<sub>2</sub>, and AlO<sub>3</sub> than Minjingu PR. Higher concentrations of these  
259 oxides are undesirable as far as reactivity of the PR is concerned because Fe, Si, and Al have  
260 high affinity for P and therefore tend to form complex compounds with P, making it difficult  
261 to be released from the PR for plant uptake. High concentrations of these metal oxides  
262 explains the reason for low reactivity of Panda Hill PR as compared with Minjingu PR and  
263 associated differences in crop response following applications of the two PRs on soils with  
264 similar characteristics.




265

266 Figure 4. Concentrations of FeO<sub>3</sub>, SiO<sub>2</sub>, and AlO<sub>3</sub> in Minjingu and Panda Hill PRs.

267

268 With the exception of MnO<sub>3</sub> content of Panda Hill PR, all oxides determined in the  
 269 two PRs indicate low concentrations of micronutrients Zn and Cu for the two PRs to be  
 270 considered as promising source of micronutrients (Figure 5). This implies that direct  
 271 application of Minjingu or Panda Hill PR as source of P for crops will require an alternative  
 272 source of micronutrient for a balanced fertilization. Co-application of the PRs with manure or  
 273 composts may benefit plants more than just PR application alone or with industrial N  
 274 fertilizers because animal manures and composts contain most nutrients though in small  
 275 amounts.




276

277

278

Figure 5. Concentrations of ZnO<sub>3</sub>, CuO<sub>3</sub>, and MnO<sub>3</sub> in the PRs

### 279 3.6 Chemical composition of the PR-biomass Composts

280 Organic C , and P content of the composts produced are presented in Table 3.

281 Chemical analysis results indicate that OC content of the composts produced from Mucuna

282 biomass mixed with either Minjingu or Panda Hill PR was different (P=.05) from OC

283 determined in the composts of Crotalaria and Lablab biomass mixed with the same PRs.

284 Panda Hill PR composted with Mucuna biomass was found to have the highest and

285 significant (P=.05) total N concentration, followed by Minjingu PR composted with

286 Crotalaria biomass. Lablab composted with Panda Hill PR had the lowest N content of all

287 composted materials (P=.05).

288

289

290

291

292 Table 3. Selected chemical properties of composts used  
 293

Compost composition	OC	N	P	C:N	C:P	N:P
	----- % -----					
Minjingu PR + <i>Crotalaria juncea</i>	22.4 ab	1.98 b	0.51 a	11.3	42.7	4.02
Minjingu PR + <i>Lalab purpureus</i>	24.0 a	1.64 c	0.52 a	14.7	46.4	3.18
Minjingu PR + <i>Mucuna pruriens</i>	22.1 b	1.69 c	0.55 a	13.1	40.9	3.12
Panda Hill PR + <i>Crotalaria juncea</i>	23.3 a	1.70 c	0.49 a	14.1	45.4	3.80
Panda Hill PR + <i>Lablab purpureus</i>	23.2 a	1.36 d	0.48 a	16.6	49.0	2.96
PandaHill PR + <i>Mucuna pruriens</i>	21.8 b	2.16 a	0.38 b	10.8	58.0	5.37
LSD (P=0.05)	0.87	0.15	0.07	-	-	-

294 †, Values in the same column followed by different letter(s) are statistically different (p=0.05)  
 295

296  
 297 In general, the composted materials showed  $\frac{3}{2}$ ,  $\frac{3}{2}$ , and  $\frac{1}{2}$  lower contents of OC, total  
 298 N and C:N ratio as compared with the initial biomass. A decrease in OC, N and C:N ratio as  
 299 shown in Table 3 for the composts as compared with the initial biomass (Table 2) was caused  
 300 by oxidation of OC to produce carbon dioxide that was lost as CO<sub>2</sub> gas while portion of the  
 301 OC is incorporated into microbial cells. However lower total N content in the compost than  
 302 previously determined in the biomass probably caused by a dilution effect due to addition of  
 303 PR to the compost material.


304 Similar trends in total N content was reported when coffee pulp was composted with  
 305 Minjingu PR using surface soil for inoculation of the compost mix [34]. Other research  
 306 findings [35] reported a slight increase in total N of the compost relative to N content of the  
 307 raw material when coffee pulp and coffee husks were mixed with cow dung and composted  
 308 with phosphate rock following inoculation with P-solubilizing bacteria (*Bacillus*  
 309 *megatherium*). However, the increase in N content reported [35] could be due to relatively  
 310 high amount of cow dung (12 kg) equivalent to 20% of total weight of the compost mix used  
 311 to enrich the compost.



### 312 3.8 Effect of Leguminous Crop Strips on Maize Grain Yield

313 Leguminous crop strips had a significant effect on maize grain yield only when Panda  
 314 Hill PR was used as P source and the yields under Crotalaria strip was significantly greater  
 315 than those under Mucuna strip. (Figure 6). Maize grain yield obtained from the three  
 316 leguminous crop strips were 5.3, 4.5, and 4.0 t ha<sup>-1</sup> from Crotalaria, lablab and Mucuna  
 317 strips, respectively.

318




319  
 320

321 Figure 6. Effect of cover crop strips (species) on maize grain yield

322 † Values for the same PR type followed by the same letter(s) are statistically similar (p=0.05)

323

324 In Bukoba District of Tanzania, maize grain yield of 0.7 t ha<sup>-1</sup> was reported following  
 325 incorporation of crotalaria residues while lablab residues increased maize grain yield by 57-  
 326 103% above the control crop yield although the effect of lablab was below yield increase  
 327 obtained from crotalaria strips [36]. Other studies conducted in Tanzania reported maize  
 328 grain yield ranging from 1.2 to 4.0 t ha<sup>-1</sup> following incorporation of crotalaria as green  
 329 manure [37]. In South Africa, maize grain yields ranging from 2.6 to 10.6 t ha<sup>-1</sup> were reported  
 330 following incorporation of crotalaria, lablab and Mucuna [38]. Among all the leguminous

331 crops tested, maize biomass and grain yields were highest on Crotalaria plots [38]. Superior  
332 influence of Crotalaria on maize grain yields over Lablab and Mucuna was also reported in  
333 Malawi [38]. Maize grain yield obtained in this work is therefore within the range reported by  
334 other researchers in Sab Saharan Africa (SS  Superior performance of crotalaria over  
335 lablab and Mucuna also agrees with majority of research works conducted in Tanzania and  
336 neighbor countries using these leguminous crops as source of N for maize.


337

338 Other studies [39; 40] obtained results showing that incorporation of Lablab produced  
339 more maize grain yield than Crotalaria and Mucuna. Variations reported in different studies  
340 could be attributed to differences in soil property, local climatic conditions, yield potentials  
341 of maize varieties used and management practices such as timing of biomass incorporation as  
342 green manure vs. composting.

### 343 **3.9 Effect of treatments on maize grain yield**

344 Figure 7 presents maize grain yield obtained following application of different  
345 treatments. Application of Minjingu or Panda Hill PR alone failed to increase maize grain  
346 yield above the control. This observation is in agreement with findings reported by other  
347 researchers [29 and 41] following direct application of Minjingu and Panda Hill PRs on soils  
348 with varying properties. Such observations were attributed to application of PRs on soils  
349 where P is not the primary limiting factor for crop performance, as well as masking effect of  
350 moisture stress, soil acidity and deficiencies of other nutrients which affect maize yield [29,  
351 41].

352



353

354 Figure 7. Maize grain yield obtained with different treatment combinations following  
 355 leguminous crop biomass removal. † Values for the same PR type followed by the same  
 356 letter(s) are statistically similar (P=.05)

357


358 Addition of urea with Minjingu PR and Minjingu PR composted with leguminous  
 359 crop biomass increased maize grain yield by 2.40 and 1.58 t ha<sup>-1</sup>, respectively above the  
 360 control while addition of urea to Panda Hill PR and Panda Hill PR composted with  
 361 leguminous crop biomass increased grain yield only by 1.20 and 1.06 t ha<sup>-1</sup>, respectively  
 362 above the control. Difference observed in maize grain yields following the application of  
 363 Minjingu PR or Panda Hill PR alone were not significant (P =.05) even though the two PRs  
 364 have different reactivity and chemical composition.

365 Average maize grain yield produced when legume biomass was removed but  
 366 Minjingu or Panda Hill PR + urea was applied reached 5.62 t ha<sup>-1</sup> compared with 5.15t ha<sup>-1</sup>  
 367 when biomass-PR compost was applied. However, the observed difference (0.47 t ha<sup>-1</sup>) was  
 368 also not statistically significant (P= .05) indicating that biomass composted with PR was as  
 369 effective as the PR applied with urea. This suggests that legume biomass composted with PRs  
 370 could effectively substitute for the application of PRs with urea at Wang'waray FTC and  
 371 other areas with similar soil type and climatic conditions in the long run.

372

373 **3.10 Effect of Treatments on Maize Stover Yield**

374 Figure 8 indicates that stover yield was significantly different ( $P=0.05$ ) between PR  
 375 alone and PR + urea treatments.



376

377

378 Figure 8. Effects of treatments on maize stover yield following leguminous crop biomass  
 379 removal. Values for the same PR type followed by the same letter(s) are statistically similar  
 380 ( $P=0.05$ )


381

382 Application of Minjingu or Panda Hill PR with Urea produced the highest ( $2.76 \text{ t ha}^{-1}$   
 383 and  $2.54 \text{ t ha}^{-1}$ ) yield of maize stover, respectively as compared with Minjingu or Panda Hill  
 384 PR alone ( $1.92$  and  $1.66 \text{ t ha}^{-1}$ ). However, maize stover yield obtained following application  
 385 of PRs with urea and PRs composted with cover crop biomass were not statistically different  
 386 ( $P=0.05$ ) from stover yields obtained in the control plots. The lowest stover yield obtained  
 387 following application of PR alone could be due to limited supply of N and further distortion  
 388 of the balance between nutrient supply levels in the soil. This observation is in agreement  
 389 with the lowest maize grain yield obtained when PRs were applied alone and highest grain

390 yield following application of PR with urea. Application of PR alone is therefore not  
 391 economical for maize production.

### 392 3.11 Interaction of legume crop strips x fertilizer treatments effect on maize grain yield


393 With the exception of *Crotalaria* strips, when above ground crop biomass was  
 394 removed and no external input was applied, maize grain yield was below 4 t/ha. (Figures 9  
 395 and 10).



396  
 397  
 398  
 399  
 400

Figure 9. Interactional effect of leguminous crop strips and treatments with Minjingu PR on maize grain. MPR = Minjingu PR; Values followed by the same letter(s) are similar ( $P=0.05$ )

401 Following removal of *Mucuna pruriens* and *Lablab purpureus* above ground biomass, the  
 402 application of PRs without urea or compost did not increase maize grain yield compared with  
 403 the control plot. Although not significant ( $P=0.05$ ), higher maize yield was obtained on  
 404 *crotalaria* strips following removal of cover crop biomass and application of Minjingu PR  
 405 with urea, while Minjingu PR composted with biomass performed better on *lablab* strips.  
 406 Similarly *crotalaria* strips produced higher yield but not significantly different from all  
 407 treatments evaluated.



408

409 Figure 10. Interactional effect of cover crop strips and treatments with Panda Hill PR on  
 410 maize grain yield. PPR = Panda Hill PR; Values followed by the same letter(s) are  
 411 statistically similar ( $P=0.05$ )

412

413

#### 414 4.0 Conclusion

415 This study investigated the effect of three leguminous crops (*Crotalaria juncea*,  
 416 *Lablab purpureus* and *Mucuna pruriens*) biomass composted with Minjingu (medium  
 417 reactivity) or Panda Hill (low reactivity) PR on maize yield. The effect of each PR composted  
 418 with leguminous crop biomass on maize grain and stover yield was found to be similar to that  
 419 of the PRs applied with urea, while PRs applied alone failed to increase maize yield above  
 420 the controls. Similar maize yields obtained with PR-urea and PR-biomass compost treatments  
 421 imply that leguminous crop biomass composted with PRs was as effective as PRs applied  
 422 with urea in terms of P and N supply for maize. Based on these results, it was concluded that  
 423 leguminous crop biomass composted with PRs have a potential for improving maize yield  
 424 and could replace the use of highly expensive urea for maize production in the long run.  
 425 Cost-benefit analysis is however required to justify substituting urea for PR – biomass  
 426 composts in maize production.

#### 427 Reference

- 428 1. Saidia PS, Chilagane DA, Alexander W, Janet FM. Evaluation of EM technology on  
 429 maize (*Zea mays* L.) growth, development and yield in morogoro Tanzania. 2010;  
 430 Accessed 10 Aug. 2015. Available:  
 431 [kilimo.org/.../wp.../Saidia-Paul-Sabas-Chilagane-Daudi-Amos-et-al.pdf](http://kilimo.org/.../wp.../Saidia-Paul-Sabas-Chilagane-Daudi-Amos-et-al.pdf)  
 432

- 433 2. Mboya R, Tongoona, P, Derera J, Mudhara M, Langyintuo A. The dietary importance of  
434 maize in Katumba ward, Rungwe district, Tanzania, and its contribution to household  
435 food security. *African journal of Agricultural Research*. 2011; Vol. 6 (11) 2617 –  
436 2626
- 437  
438 3. USAID. The legal, regulatory, and institutional constraints to the growth of maize and  
439 rice in Tanzania. Tanzania MicroCLIR Report. 2010; Accessed 25 September 2015.  
440 Available: [egateg.usaid.gov/sites/default/files/Tanzania\\_MicroCLIR.pdf](http://egateg.usaid.gov/sites/default/files/Tanzania_MicroCLIR.pdf)
- 441 4. Rowhani P, Lobel DB, Linderman M, Ramankutty N. Climate variability and crop  
442 production in Tanzania. *Agric.Forest Meteor*. 2011; 151:449-460
- 443 5. Bekunda MA, Bationo A, Sali H. Soil fertility management in Africa. A review of selected  
444 research trials. In: Buresh RJ, Sanchez PA, Calhoun F. Eds. Replenishing soil fertility  
445 in Africa. Soil Science Society of America Special Publication No. 51. Madison,  
446 Wisconsin; 1997.
- 447  
448 6. Sizlas C, Semoka JMR, Borggard, OK. Establishment of an agronomic database for  
449 Minjingu phosphate rock and examples of its potential use. *Nutr Cycl Agroecosyst*.  
450 2007; 78:225-237.  
451
- 452 7. Kamhabwa F, Consumption of Fertilizers and Fertilizer Use by Crop in Tanzania.  
453 International Fertilizer Development Center (IFDC); 2014.  
454
- 455 8. Kpombrekou-A K, Tabatabai MA. Effect of low-molecular weight organic acids on  
456 phosphorus release and phytoavailability of phosphorus in phosphate rocks when  
457 added to soils. *Agric. Ecosyst. Environ*; 2003; (100):275-284.
- 458 9. Mowo JG, Janssen BH, Oenema O, German LA, Mrema, JP, Shemdoe RS. Soil fertility  
459 evaluation and management by smallholder farmer communities in the Northern  
460 Tanzania. *Agriculture, Ecosystems & Environment*. 2006; 116 (1-2): 47-59  
461 <http://www.wsi.nrcs.usda.gov/products/w2q/awm/docs/neh637c2.pdf>.  
462
- 463 10. National Bureau of Statistics (NBS); Office of Chief Government Statistician (OCGS)  
464 Zanzibar. 2012 Population and Housing Census: Population Distribution by  
465 Administrative Units; Key Findings. Dar es Salaam, Tanzania; 2013.  
466
- 467 11. Kpombrekou K, Tabatabai MA. Effect of organic acids on release of phosphorus from  
468 phosphate rocks. *Soil Sci*. 1994; (158) 442–453.  
469

- 470 12. FAO. Guide line for soil description. FAO, Rome, Italy; 2006  
471
- 472 13. Soil survey staff. Keys to Soil Taxonomy 12<sup>th</sup> Edition. USDA-NRSA; 2014
- 473 14. Nelson DW and Sommers L E. Total carbon, organic carbon and organic matter. In:  
474 Sparks DL, editor. Methods of Soil Analysis Part 3. Chemical Methods. SSSA Book  
475 series No.5. SSSA and ASA, Madson, Wisconsin; 1996.
- 476 15. Bremner JM. Nitrogen Total. In: DL Sparks, editor. Methods of Soil Analysis Part 3.  
477 Chemical Methods. SSSA Book series No.5. SSSA and ASA, Madson, Wisconsin;  
478 1996.
- 479 16. Kuo S. Phosphorus. In: Sparks DL, editor. Methods of Soil Analysis Part 3. Chemical  
480 Methods. SSSA Book series No.5. SSSA, ASA. Madson, Wisconsin; 1996.
- 481 17. Okalebo JR, Gathua KW, Woomer PL. Laboratory methods of soil and plant analysis.  
482 Working manual. UNESCO Press, Nairobi, Kenya; 1993.
- 483 18. FAO. On farm composting methods. Land and water discussion paper 2. Rome; 2003.
- 484 19. Nkonya E, Xavery P, Ankomay H, Mwangi F, Ponia, A, Moshi A. Maize production  
485 technologies in the Northern Zone Tanzania. Mimeo. Kansas University; 1998.
- 486 20. Landon J R, editor. Booker Tropical Soil Manual. A handbook for soil survey and  
487 agricultural land evaluation in the tropics and subtropics. Addison Wesley Longman  
488 Limited, England; 1991.
- 489 21. Dierolf T S, Fairhurst TH, Mutert EW. Soil Fertility Kit: A Toolkit for Acid Upland Soil  
490 Fertility Evaluation and Management in South East Asia. Potash and Phosphate  
491 Institute, Canada; 2001.
- 492 22. Reisenauer HM, Walsh LM, Hoefl RG. Testing soil for S, B, Mo, and Cl. In: Walsh, M.L.  
493 and Beaton JD, editors. Soil testing and plant analysis, SSSA Inc. Madson,  
494 Wisconsin; 1973.
- 495 23. Gee GW, Bauder J.W. Particle-size analysis. In: Klute A, editor. Methods of soil analysis:  
496 Part 1—physical and mineralogical methods. SSSA Book Series 5.1. SSSA, ASA.  
497 Madson, Wisconsin; 1986.  
498
- 499 24. Maranguit D, Nguillaume T, Kuzyakov Y. Land use change affects phosphorus fractions  
500 in highly weathered tropical soils. *Catena*. 2017; 149: 385-393.  
501
- 502 25. Sakala WD, Kumwenda JDT, Saka AR. The potential of green manure to increase soil  
503 fertility and maize yield in Malawi. *Biol Agr & Hort*. 2002; 21(2):121-130.  
504
- 505 26. Graves ER, Hattemer G M. Composting: National Engineering Handbook. 2000.  
506 Accessed 30 June 2009. Available:  
507 [www.springerlink.com/index/11P842T5VW685N40.pdf](http://www.springerlink.com/index/11P842T5VW685N40.pdf)



- 508 27. Van Straaten P. Rocks for crops: Agrominerals of sub-Saharan Africa. ICRAF, Nairobi,  
509 Kenya; 2002.  
510
- 511 28. FAO *Use of phosphate rock for sustainable agriculture*. FAO Fertilizer and Plant  
512 Nutrition Bulletin No. 13. Rome; 2004.  
513
- 514 29. Semoka JMR, Mnkeni PNS, Ringo HD. Effectiveness of Tanzania phosphate rocks of  
515 igneous and sedimentary origin as source of phosphorus for maize. *Zimb Jour Agr Res*.  
516 1992; 30 (2):127 - 136.
- 517 30. Van Straaten P, Editor. Rock for Crops. Agro minerals of sub-saharan Africa. Fidelity  
518 National Information Solutions Canada, Scarborough, ON, M1B 3C3, Canada; 2002.
- 519 31. Mnkeni PNS, Semoka JMR, Kaitaba EG. Effects of Mapogoro phillipsite on availability  
520 of phosphorus in phosphate rocks. *Trop. Agric*. 1994;71:249-253.  
521
- 522 32. Szilas C, Semoka JMR, Borggaard OK. Can local Minjingu phosphate rock replace super  
523 phosphate on acid soils in Tanzania? 2006. Accessed 8 June 201. Available:  
524
- 525 33. Szilas C. The Tanzanian Minjingu phosphate rock - possibilities and limitations for direct  
526 application. PhD Thesis. Royal Veterinary and Agricultural University, Copenhagen,  
527 Denmark; 2002.  
528
- 529 34. Shitindi MJ. Response of tomato (*Lycopersicon esculentum M.*) to coffee pulp compost,  
530 Minjingu phosphate rock and coffee pulp - minjingu phosphate rock compost applied  
531 to a Chromic Acrisol. MSc. Thesis . Sokoine University of Agriculture. Morogoro,  
532 Tanzania; 2011.
- 533 35. Preethu DC, Bhanu Prakash BNUH, Srinivasamurthy CA, Vasanthi BG. Maturity indices  
534 as an index to evaluate the quality of compost of coffee waste blended with other  
535 organic wastes. Proceedings of the International Conference on Sustainable Solid  
536 Waste Management.5-7 September 2007, Chennai India. 2007.  
537
- 538 36. Bajjukya FP, de Ridder N. Giller KE. Nitrogen release from decomposing residues of  
539 leguminous cover crops and their effect on maize yield on depleted soils of Bukoba  
540 District, Tanzania. *Pant and Soil*. 2006; 279: 77-93.
- 541 37. Lupatu M, Kilimwiko L. Natural fertilizers: New life for tired soils. Africa Farmer No.6;  
542 December 1991. Hunger Project Savaccon gallery, 240 E. 13<sup>th</sup> street New York; 1991.
- 543 38 Odhiambo JJO, Ogola JBO, Madzivhandila T. Effect of green manure legume – maize  
544 rotation on maize yield and weed infestation levels. *Afr. J. Agric. Res*. 2010; 5(8):  
545 618-625.  
546

- 547 39. Mureiythi JG Gachene CKK, Ojien J. The role of green manure legumes in smallholder  
548 farming systems in Kenya. The legume research network project. *Tropical and*  
549 *Subtropical Agro ecosystems*. 2003; 1:57-70.
- 550 40. Odhiambo JJO. Potential use of green manure legume cover crops in smallholder maize  
551 production systems in Limpopo province, South Africa. *Afr Jour Agr Res*. 2011; Vol.  
552 6(1): 107-112.