

Original Research Article
Biometrical assessment of urucum quantitative features (*Bixa orellana* L.) in the northeastern Brazil

ABSTRACT

Urucum (*Bixa orellana* L.) is a shrub plant occurring in the caatinga biome. This study evaluated the biometrical assessment of urucum quantitative features of Urucum (*Bixa orellana* L.) in the northeastern Brazil. We Fruit samples d the fruits were taken from within an area of native vegetation of the Federal Rural Semi-Arid University (UFERSA), Mossoró, RN (full form?). The measurements on the seed length and width of fruit samples s were done at taken to the plant science laboratory to measure the seeds length and width. Descriptive and graphical analysises were carried out using the R statistical software R. The length and width showed a small range of variation with, resulting in an excellent and reasonable values of coefficients of variation, respectively. We found a regular degree of symmetry and a mesokurtic distribution was observed for length and width of seeds. There was non-significant linear correlation between length and width. The features of Urucum seeds did not fit to the normal probability distribution of probability.

Comment [di1]: Don't use we word

Key words: Medicinal plant; descriptive statistical; inference.

1. INTRODUCTION

The Urucum or urucu (*Bixa orellana* L.), a shrub species native to from tropical region of America, has gained importance in the world agricultural market every year and is cultivated in the tropical areas all over the world [1]. The fruits are ovoid capsules containing 30 to 40 seeds [2]. The Urucum produces particular atoxic coloring substances with the indication as a hypolipidemic agent, which makes it one of the few pigments that is allowed by the World Health Organization [3].

The species has multiple uses for, such as ornamental and medicinal values; and plant and soil restorer properties. It Urucum is a pioneer species, native to the Amazon Rainforest. Its seeds are used to produce condiments and tinctures that are widely used in the industry. The fast growth nature of the uUrucum allows its the cultivation together with other species in degraded areas destined to reforestation [2].

Morphological analysises of fruits and seeds help to understand the germination process and the characterization of vigor and viability characteristics [4]. Biometric analysises comprises an essential tool to detect genetic variability within and between populations and to define the relationships between the variability and environmental factors, thus contributing to genetic improvement programs [5].

Biometrics of fruits and seed are central in studies of species, providing data for differentiation of species within the same genus. For example, [6] showed that the seeds of *Hymenaea courbaril* L weight almost twice the seeds of *Heritiera parvifolia*. Also, the seeds characteristics influence its patterns of dispersion and establishment of seedlings [7], which are used to differentiate pioneer and non-pioneer species in tropical forests [8]. In most shrubs and trees, there is an inverse relation between seed size and the number of seeds per fruit [9].

Urucum stands out for its therapeutic **and food properties** **potential and as food**. However, there is no record of its exploration in the region of Mossoró, RN, **as well as, studies of urucum biometry, showing the need for studies concerning this species.**

This work evaluated the biometry of **U**urucum seeds as a subsidy **for** studies **for** comparing their characteristics when submitted to different environments.

2. MATERIAL AND METHODS

The study was carried out in Mossoró, RN, geographic coordinates: 5°11'S and 37°20'W at 18 m of altitude, with an annual mean temperature of 27.5°C and relative humidity of 68.9% [10]. **According to Köppen's classification, T**the climate in Mossoró is BSw', hot and dry. **Of total, During May 2018, we collected a sample of 300 urucum seed** **samples were collected from s in an area of the n**native vegetation **located inside the campus** of the Federal Rural Semi-Arid University (UFERSA). **Brazil during May 2018., in the municipality of Mossoró, State of Rio Grande do Norte, Brazil, and were taken to the plant breeding laboratory, where t**The morphological characterization of the seed was evaluated **in the plant breeding laboratory., and t**The width and length of well-developed seeds were determined **in millimeters** with the precision of a 0.1 mm pachymeter. The descriptive analysis and graphs were done with the software R version 3.1.1. [11].

3. RESULTS

To determine and compare quantitative aspects of the distributions of values of the variables seeds length and width **we** based the analysis on specialized biometric literature [12,13,14,15,16,17,18]. Thus, **we** adopted the exploratory data analysis using frequency distributions, box plots, as well as the statistical estimators of the variables under study, which are the main descriptive and inferential statistical measures, such as arithmetic mean, median, total range, variance, standard deviation, standard error of mean, coefficient of variation, asymmetry coefficient, kurtosis coefficient, quartiles and interquartile deviation. We used the Pearson

Comment [di2]: Don't use we, so paraphrase this sentence

Comment [di3]: Don't use we

correlation coefficient and statistical inference such as the hypothesis tests, T-test or Z-test, at a significance level of 5% probability, based on Student's t-distribution and Normal distribution, respectively, for the construction of confidence intervals with 95% probability (Tables 1 to 3 and Figures 1 to 7).

Table 1. Frequency distribution of seed length (mm) of Urucum (*Bixa orellana* L.)

Class (Length in mm)	f_i	X_i	$f\%$
[2.9-----3.2)	2	3.1	0.67
[3.2-----3.4)	5	3.3	1.67
[3.4-----3.7)	9	3.5	3.00
[3.7-----3.9)	21	3.8	7.00
[3.9-----4.1)	45	4.0	15.00
[4.1-----4.4)	72	4.3	24.00
[4.4-----4.6)	84	4.5	28.00
[4.6-----4.9)	51	4.7	17.00
[4.9-----5.1)	11	5.0	3.67
Total	300	-----	100

Table 2. Frequency distribution of seed width (mm) of Urucum (*Bixa orellana* L.)

Classes (Width in mm)	f_i	X_i	$f\%$
[1.7-----1.9)	4	1.8	1.33 %
[1.9-----2.2)	28	2.1	9.33
[2.2-----2.5)	41	2.4	13.67
[2.5-----2.8)	47	2.6	15.67
[2.8-----3.1)	58	2.9	19.33
[3.1-----3.3)	44	3.2	14.67
[3.3-----3.6)	32	3.5	10.67
[3.6-----3.9)	33	3.8	11.00
[3.9-----4.2)	13	4.0	4.33
Total	300	-----	100

Table 3. Results of the descriptive and inferential statistics of length (mm) and width (mm) of the 300 seeds of Urucum (*Bixa orellana* L.)

Statistics or Estimator	Length	Width
Sample size (number of seeds)	300	300
Minimum value	2.94	1.65
Maximum value	5.05	4.17
Total range	2.11	2.52
Arithmetic mean	4.32	2.94
Median or second quartile	4.38	2.94
Variance	0.13	0.33
First quartile	4.13	2.51
Third quartile	4.57	3.36
Standard deviation	0.37	0.57
Standard Error	0.02	0.03
Coefficient of Variation (%)	8.44	19.40
Asymmetry Coefficient	-0.85	0.08
Kurtosis Coefficient	1.12	-0.86
Interquartile Range (IR)	0.44	0.85
Z Test for mean, at 0.1% of probability	216.00***	98.00***
Confidence interval for mean at 95% of probability	4.29 to 4.36	2.88 to 3.00
Theoretical Normal Distribution or Gaussian	p -value = 0.01	p -value = 0.06
Probabilities, D'Agostino-Pearson's Test	Fitted to normal distribution	Fitted to normal distribution
Pearson's Simple Linear Correlation Coefficient (r)	+ 0.16 (p-value of Student's T test = 0.10)	




Figure 1. Histogram and polygon of frequencies representative of the length distribution (mm) of urucum seeds (*Bixa orellana* L.)




Figure 2. Boxplot using median and quartiles of length of seeds (mm) of Urucum (*Bixa orellana* L.)




Figure 3. Box plot using the mean and standard deviation of seeds length (mm) of urucum (*Bixa orellana* L.)




Figure 4. Histogram and polygon of frequencies of the width (mm) distribution of urucum seeds (*Bixa orellana* L.)




Figure 5. Box plot using median and quartiles of the seeds width (mm) of urucum (*Bixa orellana* L.)




Figure 6. Boxplot using the mean and standard deviation of seed width (mm) of Urucum (*Bixa orellana* L.)




Figure 7. Pearson linear correlation dispersion diagram for length and width (mm) of 300 urucum seeds (*Bixa orellana* L.).

Comment [di4]: You have not described the results/data of table and figures?

4. DISCUSSION

The length and width of seeds had a small amplitude of variation (Table 3) and good coefficients of variation, showing a high degree of homogeneity. Data dispersion was higher for width than length. There is not a great similarity in the location of the length and the width of the seeds, since the means 4.32 and 2.94 mm for the average length and the average width, respectively, are not very close. It is worth noting that the calculation of the average does not have to make real sense. However, it can be stated that the seeds have a center of gravity in the distribution of these variables, which shows that the length of the seeds is close to 4.32 millimeters and the width around 2.94 millimeters. The relative variation between the variables measured by the coefficients of variation, in this case, was the most adequate, since they are independent of the magnitude of the variable, as well as of its unit of measure, being, therefore, the most correct procedure. The coefficients of variation (8.44 e 19.40% for length and width, respectively) showed a similar or close relative dispersion for the two characteristics. This coefficient was the most adequate to measure data variability since they are independent of the magnitude of variables and their units. Even though there was less relative variability for the length with a significant difference in the magnitude compared to the width [12,14,16,19].

Other results obtained, as can be seen in Table 3, were that the mean and median were very close to the length and equal to the width, which indicated a reasonable and a high degree of symmetry for length and width, respectively, which were -0.85 and 0.08 for the length and width, respectively. However, the momentum coefficients of kurtosis, which were 1.12 and -0.86, respectively, showed that both length and width had a mesocuric distribution, and the

Comment [di5]: Discuss about the similar works results done previously, only keeping the references would not be sufficient?

characterization of the degree of asymmetry and kurtosis of a distribution according to [13,16,20,21,19], can not be done by looking only at measures of position or central tendency such as the average and the median, but also through the asymmetry and kurtosis coefficients, as well as through histograms and frequency polygons (Figures 1 and 4).

With respect to quantiles, the first quartile, 25% of the lowest values, showed a maximum value of 13.24 mm length and 9.78 mm for width. The 25% larger seed lengths and widths had at least 14.67 and 10.97 mm, respectively, corroborating data of [22]. The interquartile range verifies the dispersion of the data concerning the median and thus, identify the presence of outliers in the data. The interquartile was 0.44 mm for the length and 0.85 mm for the width of the seed (Table 3).

The degree of association or simple linear correlation, whose values range from -1 to +1, including zero, and its magnitude is dimensionless, between length and seed width of urucum was 0.16 (Table 3), showing a non-significant positive relation, as evidenced by the result of Student's t-test [12,22,14,13,15,16,17,18,23].

From a graphic, visual or geometric point of view, the dispersion diagram (Figure 7), showed high dispersion which confirms the lack of correlation between length and width found in the low value of correlation coefficient (0,16). The length and width variables of urucum seed did not fit to the theoretical or special density distribution of normal or bivariate Gaussian probability so that all the conclusions obtained through statistical inference are not guaranteed or assured by this assumption (Table 3).

Reporting the quantiles comprises an excellent way to illustrate the dispersion of a distribution. Researchers are more familiar with the type of quantile called percentile because of its use in standardized tests. When a test score is reported within the 90th percentile, 90% of the scores are smaller than it, and 10% are higher. Contrary to variance and standard deviation, quantile values do not depend on arithmetic mean or median values. When distributions are asymmetric or have outliers, which are extreme values that are not characteristic of the distribution quantile box plots can portray the distribution of data more accurately than mean and standard deviation.

We obtained high values at the Z test, both for seed length and seed width, concluding that the mean values of these characteristics were highly significant (Table 3).

Boxplots, using the median and the quartiles are a widely used chart in the biological and medical sciences, showing the median, first and third quartiles. It also shows the lowest and highest scores through the lower and upper boundaries of vertical straight lines, which originate from the first and third quartiles, respectively. According to the results obtained in Figure 2, using the median and quartiles, a strong concentration of the seed length data in millimeters around the central value in the case the median was observed, which evidenced a high homogeneity of these observations, which makes its analysis and current and subsequent interpretation, including the

making of statistical inferences such as the construction of confidence intervals and the application of hypothesis test tests, as well as the adjustment of regression models for estimation and forecasting purposes.

Using the mean and standard deviation, the box plot graph, similar to the previous graph, shows the mean and standard deviation in the Box. It also shows the lowest and highest scores through the upper and lower boundary of vertical straight lines where the presence of outliers can also be verified. (Figure 3). In the case of seed length, only an atypical value occurred, showing a high similarity among the grouped observations, except for the presence of this unusual observation. Also according to the results obtained in Figure 3, most seed length data clumped around the central value, in the case the mean, which suggests a high homogeneity of these observations.

We observed a strong concentration of seed width data around the central value (median), demonstrating a high homogeneity (Figure 5). Using the mean and standard deviation (Figure 6), there was no atypical value, showing a relevant similarity in the grouped observations and a strong concentration of the seed width data around the central value (mean), which also evidenced the high homogeneity.

Comment [di6]:

In general, the results of the descriptive measures of location, variability, asymmetry and kurtosis can serve as a basis for future studies of descriptive analysis and statistical inference, for the comparison of different environments, genetic improvement studies, grouping of experiments in joint analysis, in stability analysis of cultivars, as well as in the construction of so-called components of variance [12,22,14,13,15,16,17,18,23].

Standard deviation and variance are special cases of what statisticians and physicists call the central momentum (CM). Central moment comprises the mean deviation of all observations in a data set from the mean of the observations, raised to the power r . The first central moment ($r = 1$) measures the sum of the differences between each observation minus the sample mean (arithmetic), which is always equal to zero. The second central moment ($r = 2$) is the variance. The third central moment ($r = 3$), divided by the standard deviation to the cube (s^3), is the asymmetry. The asymmetry describes how the sample differs from the form of a symmetric distribution. A normal distribution has an asymmetry coefficient equal to zero. A distribution in which the value of the asymmetry coefficient is greater than zero has asymmetry to the right, that is, there is a long tail of larger observations at the right of the mean. However, if the asymmetry coefficient is less than zero it has asymmetry on the left, there is a long tail of smaller observations to the left of the mean. The kurtosis or flattening has its basis in the fourth central moment ($r = 4$), measuring the extent or peak at which the probability density is distributed in the tails versus the center of the distribution. The distribution is classified as heavy tail or light tail when compared to a standard normal distribution

(mesokurtic). Aggregate or platykurtic (flattened) distributions have a kurtosis coefficient less than zero, compared to the normal distribution, meaning more mass of probability in the center of the distribution and less probability at the tails. In contrast, leptokurtic (tapered) distributions have a kurtosis coefficient greater than zero. Leptokurtic distributions have less mass of probability in the center and tails of relatively heavy probabilities [24].

According to [14], the law of large numbers proves that for an infinitely large number of

observations, the formula $\frac{\sum_{i=1}^n Y_i}{n}$ is an approximation of the population mean μ , where $Y_n = [Y_i]$ is the sample of size n of a random variable Y with expected value ($E(Y)$). Similarly, the variance of $y_n = \frac{\sigma^2}{n}$. Since the standard deviation is just the square root of the variance, y_n is given by $\sqrt{\frac{\sigma^2}{n}} = \frac{\sigma}{\sqrt{n}}$, what is the same as the standard error of the mean. Therefore, we have an estimate of the standard deviation of the population mean, which is the standard error.

If the conclusions based on a single sample are representative of the entire population, it is recommended using the standard error of the mean. However, if the samples limit the conclusions, it is better to use the sample standard deviation.

Large observational surveys covering large spatial scales with a substantial number of samples are likely representative of the population of interest as a whole, and the standard error of the mean should be used. Small, controlled experiments with few replicates are likely based on a single cluster, and possibly unrepresentative of individuals, as a consequence, the standard deviation should be used to characterize or measure the degree of absolute dispersion of the samples.

We conclude that the length and width of the seeds showed a low amplitude of variation, as well as an excellent value for the coefficients of variation, showing a reasonable degree of homogeneity, being the length less dispersed than the width.

There was a regular to high degree of symmetry and a mesokurtic distribution for the length and width of the seeds.

Seed length and width characteristics had no significant correlation and showed mean values with highly significant differences.

The data of urucum seed biometry did not fit to the normal distribution probability.

4. Conclusion

The results of the descriptive measures of location, variability, asymmetry, and kurtosis can aid in future studies of descriptive analysis and statistical inference, for the comparison of different environments, studies of plant genetic improvement, and support criteria used for grouping of experiments in analysis, study stability of cultivars, in multivariate analysis, as well as support the construction of so-called components of variance.

Comment [di7]: Better to use short and clear simple sentences instead long one.

References

1. Mercadante AZ, Pfander H. Carotenoids from annatto: a review. Recent Research Developments in Agriculture and Food Chemistry. 1998; 2 (1): 79-91.
Available:http://www.scielo.br/scielo.php?script=sci_nlinks&ref=000112&pid=S1983...lng=pt
2. Lorenzi H. Brazilian trees: manual of identification and cultivation of native tree plants of Brazil. New Odessa: Plantarum. 1998; 2 (1): 352 p. Portuguese.
Available:<https://www.bdpa.cnptia.embrapa.br/consulta/busca?b=ad...%22LORENZI...>
3. Silva JHV, Silva EL, Filho JJ, Ribeiro MLG, Costa FGP. Residue of annatto seed (*Bixa orellana* L.) as dye of yolk, skin, beak and ovary of laying hens evaluated by two analytical methods. Ciência e Agrotecnologia. 2006; 30 (5): 988-994. Portuguese.
Available:<https://www.scielo.br/pdf/cagro/v30n5/v30n5a24.pdf>
4. Mathus MT, Lopes JC. Morphology of fruits and seedlings and seed germination of *Erythrina variegata* L. Revista Brasileira de Sementes. 2007; 29 (3): 8-15.
Available:<https://www.redalyc.org/pdf/744/74413024004.pdf>
5. Gusmão E, Vieira FA, Júnior EMF. Fruit biometry and murici endocarps (*Byrsonima verbascifolia* Rich, Ex. A. Juss). Revista Cerne. 2006; 12 (1): 84-91. Portuguese.
Available:https://www.researchgate.net/profile/Fabio_Vieira2/publication...
6. Carpanezzi AA, Marques LCT. Germination of jutaí-açu (*Hymenaea courbaril* L.) and jutaí-mirim (*H. parvifolia* Huber) seeds scarified with commercial sulfuric acid. Circular Técnica 19. EMBRAPA-CPATU, Belém. 1981.15p. Portuguese.
Available:<https://www.embrapa.br/busca-de-publicacoes/-/publicacao/376633/...>
7. Fenner M. Seed ecology. London: Chapman & Hall, 1993. 151 p.
Available:https://link.springer.com/chapter/10.1007/978-1-4615-1619-4_4
8. Baskin CC, Baskin JM. Seeds: ecology, biogeography and evolution of dormancy and Germination. London: Academic Press, 1998. 666p.
Available:<https://link.springer.com/content/pdf/bbm:978-3-642-55974-7/1.pdf>

9. Carvalho JEU, Nascimento WMO, Muller CH. Physical characteristics and seed germination of fruit species native to the Amazon. Research Bulletin 203. EMBRAPA-CPATU, Belém. 1998.18p. Portuguese.
Available:https://www.scielo.br/scielo.php?script=sci_arttext&pid=S0100...
10. Carmo Filho F, Oliveira OF. Mossoró: a municipality of the northeastern semi-arid region, climatic characterization and floristic aspect . Mossoró: ESAM, 1995. 62 p. (Coleção Mossoroense, Série B). Portuguese.
Available:http://www.scielo.br/scielo.php?script=sci_nlinks&pid=S0102...
11. R Version 3.1.1. Viena, Áustria: Foundation for Statistical Computing, 2014. (Software). RAO KVK, Schwartz SA, Nair HK, Aalinkel R, Mahajan S, Chawda R, Nair MPN. Plant derived products as a source of cellular growth inhibitory phytochemicals on PC-3M, DU-145 and LNCaP prostate cancer cell lines. Current Science. 2004; 87 (1): 1585-1588.
12. Ferreira DF. Basic statistics. Lavras: UFLA, 2005. 625 p. Portuguese.
Available:<http://www.scholar.google.com.br/citations?user=M80iQtgAAAAJ&hl=pt-BR>
13. Spiegel MR, Stephens LJ. Statistics. 4 ed. Porto Alegre: Bookman, 2009. 597 p. Portuguese.
Available:<https://www.estantevirtual.com.br/livros/murray-r-spiegel/estatistica/3772227637>
14. Oliveira MS. Introduction to statistics. Lavras, MG: Publisher of the Federal University of Lavras (UFLA), 2009. 329 p. Portuguese.
Available:<https://www.editora.ufla.br/>
15. Bussab WO, Morettin PA. Basic statistics. 6. ed. São Paulo: Saraiva, 2010. 540 p. Portuguese.
Available:<https://www.saraiva.com.br/livros/editoras/saraiva>
16. ZAR, J. H. Biostatistical analysis. 5. ed. New York: Prentice Hall, 2010. 944 p
Available:<https://books.google.com.br/books?isbn=1119168996>
17. Claudio LC, Stein CE. Descriptive statistics and probability theory. 2. ed. Blumenau : Edifurb, 2011. 213 p. Portuguese.
Available:<https://books.google.com.br/books?isbn=853526356X>
18. Cecon PR, Silva ARS, Nascimento M, Ferreira A. Statistical methods. Viçosa: Editora UFV, 2012. 229p. Portuguese.
Available:<https://www.locus.ufv.br/bitstream/handle/123456789/9211/texto%20completo.pdf?...1...>
19. Fonseca JS, Martins GA. Statistical course. 6. ed. 15. reimp. São Paulo: Atlas, 2012. 320 p.
Available:<https://www.passeidireto.com/arquivo/46902730/livro-texto...iipdf-estatistica/13>

20. Andrade DF, Ogliari PJ. Statistics for the agricultural and biological sciences with notions of experimentation. 3 ed. Florianópolis: Editora da UFSC, 2010. 470 p. Portuguese.
Available:<https://www.passeidireto.com/.../livro-estatistica-basica-para-ciencias-agrarias-e-biolog...>
21. Casella G, Berger RL. Statistical inference. São Paulo: Editora Cengage Learning. 2010. 612 p. Portuguese.
Available:<https://www.cengage.com.br/livro/inferencia-estatistica-traducao-da-2a-edicao-norte-americana>
22. Figueiredo F, Figueiredo A, Ramos A, Teles P. Descriptive statistics and probabilities - solved and proposed problems with R. applications Escolar Editora. Lisboa: Portugal, 2007. 420 p.
Available:[https://www.wook.pt > Livros em Português > Ciências Exatas e Naturais > Matemática](https://www.wook.pt/Livros-em-Portugu%C3%AAs/Ci%C3%BAncias-Exatas-e-Naturais/Matem%C3%A1tica)
23. Costa GGO. Course of inferential statistics and probabilities: theory and practice. São Paulo: Editora Atlas, 2012. 370 p. Portuguese.
Available:<https://www.grupogen.com.br/curso-estatistica-inferencial-probabilidades>
24. Gotelli NJ, Ellison AM. Principles of statistics in ecology. São Paulo: Artmed Editora. 2010. 532 p. Portuguese.
Available:<https://www.passeidireto.com/.../principios-de-estatistica-em-ecologia---gotelli-nichola...>