

1
2
3
4
5
6
7
8
9
10
11
12

Original Research Article

Liquid *Bacillus subtilis* Formulation in Rice in for the Control of *Meloidogyne javanica* and Lettuce Development Improvement

ABSTRACT

Aim: To develop a liquid formulation based on *Bacillus subtilis*-34 using rice to evaluate the shelf life under refrigerated and room conditions and to evaluate the effect of different application addition times of the formulation on the control of *Meloidogyne javanica* and growth of lettuce plants.

Statistical Design: The design was completely randomized, with five treatments and eight replicates. The results were submitted to analysis of variance and the means were compared by the Scott Knott test with 5% error probability.

Location and Duration of the experiment: The experiment was set up during the period from 02/13/2018 to 03/20/2018 in greenhouse located at the State University of Montes Claros, municipality of Janaúba, MG, Brazil.

Methodology: Treatments consisted of irrigation drench in the substrate of tubes at 8 and 15 days; irrigation drench in the substrate of tubes at 8 and 15 days and in pot at 25 and 35 days; irrigation in pot only at 25 and 35 days after transplanting and 2 controls (Onix[®] and absolute control). All pots with plants were infested inoculated with 5000 nematode eggs. At 45 days of transplanting, the following nematological variables were evaluated: number of galls, number of egg mass, number of eggs per gram and reproduction factor, and agronomic variables fresh and dry biomass.

Results: Applications Additions in the tube and pot and in the pot only were efficient for the reduction in the reproduction of *M. javanica* and for the development improvement of lettuce plant growth.

Conclusion: *B. subtilis*-34 remains viable until 9 months in formulation stored under refrigerator and up to 7 months under room conditions.

13
14
15
16
17
18
19
20
21
22
23
24
25
26

Keywords: *Lactuca sativa* L.; rhizobacteria; root-knot nematode, technology; shelf life

1. INTRODUCTION

Lettuce (*Lactuca sativa* L.) is a vegetable economically important for Brazil, being cultivated in almost all regions of the country [1]. Lettuce is a good source of fiber, vitamins, especially B, A, C and K, low in calories and a rich source of pigments beneficial to human health [2]. In addition to the nutritional aspect, it is also a culture of great importance from the social point of view, being cultivated mainly by family farmers near large urban centers in the so-called "green belts" [3].

27 Lettuce has numerous phytosanitary problems, among which phytonematodes stand out.
28 Nematodes of the genus *Meloidogyne* are considered as limiting factor of the commercial
29 cultivation of several vegetables, since they have short cycle and are always cultivated in the
30 same area, favoring the increase in the nematode population. Losses caused by
31 phytonematodes in vegetable crops are estimated at 12.3% in developed countries and
32 14.6% in developing countries [4]. Lettuce plants, attacked by nematodes, are less
33 developed improved due to dense formation of galls in the root system and their control is a
34 difficult task.

35
36 Several strategies have been used for the control of nematodes, among them: crop rotation,
37 solarization, use of resistant cultivars and chemical control. Most lettuce cultivars present
38 high susceptibility to *Meloidogyne* species [5]. The species of this genus most important to
39 lettuce and other leafy crops are *M. javanica* and *M. incognita* [6]. The damage caused by
40 agrochemicals, crop rotation infeasibility in small areas and the cost of plastic for solarization
41 evidences the necessity of the use of biological control agents, among them rhizobacteria
42 [7].

43
44 Among rhizobacteria, the genus *Bacillus* stands out, which has the capacity to produce
45 antibiotics, enzymes and toxins, which act directly, causing the mortality of juveniles and / or
46 indirectly affecting their behavior, feeding or reproduction. Plant-host recognition, resistance
47 induction and / or plant growth promotion processes can also be performed [8]. In addition,
48 *Bacillus* produces endospores, which are spores that survive under conditions of nutrient
49 deprivation and conditions of high temperature, which favors the maintenance of the viability
50 of formulations [9].

51
52 Promising results have been obtained in the control of *M. javanica* by *B. subtilis* in banana
53 and tomato crops plants [10]. *Bacillus* has been described for producing hydrolytic enzymes
54 such as lipases, chitinases and proteases capable of degrading structural components of
55 *Meloidogyne* [11, 12, 13].

56
57 The formulation of biocontrol agents such as rhizobacteria or another microorganism is an
58 essential step for their commercial use [14]. Formulations should improve shelf life since
59 biocontrol agents are living organisms. In addition, formulations should be economical and
60 contain sufficient number of viable colony forming units (CFUs) and be easily applied to soil
61 or plants [15]. Several farmers have multiplied bacteria from organic products on their
62 properties. Thus, the development of low-cost formulations for large-scale use is essential.

63
64 In this context, the aims of this work were to evaluate the application of a liquid formulation
65 of *B. subtilis* produced in rice broth and determine its efficiency in for the control of *M.*
66 *javanica* and in for the promotion of lettuce growth.

68 2. MATERIAL AND METHODS

69 70 2.1 Development of *Bacillus subtilis* formulation in rice broth and 71 establishment of the growth curve

72 *Bacillus subtilis*-34 isolate maintained in mineral water in glass tubes under room conditions
73 was used. From this suspension, a volume of 1.85 ml was collected. This suspension was
74 placed in 1 liter erlenmeyers and the following components were added in g.L⁻¹ of distilled
75 water: 185 g of raw rice, 185 g of sugar, 55.55 g of sodium chloride (NaCl), 46.29 g of
76 phosphate of potassium monobasic (KH₂PO₄). Chemical compounds, sugar, water and rice
77 were autoclaved at 1.0 atm at 120 ° C for 30 minutes. The formulation had final pH of 7 (±
78 0.2).

79

80 For determination of the initial number of CFUs, a one-milliliter aliquot of the liquid
81 formulation was submitted to a 10^{-5} dilution and 100 μL were collected for plating in Petri
82 dishes containing Tryptic Soy Agar (TSA). Petri dishes were incubated at 25°C for 24 hours,
83 when the initial number of $\text{CFU}\cdot\text{mL}^{-1}$ was evaluated. For the establishment of the growth
84 curve, the formulation was incubated for 44 hours on an orbital shaker at 28°C at 220 rpm
85 and at 4-hour intervals; the same procedure was performed to determine the number of
86 CFUs. The growth curve of the bacteria was also determined in Tryptic Soy Broth (TSB)
87 medium, following the same methodology as that used for the liquid rice formulation.
88

89 **2.2 Evaluation of different application addition times of the liquid *Bacillus*** 90 ***subtilis* formulation in rice broth in the control of *Meloidogyne javanica* and in** 91 **lettuce development improvement**

92
93 The experiment was carried out in greenhouse at the State University of Montes Claros,
94 Janaúba, MG, with the following geographical coordinates ($43^{\circ} 16'18.2''\text{W}$ and $15^{\circ} 49'51.5''$
95 S) and average altitude of approximately 540 m a.s.l.). For the evaluation of growth
96 promotion and reduction of nematological variables, *B. subtilis*-34 bacterial isolate was used.
97 The formulation was made as reported in item 2.1.
98

99 The isolate growth was interrupted 28 hours after the beginning of incubation. At this time,
100 dilution at 10^{-5} and TSA plating was performed to determine the initial number of CFUs per
101 ml. To evaluate the survival period of the bacterium in the liquid formulation, a volume of 50
102 ml was kept at room temperature in the laboratory of Phytopathology on the bench with
103 mean temperature of 26.05°C (maximum of 29.1°C and minimum of 23°C) and another 50
104 ml kept in refrigerator at 9°C . At one-month intervals for a period of 10 months, 10^{-5} dilution
105 and TSA plating were performed to determine the number of CFUs.
106

107 Lettuce seedlings 'Aurélia' cultivars were obtained from the sowing in styrofoam tubes
108 containing Bioplant[®] substrate. After 17 days, seedlings were transplanted to 3-liter pots
109 containing: substrate composed of soil (heavy clay, 26.6% silt, 60% clay, 13.4% sand, pH =
110 5) in 3:1 proportion, respectively, previously autoclaved at 1 atm and 120°C for 30 minutes,
111 three times at 24-hour intervals. The substrate was fertilized as recommended for the
112 culture. Prior to assay setup, the substrate was submitted to liming and incubated for 30
113 days.
114

115 The experiment was set up in a completely randomized design with five treatments and eight
116 replicates. Treatments consisted of: T1- irrigation drench of the formulation to the substrate
117 in tubes at 8 and 15 days, T2 - added via irrigation drench to the substrate of tubes at 8 and
118 15 days and in the pot at 25 and 35 days, T3- irrigation drench of the formulation to the soil
119 of pots at 25 and 35 days after transplanting and 2 controls; T4-Onix[®] (Commercial product
120 based on *Bacillus methylophilicus* - Isolated UFPEDA 20) and T5- without bacterium
121 application addition and without commercial product. In Onix[®] treatment, each plant received
122 250 ml of the commercial product, previously diluted in water in the proportion of $4\text{ mL}\cdot\text{L}^{-1}$
123 one day after transplanting. Regarding the rice formulation, the volume used by application
124 addition in tubes and in pots, was two milliliters and 150 ml, respectively. At each application
125 addition, a new formulation was produced.
126

127 The inoculation of *M. javanica* occurred was carried out 24 hours after the transplanting of
128 seedlings to the pot, each one receiving received 5 ml of suspension containing 5000 eggs
129 and eventual J2 calibrated in Peters chamber, applied in three holes around the neck roots
130 of each plant. At 45 days after transplanting, the number of galls per gram of root (NG / g),
131 egg mass per gram of root (MO / g), number of eggs per gram of root (NO / g) were


132 evaluated, as well as the reproductive factor, calculated by the following formula: $FR = Pf /$
133 Pi , where Pf is the final nematode population and Pi is the initial population applied added to
134 the plant [16]. and number of second stage juveniles (J2). To count the number of egg
135 masses, roots were immersed in floxin B solution (150 mg.L^{-1}). The number of eggs was
136 determined after root extraction [17, 18]. For number of J2 in the soil, samples 200cm^3 were
137 processed [19]. Eggs and J2 of *M. javanica* were quantified in Peters counting chamber in
138 invert objective microscope.
139

140 The number of leaves, height, head diameter, fresh shoot mass, dry shoot mass and root
141 weight were also evaluated. In order to determine the shoot dry mass, leaves were placed in
142 paper bags, which were placed in a drying oven with forced air circulation at 65°C for 72
143 hours. Data were submitted to analysis of variance and means were compared by the Scott-
144 Knott test at 5% error probability. Statistical analysis was performed using the "Sisvar"
145 software [20].
146

147 3. RESULTS AND DISCUSSION

148 3.1 Development of *Bacillus subtilis*-34 formulation in rice broth and 149 establishment of the growth curve 150

151
152 Figure 1 and 2 show the growth curves of *B. subtilis*-34 in the liquid rice formulation and
153 TSB, respectively. In the liquid rice formulation, it was observed that the bacteria remained in
154 the adaptation phase up to 20 hours after plating. After 24 hours, the exponential growth
155 phase begins, culminating with higher number of CFU (6.14×10^8) at 32 hours after
156 incubation. From 36 hours, the decline phase begins began. The sudden drop in this value is
157 justified by the depletion of nutrients in the culture medium and by the increase of toxic
158 products from bacterial metabolism [21].
159




160
161

162 **Figure 1. Growth curve of *Bacillus subtilis*-34 in liquid rice formulation.**

163
164
165
166
167

In TSB, the highest number of CFUs occurred at 40 hours after incubation (6.41×10^7), that is, eight hours after occurrence in the liquid formulation and with a difference in the number of CFUs of 5.5×10^8 in relation to the liquid rice formulation (Figure 2). For the greenhouse experiment, the formulation was incubated for up to 28 hours because it was already in the logarithmic growth phase. It is important to highlight that 28 hours from the start of

168 incubation, the liquid rice formulation provided CFU of 4.72×10^8 , while in the TSB medium,
 169 CFU was 3.49×10^7 .
 170



171
 172 **Figure 2. Growth curve of *Bacillus subtilis*-34 in TSB medium.**
 173

171
 172
 173

174
 175
 176

177 **3.2 Evaluation of different application addition times of the liquid *Bacillus***
 178 ***subtilis* formulation in rice broth in the control of *Meloidogyne javanica* and**
 179 **lettuce development improvement**

180

181 Lettuce plants that received the liquid *B. subtilis*-34 formulation applied added to the pot and
 182 tube + pot showed lower number of galls per gram of root and lower number of egg mass
 183 per gram of root (Table 1). The number of eggs per gram of root was significantly lower in all
 184 treatments that received the liquid *B. subtilis*-34 formulation compared to Onix® and control.
 185 The reproduction factor of *M. javanica* was lower in treatment irrigation of formulation drench
 186 applied added to the tube followed by application addition in the pot and pot + tube.
 187 Application Addition in tube reduced the reproduction factor of the nematode by 30.55% and
 188 36.07% in relation to Onix® and control, respectively (Table 1). Positive *B. subtilis* results in
 189 reducing nematode populations, mainly of *Meloidogyne* species, in crops plants such as rice,
 190 tomato and banana have been observed in other studies [22, 23, 24]. There was no
 191 occurrence of juveniles of second stage in the soil in any of treatments; probably the J2 that
 192 hatched and infected the roots again.

193

194 **Table 1. Number of galls per gram root (NG/g), number egg mass per gram of**
 195 **root (EM / g), number of eggs per gram of root (NE / g) and reproduction factor**
 196 **(RF) of *Meloidogyne javanica* in lettuce submitted to application of inoculated**
 197 **by *B. subtilis*-34 via liquid formulation at different times.**
 198

Treatments	NG/g	EM/g	NE/g	RF
Pot	8.00a	2.25a	1,686.52a	2.90b
Pot + Tube	8.12a	3.00a	1,562.07a	3.07b

Tube	21.75b	7.75b	2,173.50a	2.41a
Absolute control	36.50c	11.62b	4,784.71b	3.77c
Onix [®]	47.00c	9.62b	5,800.99b	3.47c
Coefficient of variation	41.85	61.54	40.00	15.37

199 Averages followed by the same letter in the column do not differ from each other by the
200 Scott-Knott test at 5% error probability.

201

202 *Bacillus* species interrupt the nematode life cycle through the production of toxic metabolites
203 that restrict their mobility, prevent juvenile hatching and penetration into plant roots [25].
204 Some authors have demonstrated reduction of the *M. incognita* population in tomato
205 inoculated with *Bacillus* species [26], others have observed that metabolites produced by *B.*
206 *subtilis* trigger hypersensitivity reactions in plant cells and affect oviposition, preventing
207 nematode females from obtaining sufficient energy to produce eggs [27].

208

209 The genus *Bacillus* is described as one of the main microbial groups capable of acting in the
210 control of phytopathogens through the synthesis of secondary metabolites, which, in general,
211 present a wide range of inhibition to different phytopathogen species [28]. *Bacillus* secretes
212 many secondary metabolites, including antibiotics, antifungals and siderophores. Metabolites
213 produced by *Bacillus* may also affect the microflora in the rhizosphere, providing an
214 environment antagonistic to pathogens, or may trigger host defense responses [29]. Cry
215 proteins produced by *Bacillus* species are toxic to nematodes, both of free-living and
216 phytoparasites and the production of proteases by this group of bacteria have been
217 proposed as virulence factors in their pathogenesis against nematodes [30].

218

219 *Bacillus subtilis*-34 applied added by irrigation soil drenching and via pot + tube promoted
220 higher number of leaves, head diameter and dry shoot mass significantly higher than
221 applications additions in tube, control and Onix[®] commercial product (Table 2). Applications
222 by Additions in pot and pot + tube increased the number of leaves compared to control by
223 about 80.02 and 73.56%. In relation to Onix[®], the increase was 83.52 and 76.94%. Variable
224 head diameter increased 81.98 and 75.42% when applying the formulation in pot and pot +
225 tube, respectively, in relation to control; when compared with Onix[®], the increase was 94 and
226 87.73%. In variable dry shoot mass, considering applications in pot and pot + tube, the
227 increase was 21.74 and 17.94% in relation to control, and in relation to Onix[®], the increase
228 was 23.38 and 19.54%.

229

230 The height and fresh shoot mass of lettuce plants that received *B. subtilis*-34 in the pot were
231 significantly higher than the other treatments, with an increase of about 119 and 322.22%,
232 respectively, in relation to control. On the other hand, application addition via pot and pot +
233 tube provided an increase of 98.61 and 155% of root weight in relation to control and Onix[®]
234 commercial product. The *in situ* effect by exposure of *B. subtilis* living cells may also lead to
235 an increase in plant biometry [31], reflecting productivity gains, with the bacterium being
236 commercially used for both purposes [32, 33].

237

238 **Table 2. Number of leaves (NL), height (H), head diameter (HD), fresh shoot**
239 **matter (FSM) (g), shoot dry matter (SDM) (g) and fresh root matter (FRM) (g) of**
240 **lettuce infected by *Meloidogyne javanica* and submitted to application of**
241 ***Bacillus subtilis*-34 via liquid rice formulation at different times.**

242

Treatments	NL	H (cm)	HD(cm)	FSM(g)	SDM(g)	FRM(g)
Pot	34.87a	7.12a	41.62a	76.00a	11.87a	8.62b

Pot + Tube	33.62a	6.25b	40.12a	64.12b	11.50a	10.50a
Tube	26.25b	5.00c	28.12b	23.50c	9.87b	6.25c
Absolute control	19.37c	3.25c	22.87c	18.00c	9.75b	4.37d
Onix [®]	19.00c	3.62c	21.37c	17.12c	9.62b	3.37d
Coefficient of variation	11.39	15.32	5.53	22.235	9.76	25.41

243 Averages followed by the same letter in the column do not differ from each other by the
244 Scott-Knott test at 5% error probability.

245

246 *Bacillus subtilis* has been commercially used for the biocontrol of plant diseases and to
247 increase crop yields [32, 33]. *B. subtilis* (PRBS-1) applied added to tomato plants reduced
248 the reproduction of root-knot nematode and promoted the growth of plants under
249 greenhouse conditions [34]. Nemathel[®] treated banana seedlings reduced reproduction of
250 *Radopholus similis*, *Meloidogyne* spp., *Pratylenchus* ssp. and *Helicotylenchus* spp. with
251 efficiency similar to nematicide Carbofuran [35]. Tomato plants that received *B. subtilis*
252 applications additions showed higher shoot growth, characterizing the bacterium as a plant
253 growth promoter, and this effect may be due, in part, to the production of plant
254 phyto regulators in the rhizosphere [36].

255

256 3.3 Evaluation of the viability of liquid *Bacillus subtilis* rice formulation under 257 room and refrigerator conditions.

258

259 The initial CFU was 4.72×10^8 . Over the storage period under both conditions, oscillations in
260 the concentration of bacterial cells were observed, sometimes with higher and sometimes
261 with smaller values (Figure 3). At 4 months, the number of CFU of formulation stored under
262 room and refrigerator conditions was 12.5×10^8 and 6.3×10^8 , respectively. At 6 months, the
263 number of viable cells was similar in both storage conditions 7.7×10^8 at room temperature
264 and 6.7×10^8 at refrigerator temperature. At 7 months, it was verified that the number of
265 CFUs remained approximately constant 7.8×10^8 under room conditions, while under
266 refrigeration conditions, reduction to 3.8×10^8 UFC was observed, and this value remained
267 approximately constant until 9 months.

268

269 From 8 months, the number of CFUs stored under room conditions becomes minimal, while
270 in refrigerator, the number of CFUs is was approximately 4×10^8 at 9 months. The
271 refrigerated environment extended the "shelf life" of the bacterium in two months. Low
272 temperatures are generally used to preserve microorganisms by ensuring metabolism at low
273 activity and preventing contamination with other microorganisms from affecting the stability
274 of the biological control microorganism [37].


275

276 A biological control product to be economically viable needs to have minimum concentration
277 of 1×10^8 CFU / mL with 85% viability [38], which was achieved by the liquid *B. subtilis*-34
278 formulation stored under room conditions for up to seven months and under refrigerator
279 conditions for up to nine months.

280

281

282



283 **Figure 3. Number of CFU of *Bacillus subtilis*-34 in liquid formulation stored under**
284 **room and refrigerator conditions for twelve months.**
285

286 The results verified in the nematode control and in the promotion of lettuce growth
287 demonstrate that the liquid bacterium formulation was effective despite the lower initial
288 number of CFU.mL⁻¹ (4.72x10⁸) compared to Onix, which had 1x10⁹ CFU. It also presented
289 lower cost when compared to TSB synthetic culture medium since US\$ 128.00 are
290 necessary for the production of one liter of TSB, whereas the liquid rice formulation requires
291 only US\$ 11.64.
292

293 **4. CONCLUSION**
294

295 Higher promotion of lettuce growth and control of *Meloidogyne javanica* was were obtained
296 by applying adding the liquid *Bacillus subtilis*-34 formulation twice to the soil in the pot.
297 *Bacillus subtilis*-34 remains viable until nine months in formulation stored under refrigerator
298 conditions and up to seven months under room conditions in northern state of Minas Gerais.
299

300 **REFERENCES**
301

- 302
- 303 1. Carvalho Filho JLS de, Gomes LAA, Maluf WR. Tolerância ao florescimento precoce e
304 características comerciais de progênies F4 de alface do cruzamento Regina 71 x Salinas
305 88. Acta Scientiarum Agronomy.2009; 31: 37-42. Portuguese.
 - 306 2. Matraszek R, Nowak BH, Chwil S, Chwil M. Macroelemental composition of cadmium
307 stressed lettuce plants grown under conditions of intensive sulphur nutrition.2016; 180:24-
308 34.
 - 309 3. Villas Bôas RL, Passos JC, Fernandes M, Büll LT, Cezar VRS, Goto R. Efeito de doses e
310 tipos de compostos orgânicos na produção de alface em dois solos sob ambiente protegido.
311 Horticultura Brasileira.2004; 22:28-34. Portuguese.
 - 312 4. Anwar SA, Mckenry MV. Incidence and population density of plant parasitic nematodes
313 infecting vegetable crops and associated yield losses. Pakistan Journal of Zoology.2012;
314 44: 327-333.
 - 315 5. Fiorini CVA, Gomes LAA, Libânio RA, Maluf WR, Campos VP, Licursi V, et al.
316 Identificação de famílias F2:3 de alface homozigotas resistentes aos nematoides das
317 galhas. Horticultura Brasileira. 2007; 25: 509-513. Portuguese.
 - 318 6. Pinheiro JB, Pereira RB, Carvalho ADF, Rodrigues CS, Suinada FA. Manejo de
319 nematoides na cultura do alface. 1ed. Embrapa: Brasília .2013. Portuguese.
 - 320 7. Killani AS, Abaidoo RC, Akintokun AK, Abiala MA. Antagonistic Effect of Indigenous
321 *Bacillus subtilis* on Root-/Soil-borne Fungal Pathogens of Cowpea. Researcher.2011; 3:11-
322 18.
 - 323 8. Tian B, Yang J, Lian L, Wang C, Li N, Zhang KQ. Role of an extracellular neutral protease
324 in infection against nematodes by *Brevibacillus laterosporus* strain G4. Applied Microbiology
325 and Biotechnology.2007; 74: 372-380.
 - 326 9. Formstone A, Carballido-Lopez P, Noirot J, Errington D, Scheffers J. Localization and
327 interaction of teichoic acid synthetic enzymes in *Bacillus subtilis*. Journal
328 Bacteriology.2008; 190:1812-1821.
 - 329 10. Ribeiro RCF, Campos VP, Xavier AA, Rocha LS, Ribeiro HB, Aguiar FM, et al.
330 Rizobactérias no controle de *Meloidogyne javanica* e mal do Panamá em bananeira.
331 Nematropica. 2012; 42: 218-226. Portuguese.
 - 332 11. Sharma A, Meena KR, Kanwar SS. Molecular characterization and bioinformatics studies
333 of a lipase from *Bacillus thermoamylovorans* BHK67. International Journal of Biological
334 Macromolecules. 2018; 107:2131-2140.

- 335 12. Lee YS, Kim KY. Antagonistic potential of *Bacillus pumilus* L1 against root-Knot
336 nematode, *Meloidogyne arenaria*. Journal Phytopathology.2016; 164:29-39.
- 337 13. Zhang L, Yu J, Xie Y, Lin H, Huang Z, Xu L, et al. Biological activity of *Bacillus*
338 *thuringiensis* (Bacillales: Bacillaceae) chitinase against *Caenorhabditis elegans* (Rhabditida:
339 Rhabditidae). Journal of Economic Entomology. 2014; 107:551-558.
- 340 14. Berninger T, López OG, Bejarano A, Preininger C, Sessitsch A. Maintenance and
341 assessment of cell viability in formulation of no-sporulating bacterial inoculants. Microbial
342 Biotechnology. 2018; 11:277–301.
- 343 15. Sharma RR, SINGH D, Singh R. Biological control of postharvest diseases of fruits and
344 vegetables by microbial antagonists: A review. Biological Control. 2009; 50:205-211.
- 345 16. Oostenbrink M. Major characteristic of the relation between nematodes and plants.
346 Mededlingen van de Landbouwhogeschool;1966.
- 347 17. Hussey RS, Barker KA. A comparison of methods of collecting inocula for *Meloidogyne*
348 spp. Including a new technique. Plant Disease Reporter. 1973;57:1025-1028.
- 349 18. Boneti JIS, Ferraz S. Modificação do método de Hussey e Barker para extração de ovos
350 de *Meloidogyne exigua* de café. Fitopatologia Brasileira.1981; 6: 553. Portuguese.
- 351 19. JENKINS WR A rapid centrifugal-flotation technique for separating nematodes from soil.
352 Plant Disease Reporter.1964; 48: 692.
- 353 20. Ferreira DF. SISVAR: Um programa para análises e ensino de estatística. Revista
354 Symposium. 2008; 6: 36-41.Portuguese.
- 355 21. Instituto Superior de Ciências da Saúde do Sul. Manual Prático de microbiologia. 2018.
356 Acessado em 07 de julho de 2018.
357 Disponível:<http://www.codigopostal.cibeforma.pt/dir/0/instituto-superior-de-ciencias-da-saude-sul/>.
358
- 359 22. Ludwigi J, Mourão AB, Gomes CB. Potencial da microbiolização de sementes de arroz
360 com rizobactérias para o biocontrole do nematoide das galhas. Tropical Plant
361 Pathology.2013; 38: 264-268. Portuguese.
- 362 23. Fernandes RH, Vieira BS, Fuga CAG, Lopes EA. *Pochonia chlamydosporia* e *Bacillus*
363 *subtilis* no controle de *Meloidogyne incognita* e *M. javanica* em mudas de tomateiro.
364 Bioscience Journal. 2014; 30:194-200. Portuguese.
- 365 24. Araújo JJ S, Muniz MFS, Filho GM, Rocha FS, Castro JMC. *Bacillus subtilis* no
366 tratamento de mudas de bananeira infectadas por fitonematoides. Revista Ceres. 2018; 65:
367 99-103. 2018. Portuguese.
- 368 25. Kavitha J, Jonathan EI, Umamaheswari R. Field application of *Pseudomonas*
369 *fluorescens*, *Bacillus subtilis* and *Trichoderma viride* for the control of *Meloidogyne incognita*
370 (Kofoid and White) Chitwood on sugarbeet. Journal of Biological Control. 2007; 21:211-215.
- 371 26. Silva JO, Santana MV, Freire LL, Ferreira BS, Rocha MR. Biocontrol agents in the
372 management of *Meloidogyne incognita* in tomato. Ciência Rural.2017; 47:1-7.
- 373 27. Sharma RD, Gomes AC. Effects of *Bacillus* spp. Toxins on oviposition and juvenile
374 hatching of *Heterodera glycines*. Nematologia Brasileira. 1996; 20: 53-62.
- 375 28. Sansinenea E, Ortiz A. Secondary metabolites of soil *Bacillus* spp. Biotechnology
376 Letters. 2011; 33: 1523–1538.
- 377 29. Velusamy P, Gnanamnickam SS. The Effect of bacterial secondary metabolites on
378 bacterial and fungal pathogens of rice. Em: KARLOVSKY, P, editor. Secondary Metabolites
379 in Soil Ecology .14ª ed. Berlin: Springer; 2008.
- 380 30. Lian LH, Tian BY, Xiong R, Zhu MZ, Xul J, Zhang KQ. Proteases from *Bacillus*: a new
381 insight into the mechanism of action for rhizobacterial suppression of nematode populations.
382 Letters in Applied Microbiology. 2007; 45: 262–269.
- 383 31. Hammami I, Rhouma A, Jaodu B, Rebai A, Nesme X. Optimization and biochemical
384 characterization of a bacteriocin from a newly isolated *Bacillus subtilis* strain 14 B for
385 bioncontrol of *Agrobacterium* spp. strains. Letters in Applied Microbiology. 2009; 48: 253-
386 260.

- 387 32. Ngugi HK, Dedej S, Delaplane KS, Savelle AT, Scherm H. Effect of flower-applied
388 Serenade biofungicide (*Bacillus subtilis*) on pollination-related variables in rabbiteye
389 blueberry. *Biological Control*. 2005; 33: 32-38.
- 390 33. Yao BH, Karimov S, Boturov U, Sanginboy S, Sharipov A. Effect of FZB 24R *Bacillus*
391 *subtilis* as a biofertilizer on cotton yields in field tests. *Archives of Phytopathology and Plant*
392 *Protection*. 2006; 39: 323-328.
- 393 34. Araújo FF, Marchesii GVP. Uso de *Bacillus subtilis* no controle da meloidoginose e na
394 promoção do crescimento do tomateiro. *Ciência Rural*. 2009; 39: 1558-1561. Portuguese.
- 395 35. Araújo JJ S, Muniz MFS, Filho GM, Rocha FS, Castro JMC. *Bacillus subtilis* no
396 tratamento de mudas de bananeira infectadas por fitonematoides. *Revista Ceres*. 2018; 65:
397 99-103. 2018. Portuguese.
- 398 36. Araujo FF, Bettiol W. Supressividade dos nematoides *Meloidogyne javanica* e
399 *Heterodera glycines* em soja por adição de lodo de esgoto ao solo. *Ciência Rural*. 2005; 35:
400 806-812. Portuguese.
- 401 37. Alves LFA, Alves SB, Lopes RB, Augus NT. Estabilidade de uma formulação de *Bacillus*
402 *sphaericus* armazenada sob diferentes temperaturas. *Scientia Agricola*. 2001; 58: 21-26.
403 Portuguese.
- 404 38. D` agostino F, Morandi MAB. Análise da viabilidade comercial de produtos à base de
405 *Bacillus subtilis* e *Bacillus pumilus* para controle de fitopatógenos no Brasil. EM: Bettiol W,
406 Morandi MAB, editores. *Biocontrole de doenças de plantas: Uso e perspectivas*. 1ª ed.
407 Jaguariúna: Embrapa Meio Ambiente. 2009. Portuguese