

**Litter in potential eucalypts genotypes implanted in Eldorado do Sul, Rio Grande do Sul, Brazil**

**ABSTRACT:** The objective of this study was to estimate the amount of litter on the soil in genotypes of *Eucalyptus* at 49-month-old, located in Eldorado do Sul, Rio Grande do Sul, Brazil. For each genotype, a sampling unit of 720 m<sup>2</sup> was demarcated, where in each of them 15 random collections were carried out. The litter biomass ranged from 4.51 to 10.77 Mg ha<sup>-1</sup>, highlighting the *E. dunnii* and the hybrid of *E. urophylla* x *E. globulus* with the lowest and largest production respectively. The leaves corresponded, on average, between 48.56% and 73.03% of the total litter. The differentiation between the genotypes occurred as to the accumulated litter production.

**Keywords:** Litter biomass, Leaves, *Eucalyptus* stands, Sustainability.

## **1. INTRODUCTION**

Tree plantations have many beneficial ecological interactions with the ecosystem, such as watershed protection, increased organic matter and soil nutritional status through the production of litter [2]. This dynamic, represented by the deposition of litter via deposition and exit via decomposition is essential to the maintenance of forests or forest stands [3], especially in highly weathered soils, where plant biomass is the main nutrient reservoir [10].

The persistence of litter on the soil causes this material to be reused in the nutrient cycle of the system, through its decomposition and the release of the constituent

minerals for a later reabsorption by the roots of the plants, also increasing the organic matter content in the soil [4]. In this context, nutrient return via litter is the most important route of the biogeochemical cycle [10].

In general, an increase in litter deposition is observed until the age at which the trees reach maturity, when the crowns are closed. After this phase, a slight decrease or stabilization in the deposition may be observed [9].

The accumulation of litter varies according to the origin, species, forest cover, successional stage, age, collection season, forest type and site. In addition to these factors, others such as, edaphoclimatic conditions and water regime, climatic conditions, site, understory, silvicultural management, proportion of canopy, as well as rate of decomposition and natural disturbances such as fire and insect or artificial attack such as litter removal and crops, occurring in the forest or in the stand, also influence the accumulation of litter [5].

The litter deposition is higher in the period of greater physiological activity of the individuals, causing an intensification of foliage exchange and senescent material release to give rise to a new and photosynthetically more active foliage [13].

Knowledge of the amount of litter deposited in different eucalyptus species and provenances is of fundamental importance in order to maintain a sustainable management of soils and mineral resources [12].

The present study had as objective to estimate the biomass of litter in different *Eucalyptus* genotypes established in Eldorado do Sul, Rio Grande do Sul, Brazil.

## 2. MATERIALS AND METHODS

The research was developed with different genotypes of eucalypts (Table 1), in an area belonging to the company Celulose Riograndense - CMPC, in the city of Eldorado do Sul, RS, Brazil (Figure 1). The area is under the geographic coordinates of 30 ° 11'303 "south latitude and 51 ° 37'477" west longitude.

The climate is characterized as subtropical humid (Cfa), according to the climatic classification of Köppen, presenting average temperature corresponding to 19 °C. The average annual precipitation of 1,400 mm [1].

The soil in the area is classified as Red-Yellow Argisol. Table 2 presents the chemical and physical attributes of the soil at depths from 0 to 130 cm.

Table 1. Characterization of the studied genotypes

Genotypes	Age	Spacing	G
	(months)	(m)	(m <sup>2</sup> ha)
<i>E. benthamii</i> (Provenance 1)	49	3 x 3	24,4
<i>E. benthamii</i> (Provenance 2)	49	3 x 3	22,7
<i>E. saligna</i>	49	3 x 3	23,7
<i>E. dunnii</i>	49	3 x 3	16,7
<i>E. urophylla</i> x <i>E. globulus</i> ( <i>E. uroglobulus</i> )	49	3 x 3	22,2
<i>E. urophylla</i> x <i>E. grandis</i> ( <i>E. urograndis</i> )	49	3 x 3	26,4

*E. benthamii* (Provenance 1) is a provenance proven in Guarapuava, Paraná, Brazil; and *E. benthamii* (Provenance 2) is a source from Telêmaco Borba, Paraná, Brazil.

Figure 1. Location of the municipality of Eldorado do Sul, Rio Grande do Sul, Brazil




Table 2. Physical and chemical attributes of soil in the area implanted with different genotypes of *Eucalyptus*, at 49-months-old, in Eldorado do Sul, Rio Grande do Sul, Brazil

Granulometric composition					
Depth	Coarse sand	Thin sand	Silt	Clay	O.C
	2-0.2	0.2-0.05	0.05-0.002	<0.002	
(cm)	-----mm-----				%
0-30	24.5	16.5	29.5	29.5	0.9
30-60	40.5	8.0	6.0	45.5	0.8
60-90	33.5	6.0	5.5	55.0	0.7
90-100	15.5	6.0	15.5	63.0	0.4
100-130	15.5	6.5	13.0	65.0	0.2

Depth	V	m	T	pH	N
(cm)	-----%		--cmol <sub>c</sub> dm <sup>-3</sup> --	H <sub>2</sub> O	%
0-30	35	34	10	5.0	0.1
30-60	11	71	14	4.3	0.1
60-90	15	69	15	4.4	0.1
90-100	17	64	12	4.6	0.1
100-130	20	61	10	4.7	0.0

Depth	P	K	Ca	Mg	S
(cm)	--mg g <sup>-1</sup> --	-----cmol <sub>c</sub> dm <sup>-3</sup> -----			mg dm <sup>3</sup>
0-30	2.0	0.1	3.3	0.9	19.4
30-60	1.6	0.1	0.9	0.5	32.5
60-90	1.0	0.1	1.0	0.8	61.7
90-100	0.7	0.1	1.0	0.9	60.9
100-130	0.6	0.1	0.9	0.9	59.0

72 O.C: organic carbon; V = saturation by bases; m = saturation by aluminum; T = total

73 CTC

74

75 In the preparation of the area, the subsoiling was performed at a depth of 60 cm,  
76 and a liming treatment was applied consisting of 2 Mg ha<sup>-1</sup> of limestone, and 200 kg ha<sup>-1</sup>  
77 of single superphosphate. The fertilizer used during planting consisted of, 110 g plant<sup>-1</sup>  
78 of N-P<sub>2</sub>O<sub>5</sub>-K<sub>2</sub>O (06:30:06) + 0.3% Zn and 0.2% Cu. For coverage fertilization 200 kg  
79 ha<sup>-1</sup> of N-P<sub>2</sub>O<sub>5</sub>-K<sub>2</sub>O (12:00:20) + 0.7% of B were applied, and for the maintenance  
80 fertilization, 300 kg ha<sup>-1</sup> of N-P<sub>2</sub>O<sub>5</sub>-K<sub>2</sub>O (24:00:26) + 0,5% B were applied.

81       The litter collections were carried out in June 2016. To perform the work, in each  
82 of the genotypes studied, a plot of 720 m<sup>2</sup> was demarcated, where 15 samples were  
83 collected, randomly, totaling 90 samples.

84       The litter samples were collected using an iron frame (Figure 2) of 0.25 m x 0.25  
85 m (0.0625 m<sup>2</sup>), which was placed on the surface of the land, and all the organic material  
86 present in its soil. After collection, the sampled materials were stored in plastic bags and  
87 sent to the laboratory where they were separated into three fractions: leaves, branches  
88 and miscellaneous (peels, reproductive materials and non-identifiable residues).

89       Subsequently, the fractions were placed in paper containers to dry in a circulation  
90 oven and air renovation at 70 °C until weight stabilization. Finally, the samples were  
91 weighed in a precision scale (0.01 g) and the dry weights were extrapolated to values  
92 per hectare to obtain the mass of the litter.

93

94

95

96

97 Figure 2. A: Canopy of a stand of *Eucalyptus* sp.; Deposition of the litter on the soil;  
98 and C: Removal of the litter on the soil for quantification.



111 Statistical analyzes were performed with the aid of the statistical program Assistat  
112 7.7 ® [14] at the level of 5% probability of error. The Tukey test was used for the  
113 comparison of means.

### 116 3. RESULTS AND DISCUSSIONS

117 In Table 3 it is possible to verify the distribution of litter for the different fractions  
118 of the studied genotypes. The litter biomass was higher in hybrid *E. urophylla* x *E.*  
119 *globulus* (10.77 Mg ha<sup>-1</sup>) and lower in *E. dunnii* (4.51 Mg ha<sup>-1</sup>), reaching intermediate  
120 values in *E. benthamii* (P1) (8, 38 Mg ha<sup>-1</sup>).

Table 3. Litter production for the different fractions of the eucalypts genotypes at 49-  
months-old

Genotypes	Leaves	Miscellaneous	Branches	Total
	(Mg ha <sup>-1</sup> )			
<i>E. benthamii</i> (P1)	6.12ab *(2.15)	0.53 <sup>a</sup> (0.45)	1.73bc (1.74)	8.38a (3.31)
<i>E. benthamii</i> (P2)	3.27cd (1.41)	0.65 <sup>a</sup> (0.61)	1.14c (0.57)	5.06b (1.64)
<i>E. saligna</i>	4.63bc (1.26)	0.43 <sup>a</sup> (0.25)	4.10a (2.59)	9.13a (3.04)
<i>E. dunnii</i>	2.19d (1.24)	0.95 <sup>a</sup> (1.17)	1.46bc (1.35)	4.51b (2.81)
<i>E. urophylla</i> x <i>E. globulus</i>	6.72a (2.93)	0.32 <sup>a</sup> (0.27)	3.82a (2.08)	10.77a (3.95)
<i>E. urophylla</i> x <i>E. grandis</i>	5.88ab (1.65)	0.53 <sup>a</sup> (0.44)	3.11ab (2.03)	9.49a (2.87)
CV %	38.97	108.2	71.67	38.26

CV: Coefficient of variation.

Mean of each variable in the different treatments followed by equal letters, do not differ significantly by the Tukey test at the 5% level of error. \* Values in parentheses are the standard deviation of the mean.



129 The litter presented a value lower than that found by [15], in a plantation of hybrid  
130 *E. urophylla* x *E. globulus* at four-years-old in Eldorado do Sul, RS, Brazil (14.0 Mg ha<sup>-1</sup>);  
131 by [12], in stands of *E. grandis*, *E. cloesiana* and *E. urophylla*, with nine-years-old in  
132 Santa Maria, RS, Brazil (16.8, 16.5 and 12.6 Mg ha<sup>-1</sup>); by [11] in a stand of *E. saligna* at  
133 four and five years of age in São Gabriel, RS, Brazil (12.76 and 12.00 Mg ha<sup>-1</sup>); and by  
134 [4] in a planting of *E. uroglobulus*, with 5.5-years-old in Eldorado do Sul, RS, Brazil  
135 (19.5 Mg ha<sup>-1</sup>).

136 In another stand of *E. grandis*, at seven-years-old, in Seropédica, RJ, Brazil, [3],  
137 reported an amount of 11.84 Mg ha<sup>-1</sup> litter. In the Southwest region of Brazil, in  
138 different forest sites, [7] evaluating litter production in eucalypts plantations, also at  
139 seven-years-old, observed that the amount of litter ranged from 4.2 to 37.6 Mg ha<sup>-1</sup>. A  
140 similar result was found by [6], in a stand of *E. grandis*, at the nine-years-old, in the city  
141 of Alegrete, RS, Brazil (5.41 Mg ha<sup>-1</sup>).

142 The production and accumulation of litter presents a great variability, which can  
143 be due to variation in the climatic conditions, the quality of the site, the age of the stand,  
144 the characteristics of the species, as well as in the inclusion of the understory litter, and  
145 the degree of forest stability [10].


146 In relation to total biomass, the hybrid *E. urophylla* x *E. globulus* presented a  
147 production of 58.12% and 53.02% higher than the clones *E. dunnii* and *E. benthamii*  
148 (P2). According to [6], the litter rates accumulated in forest plantation soils can vary  
149 significantly between different species in the same sites.

150 The leaves fraction corresponded on average, between 48.56% and 73.03% of the  
151 total litter (Figure 3). *E. dunnii* was the clone that presented the lowest percentage of  
152 leaves deposited among the species studied, and *E. benthamii* (P1) the largest. The

153 leaves usually constitute the largest proportion of the biomass of the residues that fall to  
 154 the soil. This proportion grows with age to a certain extent, and then decreases due to  
 155 the increase in the fall of branches and bark [10].

156

157 Figure 3. Relative litter of different *Eucalyptus* genotypes



158

159 In this context, [8] studied the litter in a twelve-year-old eucalypts stand, observed  
 160 that the branches presented greater accumulation on the ground, with 38.8%. According  
 161 to the author, this occurred because the eucalypts stands suffer a more intense process of  
 162 natural desrama, in relation to other species, which leads to greater accumulation of  
 163 branches on the ground.

164 [12], studying the litter in three species of eucalypts (*E. urophylla*, *E. cloesiana*  
 165 and *E. grandis*), at nine-year-old, in Santa Maria, RS, Brazil, observed that the branches  
 166 fraction was the most representative in all species. According to the authors this may  
 167 mean a propensity of these species for the natural desrama.

168 For the branches fraction, the highest yields occurred in the clones *E. saligna* and  
169 *E. uroglobulus*, with 2.48 and 2.24 Mg ha<sup>-1</sup>, and lower production occurred in *E. dunnii*,  
170 with 2.19 Mg ha<sup>-1</sup>. The miscellaneous fraction varied from 0.32 to 0.95 Mg ha<sup>-1</sup>, the  
171 lowest value found in hybrid *E. urophylla* x *E. globulus* and the highest in *E. dunnii*, not  
172 statistically different from the other clones.

173 The existence of a great variability in the accumulation of organic litter blanket in  
174 relation to other works carried out with eucalypts species can be explained based on the  
175 variation of climate, sites, age and forest density, different genetic characteristics of  
176 each species and the stability achieved by the stand, depending on the time elapsed  
177 since the last intervention that may have influenced the litter. These factors will affect  
178 the balance between the amount of material deposited and the time required for its  
179 decomposition, reflecting the greater or less accumulation of litter in the forest soil and  
180 the percentage composition of the different fractions that compose it [12].

181

#### 182 4. CONCLUSIONS

183 The highest production of litter occurred in hybrid *E. urophylla* x *E. globulus*,  
184 providing greater protection to the soil, in contrast the lower production of litter  
185 occurred in *E. dunnii*.

186 The leaf fraction presented the highest contribution in all genetic materials  
187 studied.

188 The differentiation between the genotypes occurred as to the accumulated litter  
189 production.

190

191

192

193

194 **COMPETING INTERESTS DISCLAIMER:**

195

196 Authors have declared that no competing interests exist. The products used for this  
197 research are commonly and predominantly use products in our area of research and  
198 country. There is absolutely no conflict of interest between the authors and producers  
199 of the products because we do not intend to use these products as an avenue for any  
200 litigation but for the advancement of knowledge. Also, the research was not funded by  
201 the producing company rather it was funded by personal efforts of the authors.

202

203

204 **6. BIBLIOGRAPHIC REFERENCES**

205 1. Alvares CA, Stape JL, Sentelhas PC, Gonçalves JLM, Sparovek G. Köppen's climate  
206 classification map for Brazil. Meteorologische Zeitschrift. 2013; 22 (6): 1-18.

207

208 2. Ashagrie Y, Zech W. Litter production and nutrient cycling in two plantations and  
209 Podocarpus falcatus dominated natural forest ecosystems in south-eastern highlands of  
210 Ethiopia. African Journal of Agricultural Research. 2013; 8 (38): 4810-4818.

211

212 3. Balieiro FC, Franco AA, Pereira MG, Campello EFC, Dias LE, Faria SM et al. Litter  
213 dynamics and nitrogen transfer to the soil, in plantations of Pseudosamanea guachapele  
214 and Eucalyptus grandis. Pesquisa Agropecuária Brasileira. 2004; 39 (6): 597-601.

215

216 4. Brun EJ, Ferraz MO, Araújo EF. Relation between soil litter accumulation and  
217 dendrometric variables in hybrid Eucalyptus urophylla x E. globulus Maidenii stands, in  
218 Eldorado do Sul / RS. Revista Ecologia e Nutrição Florestal. 2013; 1 (1): 24-31.

219

- 220 5. Boiler MVW. Quantification of litter and nutrients - Mixed Ombrophylous Forest  
221 Montana - Paraná. Academic Review. 2007; 5 (2): 101-116.  
222
- 223 6. Freitas R, Schumacher MV, Boiler MVW, Spathelf P. Biomass and nutrient content  
224 in *Eucalyptus grandis* stands W. Hill ex Maiden planted in soil subject to sand blasting,  
225 in the municipality of Alegrete-RS. Biomass & Energy. 2004; 1 (1): 93-104.  
226
- 227 7. Gama-Rodrigues EF, Barros NF, Viana AP, Santos, GA. Changes in biomass and  
228 microbial activity of litter and soil, due to the replacement of native forest cover by  
229 eucalyptus plantations, in different sites of the Southeastern Region of Brazil. Brazilian  
230 Journal of Soil Science. 2008; 32 (4): 1489-1499.  
231
- 232 8. Kleinpaul IS, Schumacher MV, Brun EJ, Brun FRK, Kleinpaul JJ. Sample  
233 sufficiency for accumulated litter collection on the soil in *Pinus elliottii* engelm,  
234 *Eucalyptus* sp. and Seasonal Deciduous Forest. Tree Review. 2005; 29 (6): 965-972.  
235
- 236 9. Poggiani F, Schumacher MV. Nutrient cycling in native forests. In: Gonçalves JLM,  
237 Benedetti V. (Ed.). Forest nutrition and fertilization. Piracicaba; 2004.  
238
- 239 10. MGF Kings, Barros NF. Nutrient cycling in eucalyptus plantations. In: Barros NF,  
240 Novais RF. (Eds.). Relationship between soil and eucalyptus. Viçosa; nineteen ninety.  
241

- 242 11. Santos JC, Schumacher MV, Witschoreck R, Araújo EF, Lopes VG. Nutrients in  
243 litter accumulated in a settlement of *Eucalyptus saligna* Smith in São Gabriel, RS.  
244 *Revista Ecologia e Nutrição Florestal*. 2014; 2 (1): 1-8.
- 245
- 246 12. Schumacher MV, Bauermann, GC, Copetti L, Brun EJ, König F. Litter fractionation  
247 in three eucalyptus species in the municipality of Santa Maria - RS: *Eucalyptus*  
248 *urophylla*, *Eucalyptus cloesiana* and *Eucalyptus grandis*. In: CONESUL forest update  
249 cycle. Santa Maria; 2002.
- 250
- 251 13. Schumacher MV, Brun EJ, Rodrigues LM, Santos EM. Return of nutrients via  
252 deposition of litter in a settlement of *Acácia* in Rio Grande do Sul. *Revista Árvore*.  
253 2003; 27 (6): 791-798.
- 254
- 255 14. Silva FAS, Azevedo CAV. Main components analysis in the software assistat  
256 statistical attendance. In: Word congress on computers in agriculture 7, Reno-NV-USA:  
257 American Society of Agricultural and Biological Engineers, 2009.
- 258
- 259 15. Viera M, Schumacher MV, Boiler MVW. Decomposition and Nutrient Dynamics in  
260 Planting of *Eucalyptus urophylla* × *Eucalyptus globulus* in Southern Brazil. *Forest and*  
261 *Environment*. 2013; 20 (3); 351-360.

262

263

264