

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32

3  
4

5  
6  
7  
8  
9  
10  
11  
12  
13  
14

15  
16  
17  
18  
19  
20  
21  
22  
23  
24

25

26

27

28  
29  
30  
31  
32

33 Nigeria where it is grown as sole crop or as an intercrop with other crops such as root crops, grain  
34 legumes, cereals (sorghum, millet, rice) and even vegetables (Gwinner *et al.*, 1996). Despite the  
35 importance and wide cultivation of this cereal, its growth and productions are usually impaired by  
36 diseases caused by fungal, bacterial, viral or nematode which results in considerable yield loss and  
37 decrease grain quality (Akande and Lamidi, 2006).

38 Fungi were ranked as the second most important cause of maize diseases and the major genera  
39 commonly encountered on maize in tropical regions are *Fusarium*, *Aspergillus* and *Penicillium*  
40 (Ominski *et al.*, 1994; Orsi *et al.*, 2000). *Fusarium* is considered as the most devastating fungal  
41 menace of maize while *Fusarium verticillioides* is the prevalent specie causing root rot, stalk rot and  
42 ear rots in maize (Masuka *et al.* 2003; Alankoya *et al.*, 2008). Meanwhile, the severe economic  
43 losses in corn and other cereal crops worldwide, in addition to the potential occurrence of  
44 fumonisins and other mycotoxins in consequence of *F. verticillioides* infection are a matter of  
45 concern in current mycotoxicology (Ono *et al.* 2010).

46 In the recent times, the use of fungicide which is a chemical control measures commonly employed  
47 in the management of diseases caused by fungi have been discouraged due to its environmental and  
48 food contamination (Akanmu *et al.* 2013). Hence, there is need to embrace a safer and novel  
49 biological approach to plant disease management. In order to improve plants' resistance to diseases  
50 effort has been directed at the search for new anti-microbial materials from natural sources, which  
51 are mostly low-molecular weight secondary metabolites essential for plant disease resistance (Singh  
52 *et al.*, 2010; Maddox *et al.* 2010). While accumulation of certain plant secondary metabolites such as  
53 phytoalexins is induced upon pathogen attack, the increase accumulation of phenolic phytoalexins in  
54 plants can promote host defense against pathogens (Boudet, 2006; Maddox *et al.* 2010). More so,  
55 phenolic compounds with less complex structures, such as catechol and coumarin, have exhibited  
56 bactericidal and fungicidal activities (Cowan, 1999). Hence, the ability of plants to release phenol as  
57 a defense mechanism against infection by pathogens is of importance in determining its resistance  
58 (Samapundo, 2007). This study therefore investigates the phenol production by five maize varieties  
59 in response to infection caused by *Fusarium verticillioides*

## MATERIALS AND METHODS

**Experimental Site:** The experiment was conducted in the screen house of the Department of Botany University of Ibadan, Ibadan, Nigeria.

**Sources of materials used:** Seeds of maize varieties; SWAM1-SR, BR-9928-DMR-SR, ART-98-SW1, ART-98-SW6, BR-9943-DR-SR were obtained from maize germplasm of the Institute of Agricultural and Research Training (IAR&T), Apata, Ibadan. Pure culture of the characterised *Fusarium verticillioides* isolate was obtained from the Plant Pathology Laboratory, International Institute of Tropical Agriculture (IITA), Ibadan, Nigeria.

**Multiplication of the inoculum:** *Fusarium verticillioides* isolate was sub-cultured, maintained on solidified potato dextrose agar (PDA) and incubated at room temperature ( $28\pm 2$ ) °C for 7 days to produce multiple cultures of the organism. Each of the cultured plates was flooded with 2ml sterile distilled water. A sterile scalpel was used to harvest the spores and mycelia growths by scraping them into a sterilized flask. The solution was adjusted with sterile distilled water, stirred and filtered out the spore suspension with the use of muslin cloth. The inoculum load of  $1.4\times 10^7$  spores/ml was counted using haemocytometer.

**Soil preparation and planting:** Prior to the conduct of the experiment, the screenhouse was adequately sanitized. Agricultural soil of between 0 to 15 cm depth was collected from the experimental site of Botany Department, University of Ibadan, Ibadan. The soil was sterilized using electric soil sterilizer at 120°C for 1 hour, and was filled in polythene bag at 5kg per bag after cooling. Maize seeds were surface sterilized in a beaker containing sodium hypochlorite (3.5% w/v) for 3 minutes after which the seeds were dried in between layers of whatmann filter paper for another 5 minutes. The seeds were planted at three seed per hole across the varieties.

**Experimental Design:** The experiment was laid out in a completely randomized design (CRD) in ten replications. Treatments across the five maize varieties consisted of two inoculation methods; soil and seedling inoculations. These were carried out at varying volume of the standardized ( $1.4\times 10^7$  spores/ml) inoculum at 0 ml, 10 ml and 20 ml of the spore suspensions. The treatment

with soil inoculation was conducted at the period of planting while seedling inoculation was carried out on the respective plots at 2 weeks after planting. The control experiments were treated with respective volume of sterile distilled water. Can include number of replication

**Determination of disease incidence and severity:** The percentage of disease incidence and severity were determined using the formulas:

$$\text{Disease incidence (\%)} = \frac{\text{number of infected maize plants}}{\text{number of maize plants}} \times 100$$

$$\text{Disease severity (\%)} = \frac{\text{area of plant tissue affected}}{\text{total area}} \times 100$$

**Extraction of plant samples:** Two replicates from each treatment across the varieties were carefully uprooted at 4<sup>th</sup>, 5<sup>th</sup> and 6<sup>th</sup> week after planting (WAP). Soil adhering to the root of maize plant was removed by agitation in water, the plants were then air dried at room temperature. Two gramme (2g) of leaf sample per treatments was treated with 10ml of 80% methanol in a tightly clogged flask. The treated leaf was allowed to stand for 3 minutes after which the liquid fraction was filtered using muslin cloth. The step was repeated three times with 80% ethanol and the supernatants were collected into volumetric flasks. Final volume of the extracts was made to 50 ml with 80% ethanol and all the extracts from each sample was combined and centrifuged at 2000 revolutions per minutes for 20 minutes in a centrifuge.

**Determination of phenol content:** This was done using the Folin-Ciocalteu method as described by Singh *et al.* (2012) in which gallic acid served as a standard and total phenol is expressed as mg/g gallic acid equivalents (GAE). The amount of total soluble phenol present in the sample was calculated according to Singh *et al.* (2012).

**Data collection and statistical analysis:** Data were collected on plant height (cm), leaf number (cm), disease incidence (%) and disease severity (%) weekly for 9 weeks. All the data on growth, disease and phenol contents collected were subjected to ANOVA using SAS 9.1 statistical analysis software and Means were separated by Duncan Multiple Range System at 95% confidence interval.

## RESULTS

The fitted model for the interactive effect of maize varieties, inoculation methods, inoculums' quantity and duration of experiment produced a significant ( $p < 0.01$ ) result on the growth and disease parameters measured. All the factors evaluated were significant with plant height, number of leaves and disease severity while only inoculum volume and duration of the experiment (WAP) was significant with disease incidence caused by *F. verticillioides* (Table 1).

Maize variety; BR-9928-DMR-SR followed by ART-98-SW6 recorded the most significant ( $p < 0.05$ ) growth on plant height and number of leaves across the varieties evaluated. This was followed by ART-98- SW1 and BR-9943-DR-SR which also recorded similar level of significance while SWAM 1-SR showed the least growth performances (Table 2).

All the maize varieties recorded more than 70% stalk rot infections with BR-9928-DMR-SR having highest rate of 73.11%. The stalk rot severity ranged from 19.01% in variety ART-98-SW1 to 25.21% in ART-98-SW6 (Figure 1).

Table 3 showed the effect of different inoculation methods and inoculums volume on the growth and disease occurrence in maize plants. While no significant differences was recorded between the inoculation methods with respect to plant height and number of leaves, soil inoculation method produced a significant ( $p < 0.05$ ) disease incidence and severity compare to seedling inoculation which showed no significant difference from the control. Whereas, significant ( $p < 0.05$ ) reduction was recorded in the growth rate with increasing inoculums volume. No significant result was obtained in the disease incidence with respect to inoculums levels while at 20 ml, there was significant increase in stalk rot severity (Table 3).

There was a significant increase in plant height and leaf number with the increasing weeks after planting (Table 4)

Figure 2 showed a linear relationship between the disease incidence caused by *F. verticillioides* in maize and the period of experiment. A consistent increase in the disease progression was observed

140 until the 5<sup>th</sup> week when the infection reached its climax and maintained this position till the 9<sup>th</sup> week  
141 of experiment (Figure 2).

142 The  $r^2$  value of 0.8694 reinstated the reliability of result obtained in the severity of stalk rot with  
143 respect to period of experiment. Having received *F. verticillioides* inoculation in the 2<sup>nd</sup> WAP,  
144 disease severity increased consistently from 3<sup>rd</sup> week (9.95%) to 9<sup>th</sup> week (36.29%) after planting  
145 (Figure 3).

146

UNDER PEER REVIEW

147 **Table 1:** ANOVA table of growth and disease occurrence in maize plants after inoculations with *Fusarium verticillioides*

Sources of variation	df	Plant height (cm)		Leaf number		Disease incidence		Disease severity	
		F value	P value	F value	P value	F value	P value	F value	P value
Model	21	162.46	0.0001**	157.48	0.0001**	210.05	0.0001**	72.74	0.0001**
Maize varieties	4	12.48	0.0001**	14.55	0.0001**	0.30	0.8765	8.82	0.0001**
Inoculums volume	2	43.49	0.0001**	39.33	0.0001**	11.00	0.0011**	34.64	0.0161*
Inoculation method	2	16.16	0.0001**	9.44	0.0001**	0.29	0.5882	5.81	0.0001**
Weeks After Planting	8	405.5	0.0001**	395.34	0.0001**	392.13	0.0001**	172.33	0.0001**
Replicates	9	1.56	0.1693	1.05	0.3864			0.42	0.8372
Error	1327								
Corrected Total	1348								


148

149 \*Significant \*\* Highly Significant.

150

151

152



153

154

155 Figure 1: Disease incidence (DI) and severity (DS) across the maize varieties caused by *F.*  
156 *verticillioides*.

157

158

159

160

161

162

163

164

165

166

167

168

169

170



Table 2: Effect of *F. verticillioides* on plant heights and leaf number across the maize varieties

Varieties	Plant height (cm)	Number of leaves
SWAM 1-SR	43.27 <sup>c</sup>	4.53 <sup>c</sup>
BR-9928-DMR-SR	51.31 <sup>a</sup>	5.47 <sup>a</sup>
ART-98-SW6	54.56 <sup>a</sup>	5.68 <sup>a</sup>
ART-98- SW1	49.52 <sup>b</sup>	5.07 <sup>b</sup>
BR-9943-DR-SR	48.94 <sup>b</sup>	4.97 <sup>b</sup>
R <sup>2</sup>	0.72	0.71

Mean with different letters are significantly different ( $p \leq 0.05$ )

187 Table 3: Effect of inoculation method and inoculum volume on growth and disease occurrence in  
188 maize plants

Parameters	Variables	Plant height (cm)	Number of leaves	Disease incidence (%)	Disease severity (%)
<b>Inoculums' volume (ml)</b>	Control	54.11a	5.63a	70.18a	21.51b
	10	48.88b	4.90b	71.78a	22.18ab
	20	41.58c	4.63c	72.65a	23.99a
<b>Inoculation method</b>	Control	52.99a	5.23a	69.56b	20.69b
	Seedling inoculation	46.72b	5.06a	70.18b	21.51b
	Soil inoculation	50.58b	5.17a	74.87a	25.59a
	R <sup>2</sup>	0.72	0.71	0.94	0.53

189

190 Means with different letters are significantly different ( $p \leq 0.05$ )

191


192 Table 4: Effect of time (WAP explain) on the growth of treated maize plants

	Weeks After Planting	Plant height (cm)	Leaf number
	1	2.08i	0.01i
	2	11.78h	1.59h
	3	24.61g	3.07g
	4	39.63f	4.35f
	5	54.69e	5.61e
	6	66.92d	6.50d
	7	75.94c	7.43c
	8	82.57b	8.38b
	9	87.65a	9.36a
	R <sup>2</sup>	0.72	0.71

193

194 Means with different letters in a column are significantly different

195



197


198 Figure 2: Effect of time (WAP) on disease incidence caused by *F. verticillioides* in maize plants.

199

200

201

202



203

204 **Figure 3:** Effect of time (WAP) on disease severity of *F. verticillioides* in maize plants

The model for the reaction of quantity of phenol produced was significant ( $p<0.01$ ) through the period of experiment. The effect of maize varieties and inoculums volume also recorded significant result at the 4<sup>th</sup>, 5<sup>th</sup> and 6<sup>th</sup> (WAP). The inoculums' method used showed significance ( $p<0.01$ ) at 4<sup>th</sup> and 5<sup>th</sup> WAP while it was significant at  $p<0.05$  at 6<sup>th</sup> WAP (Table 5).

Maize variety ART-98-SW6 showed the most significant ( $p<0.05$ ) phenol content across the period of this study. This result was followed by BR-9928-DMR-SR while ART-98-SW1 and BR-9943-DMR-SR showed no significant difference (Table 6).

The seedling inoculation method showed significantly higher phenol contents than soil inoculation, although the two methods produced results which were significantly higher than the control. Moreover, inoculums volume of 10ml recorded a significantly higher phenol contents than at 20ml (Table 7).

The inoculation method were positively and significantly ( $p<0.01$ ) correlated with inoculums volume ( $r=0.64$ ). Also, inoculation method was significantly associated with phenol production at week 4 ( $r=0.60$ ), week 5 ( $r=0.68$ ) and week 6 ( $r=0.71$ ) after planting. Similarly, inoculums' volume was significantly correlated with phenol content obtained at week 4 ( $r=0.81$ ), week 5 ( $r=0.90$ ) and week 6 ( $r=0.89$ ) of the experiment (Table 8).

**Table 5:** ANOVA table for Phenol (mg/g) produced by the maize plants after days of infection

Source	df	4th WAP		5th WAP		6th WAP	
		F value	P>F	F value	P>F	F value	P>F
Model	7	90.24	0.0001**	227.49	0.0001**	310.19	0.0001**
Maize varieties	4	49.92	0.0001**	67.34	0.0001**	74.98	0.0001**
Inoculation methods	1	9.59	0.0065**	9.79	0.0061**	4.21	0.0560*
Inoculums volume	1	139.13	0.0001**	306.88	0.0001**	267.68	0.0001**
Error	17						
Corrected Total	24						

\*\*=Highly significant \*= Significant

**Table 6:** Phenol (mg/g) content of the maize plants after days of infection

Varieties	4th WAP	5th WAP	6th WAP
SWAM1-BR-SR	7.40 <sup>d</sup>	11.12 <sup>d</sup>	14.04 <sup>d</sup>
BR-9928-DMR-SR	13.80 <sup>b</sup>	16.92 <sup>b</sup>	20.96 <sup>b</sup>
ART-98-SW6	16.16 <sup>a</sup>	19.04 <sup>a</sup>	21.28 <sup>a</sup>
ART-98-SW1	11.20 <sup>c</sup>	15.04 <sup>c</sup>	19.22 <sup>c</sup>
BR-9943-DMR-SR	10.52 <sup>c</sup>	14.54 <sup>c</sup>	18.74 <sup>c</sup>
R <sup>2</sup>	0.97	0.99	0.99

Means with different letters are significantly different ( $p \leq 0.05$ )

**Table 7:** Phenol (mg/g) content of the maize plants days after inoculation with different volumes of *Fusarium verticillioides*

Parameter	Variable	4th WAP	5th WAP	6th WAP
Inoculation method	Control	4.72 <sup>c</sup>	5.94 <sup>c</sup>	7.88 <sup>c</sup>
	Seedling	14.32 <sup>a</sup>	16.33 <sup>a</sup>	17.58 <sup>a</sup>
	Soil	12.86 <sup>b</sup>	14.03 <sup>b</sup>	15.48 <sup>b</sup>
	R <sup>2</sup>	0.97	0.99	0.99
Inoculums volume	Control	4.97 <sup>c</sup>	5.94 <sup>c</sup>	7.88 <sup>c</sup>
	10ml	13.37 <sup>a</sup>	15.32 <sup>a</sup>	18.61 <sup>a</sup>
	20ml	10.81 <sup>b</sup>	13.04 <sup>b</sup>	15.63 <sup>b</sup>
	R <sup>2</sup>	0.97	0.99	0.99

Means with different letters are significantly different ( $p \leq 0.05$ ).

233 **Table 8:** Extent of association between the maize varieties, pathogen inoculation, and time with  
 234 phenol content

Correlation	Maize varieties	Inoculation method	Inoculums' volume	Phenol contents		
				Week 4	Week 5	Week 6
Maize varieties						
Inoculation method	0.00					
Inoculums' volume	0.00	0.64**				
Week 4	-0.13	0.60**	0.81**			
Week 5	-0.11	0.68**	0.90**	0.98**		
Week 6	-0.72	0.71**	0.89**	0.94**	0.99**	

236 \* Significant, \*\*Highly Significant

## 238 DISCUSSION

239 High incidence of stalk rot infection recorded across the *Fusarium verticillioides* inoculated maize  
 240 varieties supported the claim that *F. verticillioides* is the major causal agent of highly variable  
 241 disease symptoms ranging from asymptomatic plants to severe rotting and wilting in maize (Oren et  
 242 al., 2003; Sobowale et al., 2005; Aliakbari, 2007). Virulence of *F. verticillioides* that ranged from  
 243 25.21% in ART-98-SW6 to 19.01% in ART-98-SW1 demonstrated the variations in the severity of  
 244 systemic infection on the host, while environmental factors also play a major in the activities of this  
 245 cosmopolitan pathogen (Murillo-Williams and Munkvold, 2008; Thompson and Raizada, 2018).  
 246 More so, (delete) variation recorded in the growth rate and resistance of maize varieties to *F.*  
 247 *verticillioides* infections is in consonance some earlier findings in which host resistance was  
 248 reported as a major determinant of variations in the activities of the pathogen (Sharma, 2003; Olowe  
 249 et al. 2017).

250 Similar rate of maize growths observed in both seedling and soil inoculation methods is contrary to  
 251 the results obtained by Khan et al. (2012) in which a significant increase in the growth parameters of  
 252 plants that received *Alternaria inocula* through different methods was reported. Also, higher

incidence and severity of stalk rot recorded in the soil inoculation method conforms to the earlier claims that attributed varying severities caused by artificial inoculation of *F. verticillioides* with different inoculation methods (Drepper and Renfro, 1990; Sobowale, 2011). Furthermore, the increase in inoculums' quantity which resulted in corresponding decrease in growth rate of maize plants revealed varied efficiency as regards inoculum delivery with different inoculums' level (Sobowale et al., 2007).

The prevalence of stalk rot incidence observed to reach its peak at 5<sup>th</sup> week after planting affirmed that fungus colonizes maize stalks systemically without necessarily causing visible disease symptoms and that *F. verticillioides* strains can be vertically transmitted through seed-to-plant transmission and systemic stalk infection (Munkvold, 1997; Murillo-Williams and Munkvold, 2008). Meanwhile, the severity rate of 36.29% at 9<sup>th</sup> week after inoculation justifies the epidemiological claim that *Fusarium* stalk rot reduces output in maize by 10% typically and by 30–50% in severely affected areas (Li et al. 2010).

The result of maize variety ART-98-SW6, followed by BR-9928-DMR-SR, ART-98-SW1, BR-9943-DMR-SR and SWAM1-BR-SR as the decreasing order of phenol production after inoculated with *F. verticillioides* was in agreement with some earlier reports that both constitutive or induced synthesis contribute to abundance and composition of phenolic compounds in cereal grains and this is highly variable depending on the species, variety and environmental conditions (Adom and Liu 2002; Lattanzio et al. 2006). The observation of Reddy and Sireesha (2013) that nutritional status and concentration of biochemical constituents in plants prior to infection determines the severity of disease possibly explains the reasons maize varieties ART-98-SW6 and BR-9928-DMR-SR with higher stalk rot severities also produced higher phenol contents than other varieties tested. This further corroborated the reports that plants respond to pathogen invasion through the activation of complex defense strategies such as the accumulation of flavonoids, phytoalexins and phenolic compounds (Delledonne et al., 2001; Hefny et al., 2012). In this work, total phenol produced by the maize varieties acts as biochemical markers to analyse disease incidence and severity which

conforms to the work done by Singh et al. (2012) on biochemical response and host-pathogen relationship of stalk rot fungi in early stages of maize (*Zea mays*).

Higher phenol contents produced in seedling inoculation method compare to soil inoculation could be associated with plants' reaction to direct inoculation which possibly resulted into a more effective delivery of *F. verticillioides* inoculums. Thus, the possibility of the pathogen inducing phenol production could be substantiated by the report that biochemical resistance, tolerance or susceptibility in plants against any disease depends mainly on preexisting, preformed or induced substances by the pathogen in the host (Reddy and Sireesha, 2013). In this study, the higher phenol production that occurred at lower inoculum volume was consistent with the findings of Perveen et al. (2010) who reported a decrease in total phenol of the leaves of *M. arvensis* with increase in initial inoculum of *S. sclerotiorum*, a situation attributed to the altered rates of synthetic activity because of infection by pathogen (Howlett, 2006).

The inoculation methods and inoculums volume used in this study effectively delivered infective stalk rot inoculum dosage, the rate of which were found strongly correlated with phenol production in maize plants at 4<sup>th</sup>, 5<sup>th</sup> and 6<sup>th</sup> week after inoculation. Thus, agreed with the claim that majority of phenolic compounds are a part of the preformed general defense system against potential pathogens (Stuper-Szablewska et al. 2017).

In general, phenol production in the maize varieties corresponds with the rate of severity of *F. verticillioides* infections. Therefore, close association exists between stalk rot disease and maize plants' resistance to infection through phenol production. Hence, researches that will increase phenol production in crops especially maize, may thus be suggested.

300

#### 301 **REFERENCES-Please check with the journal format, Journal representation uniform**

302 Adom KK, Liu RH. (2002). Antioxidant activity of grains. *J Agric Food Chem*. 50:6182–6187

303

304 Akande S.R. and Lamidi G.O. (2006). Performance of quality protein maize varieties and disease  
305 reaction in the derived-savanna agro-ecology of South-West Nigeria. *African Journal of*  
306 *Biotechnology*. 5(19), 1744-1748.



307 Akanmu A. O., Abiala M. A., Akanmu A.M., Adedeji A. D, Mudiaga P. M and Odebode A. C.  
 308 (2013) Plant Extracts Abated Pathogenic *Fusarium* Species of Millet Seedlings. *Archives of*  
 309 *Phytopathology And Plant Protection*, 46 (10) 1189-1205,  
 310  
 311 Alankoya A.E. Monda E.O. and Ajanga S. (2008). Variation in *in vitro* fumonisin B1 production by  
 312 different *Fusarium verticillioides* isolates in Kenya. *American-Eurasian Journal for Agriculture and*  
 313 *Environmental Science*. 4: 368-371.

314 Aliakbari F., Mirabolfathy M., Emami M., Mazhar S.F. and Karami-Osboo R., (2007). Natural  
 315 Occurrence of *Fusarium* species in Maize Kernels at Gholestan Province in Northern Iran. *Asian*  
 316 *Journal of Plant Sciences*, 6: 1276-1281.

317 Boudet A (2006) Evolution and current status of research in phenolic compounds. *Phytochemistry*  
 318 68:2722–2735. (fullstop)  
 319

320 Cowan M (1999) Plant products as antimicrobial agents. *Clin Microbiol Rev* 12:564–582. (fullstop)  
 321

322 Delledonne, M., J. Zeier, A. Marocco and C. Lamb (2001). Signal interactions between nitric oxide  
 323 and reactive oxygen intermediates in the plant hypersensitive disease-resistance response. *Proc.*  
 324 *Natl. Acad. Sci. USA.*, 98: 13454-13459.  
 325

326 Drepper W.J. and Renfro B.L. (1990). Comparison of Methods for Inoculation of Ears and Stalks of  
 327 Maize with *Fusarium moniliforme*. *Plant Dis.* 74:952-956.  
 328

329 Gwinner J., Harnisch R., Much O. (1996). Manuel sur la manutention et la conservation des grains  
 330 apres-recolte. GTZ, Eschborn, Germany. P 38.

331 Hefny M., Attaa S., Bayoumi T., Ammar S.H. and El- Bramawy M. (2012). Breeding Maize for  
 332 Resistance to Ear Rot Caused by *Fusarium moniliforme*. *Pakistan Journal of Biological Sciences*,  
 333 15: 78-84.

334 Howlett B.J. (2006) Secondary metabolite toxins and nutrition of plant pathogenic fungi *Cur.*  
 335 *Opinion Plant Biol.*, 9, pp. 371-375. (fullstop)  
 336

337 Khan M.M., Khan M.R. and Mohiddin F.A. (2012). The Relative Performance of Different  
 338 Inoculation Methods with *Alternaria brassicae* and *A. brassicicola* on Indian Mustard. *Plant*  
 339 *Pathology Journal*, 11: 93-98.

340 Lattanzio V, Lattanzio V.M., Cardinali A. (2006). Role of phenolics in the resistance mechanisms of  
 341 plants against fungal pathogens and insects. *Phytochem: Adv Res.* 661:23–67.

342 Li W.J., He P and Jin J.Y. (2010) Effect of potassium on ultrastructure of maize stalk pith and young  
 343 root and their relation to stalk rot resistance. *Agricultural Sciences in China*, 9:1467–1474.

344 Maddox C.E., Laur L.M. and Tian L. (2010) Antibacterial activity of phenolic compounds against  
 345 the phytopathogen *Xylella fastidiosa*. *Curr Microbiol.* 60(1):53-58.  
 346

347 Masuka A.J., Cole D.L., and Mguni C. (2003). List of plant diseases in Zimbabwe. Plant Protection  
 348 Research Institute.

349 Munkvold, G. P., McGee, D. C., and Carlton, W. M. (1997). Importance of different pathways for  
 350 maize kernel infection by *Fusarium moniliforme*. *Phytopathology* 87:209-217. (fullstop)

351 Murillo-Williams A. and Munkvoid G.P. (2008) Systemic Infection by *Fusarium verticillioides* in  
 352 Maize Plants Grown Under Three Temperature Regimes. *Plant Disease* 92 No. 12, pages 1695 -  
 353 1700 . (fullstop)

354 Okoro-Robinson M.O., Olawuyi O.J., Bello W.O. and Babalola B.J. (2014). Comparative evolution  
355 of organic manure on growth and yield of maize. *Agricultural and Biological Research*. 30(1): 60-  
356 73.

357 Olakojo S.A. and Iken J.E. (2001). Yield performance and stability of some improved maize (*Zea*  
358 *mays* L.) varieties. *Moor Journal of Agricultural Research*. 2: 21-24.

359 Olawuyi O.J., Bello O.B., Ntuba C.V. and Akanmu A.O. (2015) Progress from selection of some  
360 maize cultivars' response to drought in the derived savanna of Nigeria. *Agrivita*, 37 (1) 8-17.  
361 (fullstop)

362 Olawuyi O.J., Odebode A.C., Alfar A., Olakojo S.A. and Adesoye A.I. (2010). Performance of  
363 maize genotypes and arbuscularmycorrhizal fungi in Samara District of Southwest Region of Doha-  
364 Qatar. *Nigerian Journal of Mycology*, 3(1): 86-100.

365 Olowe, O.M., Odebode, A.C., Olawuyi, O.J. and Sobowale, A.A. (2017) Molecular Variability of  
366 *Fusarium verticillioides*(Sacc.) in Maize from Three Agro-Ecological Zones of Southwest Nigeria.  
367 *American Journal of Molecular Biology*, 7, 30-40. (fullstop)

368 OnoE.Y.S., Fungaro M.H.P., Sofia S.H., Miguel T.D., Sugiura Y., Hirooka E.Y. (2010). *Fusarium*  
369 *verticillioides* strains isolated from corn feed: characterization by fumonisin production and RAPD  
370 fingerprinting. *Braz. arch. biol. technol.* vol.53 no.4 Curitiba

371 Oren L., Ezrati S., Cohen D., and Sharon A. (2003) Early Events in the *Fusarium verticillioides*-  
372 Maize Interaction Characterized by Using a Green Fluorescent Protein-Expressing Transgenic  
373 Isolate. *Appl Environ Microbiol*. 2003 Mar; 69(3): 1695–1701.

374 Orsi R.B., Correa B., Possi C.R., Schammas E.A., Nogueira J.R., Dias S., Malozzi M.A.B. (2000).  
375 Mycoflora and occurrence of fumonisin of freshly harvested and stored hybrid maize. *Journal of*  
376 *storage and Product Research*. 36-87.

377 Perveen K., Haseeb A. and Shukla P.K. (2010) Effect of *Sclerotinia sclerotiorum* on the disease  
378 development, growth, oil yield and biochemical changes in plants of *Mentha arvensis*. *Saudi Journal*  
379 *of Biological Sciences*, 17, Issue 4, Pages 291-294

380 Reddy M.N. and Sireesha C.H. (2013). Role of oxidative enzymes and biochemical constituents in  
381 imparting resistance to groundnut (*Arachis hypogea* L.) against stem rot of diseases caused by  
382 *Sclerotium rolfsii*. *Bioresearch Bulletin*. 36-41pp.

383 Samapundo S., De Meulenaer B., Osei-Nimoh D., Lamboni Y., Debevere J., Devlieghere F. (2007).  
384 Can phenolic compounds be used for the protection of corn from fungal invasion and mycotoxin  
385 contamination during storage? *Food Microbiology* 24 (2007) 465–473.

386 Sharma, T.R. (2003) Molecular Diagnosis and Application of DNA Markers in the Management of  
387 Fungal and Bacterial Plant Diseases. *Indian Journal of Biotechnology*, 2, 99-109. (fullstop)

388 Singh N., Ambika R., Meena S. and Girish M. 2012. Biochemical response and host-pathogen  
389 relation of stalk rot fungi in early stages of maize (*Zea mays* L.). *African Journal of Biotechnology*.  
390 11 (82): 14837-14843.

391 Singh P., Shukla R., Prakash B., Kumar A., Singh S. and Mishra P.K.( 2010). Chemical profile,  
392 antifungal, antiaflatoxic and antioxidant activity of *Citrus maxima* Burm. and *Citrus sinensis* (L)  
393 Osbeck essential oils and their cyclic monoterpene, DL-limonene. *Food and chemical toxicology*. 48  
394 (6), 1734-1740.

401 Sobowale A.A. (2011). Determination of infective, non-lethal dosage of *Fusarium verticillioides* in  
 402 maize (*Zea mays*) stem and effective inoculation method in the screenhouse. Journal of Agriculture  
 403 and Biological Sciences Vol. 2(5) pp.118-122. (fullstop)  
 404

405 Sobowale A.A., Cardwell K.F., Odebode A.C., Bandyopadhyay R., Jonathan S.G. (2007).  
 406 Persistence of *Trichoderma* species within maize stem against *Fusarium verticillioides*. Arch.  
 407 *Phytopathol. Plant Prot.* 40, 3: 215-231.  
 408

409 Sobowale A.A., Cardwell K.F., Odebode A.C., Bandyopadhyay R., Jonathan S.G. (2005). Growth  
 410 inhibition of *Fusarium verticillioides* (Sacc.) Nirenberg by isolates of *Trichoderma Pseudokoningii*  
 411 strains from maize plant parts and its rhizosphere. Journal of Plant Protection Research, 45 (4) Pages  
 412 249-265.

413 Stuper-Szablewska K, Kurasiak-Popowska D, Nawracała J, Perkowski J. (2017). Response of non-  
 414 enzymatic antioxidative mechanisms to stress caused by infection with *Fusarium* fungi and chemical  
 415 protection in different wheat genotypes. Chem Ecol. 33:949–962.

416 Thompson M. and Raizada M. (2018). Fungal Pathogens of Maize Gaining Free Passage Along the  
 417 Silk Road. Pathogens, 7, Number 4, 81. (fullstop)  
 418