Original Research Article

Mechanical Damages and Storage Conditions Affect the Quality of Markies cv. Potatoes for Processing

5 6

1 2

3

4

7

8 ABSTRACT

Aims: Evaluate whether mechanical damage and storage conditions affect the quality of the 'Markies' potato for processing.

Study design: The experimental design was the completely randomized, in the scheme of split-plots. The plots were composed of treatments with and without mechanical damage and the subplots were made up by the evaluation times, with 5 replicates, where the experimental unit was composed of 2 tubers.

Place and Duration of Study: Tubers of the 'Markies' cultivar from the producing region of Perdizes, state of Minas Gerais, were planted in June and harvested in October 2016.

Methodology: The tubers were stored for 2 months at 8 °C, and further divided into treatment with damage, in which the tubers were subjected to impact and abrasion; and control treatment, in which the tubers were not damaged. After that, they were stored at 28 °C and evaluated for their loss of accumulated fresh mass (FML), total soluble sugar (TSS), reducing sugar (RS), non-reducing sugar (NRS) and color after frying in the periods of 0, 12, 24, 36 and 48 h after being placed at room temperature.

Results: The mechanical damage in the tubers increased FML, however, it did not affect the content of TSS, NRS, RS and color after frying. Increases were observed in the content of TSS and NRS after 12 h of evaluation. Grade 2 was assigned to the coloring scale after frying for the tubers regardless of treatment or evaluation period.

Conclusion: Mechanical damage increases the FML and the 'Markies' cultivar is suitable for

the industry of pre-fried potato processing even under the occurrence of mechanical damage and exposure to high temperatures.

9 Key-words: color; FML; POD; PPO; sugars.

10 **1. INTRODUCTION**

Potato (*Solanum tuberosum* L.) is the third most important and consumed crop in the world [451]. The consumption of processed potatoes has been increasing as a result in the changes in the diets, the increase in fast food chains and the need for semi-finished foods. To meet the market for processed potatoes, it is necessary cultivars suitable for the industry, and the execution of refrigerated storage for quality maintenance and constant supply.

16 Cultivar 'Markies' is destined to the processing industry and with qualities for cooking 17 and frying. It has high quality for the pre-fried potato market because it presents high content of 18 dry matter, low sugar content and high levels of starch, obtaining a product with desirable flavor 19 and coloration after frying [422].

20 The refrigerated storage allows the maintenance of the guality by the reduction of the 21 sprouting, respiratory activity and incidence of diseases. However, when stored at low 22 temperatures, the tubers can accumulate reducing sugars, which, when reacting with amino 23 acids, forms melanoidine pigments by non-enzymatic reaction called Maillard reaction during 24 frying [443]. This reaction promotes darkening of sticks and make them unsuitable for 25 consumption. In addition, in a secondary route, there is the formation of acrylamide, a 26 substance considered neurotoxic and carcinogenic with several studies proving a positive 27 association with human cancer [143].

The process of unloading in the cold chambers and postharvest handling of the tubers results in the occurrence of mechanical damages. The wounds develop a depreciation of the product by physiological changes, entry of pathogenic microorganisms and increase in the water loss [84]. In addition, between the storage and processing stages, the tubers stay for about 48 hours at room temperature, which reduces even more the quality of the products. Therefore, the objective of this study was to evaluate whether mechanical damage and
 storage conditions affect the quality of the 'Markies' potato for processing.

35 2. MATERIAL AND METHODS

Tubers of the 'Markies' cultivar from the producing region of Perdizes, state of Minas Gerais, were planted in June and harvested in October 2016. They were selected according to their size and lack of damage. Curing of the tubers was carried out at 14 °C (\pm 95-% <u>Relative</u> <u>Humidity (RH)</u> for 10 days. After this period, the temperature was reduced by 1 °C per day until reaching the storage temperature of 8 °C (\pm 90-% RH).

The tubers were stored for 2 months and later divided into control treatment, where the tubers were not harmed; and treatment with damage caused by impact and abrasion caused by the fall of the tubers from 1 meter of height on rough surface. This process was repeated 10 times for each tuber. After application of the treatments, the tubers were kept on benches at room conditions (28 °C and \pm 54-% RH)

The analysis of loss of accumulated fresh mass (FML), total soluble sugars (TSS), reducing sugars (RS), non-reducing sugars (NRS), color of the sticks after frying, polyPoly phenolPhenol_exidase_Oxidase (PPO) and peroxidase activity (POD) were carried out. The analyses started two hours after the treatments, with a 12-h interval between the evaluations, following the times 0, 12, 24, 36 and 48 h.

51 The FML was calculated in relation to the initial mass of the tubers and expressed in 52 percent%. The quantification of TSS and RS followed the Phenol-sulfuric method [35] and the 53 dinitrosalicylic acid (DNS) method [46], respectively. The NRSs were calculated by difference 54 between TSS and RS concentrations and the results expressed in %-percent. The color of the 55 post-fry potatoes was visually determined based on the grading scale recommended by the 'United States Standards for Grades of Frozen French Fried Potatoes' [437] and the fast food 56 57 industry color grading chart from 1 to 5. The POD and PPO were determined according to the methodology proposed by [6-8] and [59], respectively, and the results were expressed based on 58

Comment [D1]: What is this male abbreviation

59 total protein. The protein was evaluated using bevine Bovine serum Serum albumin Albumin

60 (BSA) as standard [210].

The experimental design was the completely randomized, in the scheme of split-plots. The plots were composed of treatments with and without mechanical damage and the subplots were made up by the evaluation times, with 5 <u>replicatesreplications</u>, where the experimental unit was composed of 2 tubers. Data were analyzed through analysis of variance and regression, using the [711]. The regression model was chosen based on the regression coefficients by using the t-test at the P = 0,05 probability level, at the coefficient of determination ($R^2 = SQReg / SQtrat$) and the biological behavior under study.

68 3. RESULTS AND DISCUSSION

The loss of accumulated fresh mass (FML) in ev. Markies cultivat of potato tubers with 69 70 and without damage was increasing in relation to the evaluation time (Fig. 1). The highest 71 increase in the FML in tubers occurred 48 h after the application of the damage, with values of 72 3.7-% in relation to tubers without damage (0.55-% FML), representing an increase of 6.5 times. 73 Mechanical damage increases respiratory activity and loss of water because of the exposition of 74 the tissues to the environment, facilitating gas exchange, contributing to the reduction of mass 75 loss. Observed a 1.3-fold increase in the respiratory rate of tubers of damaged potatoes in 76 comparison to the undamaged potatoes when kept at 33 °C [1112].



77

78 Figure 1. Loss of accumulated fresh mass (FML) in potato tubers cv. Markies without damage

79 $\,$ (SD) and mechanical damage (CD) after 2 months of storage.

80 The occurrence of tuber damage did not interfere with the contents of TSS and NRS. 81 However, an influence was observed in the evaluation time (Fig. 2). Transfer of tubers from 82 refrigerated storage at 8 °C to room temperature of 28 °C may have resulted in the increment in 83 respiratory activity, leading to the conversion of starch into sucrose, which caused the increase 84 in NRS and consequently of TSS. Storage of potatoes at temperatures higher than 15 °C culminating in the increment in the respiration [913]. The TSS and NRS showed a minimum 85 86 point at 12 h of evaluation (Fig. 2), indicating that it took 12 h at room temperature to begin the 87 conversion of starch to sucrose. The increase in TSS and NRS contents from the 12 h of 88 evaluation did not affect the quality of the potato for processing; the highest levels of NRS were 89 48 h with 0.14 %. For the processing industry, the NRSs should remain below 0.33 % of the 90 fresh mass [210].





93 Markies with and without mechanical damage after 2 months of storage.

	Same Davage		Cum Dana	
	Ø kenne	48 incas	() incas	48 horas
$\mathcal{O}_{\mathcal{F}}$		Man	MAM	
			Milling	MAIN
	UIII	MAR	NIMIN	Man

Formatted: Font: (Default) Arial, 11 pt, Font color: Text 1

94

91

95 Figure 3. Post-frying coloration of cv. Markies with and without mechanical damage after 2

96 months of storage and 48 hours after the mechanical damage was determined based on the

97 grading scale recommended by the 'United States Standards for Grades of Frozen French Fried

98 Petatees' (USDA, 1967) and the fact feed inductry celor grading chart from 1 to 5.

99 No effect of the mechanical damage and the evaluation period were observed on the

100 contents of RS and the color of the sticks after frying, and all the fried sticks were given score 2

101 according to the standards recommended by the USDA (fig. 3).

102



Formatted: Font: (Default) Arial, 11 pt, Font color: Text 1

Figure 3. Post-frying coloration of cv. Markies with and without mechanical damage after 2
 months of storage and 48 hours after the mechanical damage was determined based on the
 grading scale recommended by the 'United States Standards for Grades of Frozen French Fried
 Potatoes' (USDA, 1967) and the fast food industry color grading chart from 1 to 5.

Darkening is related to the increase in the contents of RS induced by refrigerated storage, in which the carbonyl or ketone group of the RS molecule react with the amino group of the amino acids, like asparagine, resulting in the formation of the melanoidine pigments by nonenzymatic reaction called Maillard reaction at the time of frying. Cultivar Markies presents low sugar content [422], which justifies the non-darkening observed in the experiment.

The acceptable content of RS for the pre-fried potato processing industry should be below 0.12 % of fresh mass [4014] to obtain a good quality product with light yellow coloration. The average RS content was 0.016 %, indicating that Markies cultivar is suitable for industrial processing even after mechanical injury and exposure to high temperature.

116 **4. CONCLUSION**

117 Mechanical damage increases the FML and the 'Markies' cultivar is suitable for the industry of 118 pre-fried potato processing even under the occurrence of mechanical damage and exposure to 119 high temperatures.

120 REFERENCES

- 121 1. Bradford MM. A rapid and sensitive method for the quantification of microgram quantities of
- 122 protein utilizing the principle of protein-dye biding. Analytical Biochemistry. 1976; 72:248-254.
- 123 2<u>10</u>. Chapper M, Bacarin MA, Pereira AS, Terrible LC. Carboidratos não estruturais em
 124 tubérculos de dois genótipos de batata armazenados em duas temperaturas. Hortic. Bras.
- 125 2002; 20 (4) : 583-588. Available: http://dx.doi.org/10.1590/S0102-05362002000400014.
- 126 3<u>5</u>. Dubois M, Gilles KA, Hamilton JK, Rebers PT, Smith F. Colorimetric method for
 127 determination of sugars and related substances. Anal. Chem. 1956; 28: 350-356. Available:
 128 http://dx.doi.org/10.1021/ac60111a017.
- 129 4<u>6</u>. Gonçalves C, Rodriguez-Jasso RM, Gomes N, Teixeira JÁ, Bello I. Adaptation of
 130 dinitrosalicylic acid method to microtiter plates. Anal. Methods. 2010; 2: 2046-2048. Available:
 131 http://dx.doi.org/10.1039/C0AY00525H.
- 132 **5.9** kavrayan D, aydemir T. Partial purification and characterization of polyphenoloxidase from
- 133 peppermint (Mentha piperita). Food Chemistry. 2001; 74: 146-154.
- 134 6.8 Marques AE, Silva F, Barbosa JB, Finger FL. Ação de inibidores de enzimas oxidativas e
- 135 crescimento bacteriano sobre a longevidade das flores de Ave-do-Paraíso (Strelitzia reginae
- 136 Aiton). Revista Brasileira de Horticultura Ornamental, 2011; 17(1):75-86.
- 137 Z<u>11</u>. SAEG. Sistema para Análises Estatísticas, Versão 9.1: Fundação Arthur Bernardes: UFV
 138 Viçosa; 2007.
- 139 84. Silva GO da, Pilon L. Colheita e pós-colheita: Sistema de Produção da Batata. Embrapa
- 140 Hortaliças: Sistema de Produção; 2016. 252.

Comment [D2]: No reference found in the text so, it must be deleted or refer at appropriate place

- 141 913. Singh J, Kaur L. Advances in potato chemistry and technology. 2nd ed. Academic press;
 142 2016.
- 143 4014. Stark JC, Olsen N, Kleink OP, Ge F, Love SL. Tuber quality. In: Stark JC, Love SL.
 144 Potato production. Aberdeen: University of Idaho; 2003. 329-343.
- 14514412. Strehmel N, Praeger U, Konig c, Fehrle I, Erban A, Geyer M, Kopka J, Dorgen JTV. Time146course effects on primary metabolism of potato (Solanum tuberosum) tuber tissue after147mechanical impact. Postharvest Biol. Technol. 2010; 56(2):109-116. Available:
- 148 <u>https://doi.org/10.1016/j.postharvbio.2009.12.008</u>

149 422. Morgan T. Potato 'Markies' (Early Maincrop) Solanum tuberosum. Accessed 30 set. 2018.

- Available from: http://www.thompson-morgan.com/vegetables/potatoes/maincrop/potatomarkies/t17927TM#additional-links.
- 152 <u>137</u>. USDA. United States Standards for Grades of Frozen French Fried Potatoes. USDA,
 153 Washington, 1967; 16.
- 154 443. Vinci RM, Mestdagh F, Meulenaer B. Acrylamide formation in fried potato products-
- 155 Present and future, a critical review on mitigation strategies. Food Chem. 2012; 133 (4): 1138-
- 156 1154. Available from: <u>https://doi.org/10.1016/j.foodchem.2011.08.001.</u>
- 157 451. Wang Q, Cao Y, Jiang CZ, Feng Y, Wei S. Effects of postharvest curing treatment on flesh
- 158 colour and phenolic metabolism in fresh-cut potato products. Food Chem. 2015; 169:246-254.
- 159 Available from: https://doi.org/10.1016/j.foodchem.2014.08.011.