

# **Original Research Article**

## **Biodiversity in Forest Fragments Under Different Forms of Environmental Conservation**

### **ABSTRACT**

The objective of this work was to evaluate the biodiversity of forest fragments at different levels of anthropization. The related research was carried out in three forest fragments, an area under reforested conditions, a natural remnant area and an area under natural regeneration. Each area was divided into three sampling units of 240 m<sup>2</sup>, constituting the replicates. The total frequency of insects, amphibians, birds, mammals, reptiles and trees was evaluated. By the cluster analysis, there was dissimilarity between the natural regeneration area and the reforestation and natural remnant areas. The analyzed variables were summarized in two main components, which explain 87.1% of the accumulated total variance. It was concluded that there are differences regarding biodiversity in the forest fragments, highlighting the areas of natural remnants and reforestation, which presented high frequency of the raised species and, consequently, greater biological diversity detriment to the natural regeneration area.

*Keywords: Environment; ecology; fauna; flora; frequency of species.*

### **1. INTRODUCTION**

In the current situation the environment makes necessary the sum of efforts to better understand the dynamic interplay between man and environment. The growing environmental awareness and the expansion of knowledge between the different knowledge areas have mobilized the scientific community and the population in favor of getting to know these interaction man/environment [1] and design strategies for the sustainable exploitation of the environment by man.

Among other things, the human exploratory actions have promoted significant changes in the dynamic equilibrium of ecosystems. In fact, the interference of man can cause disorders to environmental factors [2], notably because of the disturbances in the natural habitat of several species, such as insect eating birds [3, 4, 5], insect bioindicators[6], in addition to plant species [7, 8, 9].

To quantify the quality of habitat for wildlife is a task that is extremely challenging, this being essential to the development of quantitative techniques with robustness sufficient to express the real ability of the natural shelters [10].

On the basis of the above, it is emphasized that the stratification of areas to study their quality, is a preponderant step to understand the peculiarities of each environment [11]. Thus, [12] point out that areas in reforestation can be divided into three main categories: assisted natural regeneration, direct sowing, and planting of seedlings; these areas being, according to [13], essential for the maintenance of biodiversity. Already [14], dealing with areas of natural remnants, reports the contribution of these environments to the richness of

Comment [D1]: delete

Comment [D2]: the

Comment [D3]: change to 'use'

fauna and flora. [9] add that areas in natural regeneration derive from the interaction of processes, converging in an extremely important phenomenon, highly complex and dynamic.

Owing to the importance of the ecosystems mentioned above, [10] point out that, their quantitative analysis consists of an important strategy for the generation of local environmental quality indicators in order to subsidize the decision making on areas to be destined for legal reserve. However, it is known that the evaluation of forest fragments is usually based on a single taxonomic group, evidencing the need for rapid assessments based on multitax on indicators [15] that can be excellent tools to help conservationists and managers in the definition of environmental conservation strategies [14]. The objective of this work was to analyze the biodiversity of three forest fragments under different forms of environmental conservation.


## 2. MATERIAL AND METHODS

### 2.1 Research Coverage Area

The research was carried out in April 2015 in three forest areas in the municipality of Jaboticabal, State of São Paulo, Brazil. The areas covered by the study, consisting of 720 m<sup>2</sup> each, were characterized according to the level of anthropization, namely: Area under reforestation conditions, denominated fragment 1 (FRA-1), located at 21°14'54.7"S and 48°17'48.5"W; area of natural remnant, fragment 2 (FRA-2), located at 21°14'47.1"S and 48°17'29.4"W; and area under natural regeneration, fragment 3 (FRA-3), located at 21°15'02.5"S and 48°17'42.3"W. During the month of conduction of the research, climate variables were monitored: relative air humidity (RH%), accumulated rainfall (R mm), mean atmospheric temperature (AT °C), and mean solar radiation (SR MJ m<sup>2</sup>), as illustrated in Figure 1A and B.

**Comment [D4]:** was the study and the writing of the research paper carried out in one month. I don't think so. Kindly correct.

Is there a reason for conducting the study in April, related to climate conditions?



**Figure 1. Climate variables. (A) relative humidity (RH) and rainfall (R), (B) atmospheric temperature (AT) and solar radiation (SR), \* days the evaluations were performed. Jaboticabal, SP, 2015.**

### 2.2 Experimental Design

The research was conducted in a completely randomized design (DIC), and the treatments represented by three forest fragments [16] with three replicates. To define the sample unit, the fragments were divided into three parts of ≈240 m<sup>2</sup>, where each one represented a repetition. In each fragment, three visits were carried out at different times, the first being at

08:00, the second at 12:00, and the third at 16:00 hours, in a randomized way between the sample units.

## 2.3 Survey of Data

In the analyzed fragments, variables inherent in the frequency of occurrence of fauna and flora were included in the taxonomic groups: insects (F-INS), amphibians (F-AMP), birds (F-BIR), mammals (F-MAM), reptiles (F-REP), trees (F-TRE), and these are condensed into the variable total species frequency (F-TES). The data obtained from these variables consisted of *in loco* observation. Therefore, these were considered as "clues", in order to facilitate the visualization of copies of the groups; indicators of their existence, such as the diversity of leaves, flowers and fruits that can serve as food and water; trees and soil for shelter; besides aptitude for hunting and coexistence of populations. The research was classified as exploratory [17], of the qualitative type [18].

The collection of vegetation information was carried out based on specialized literature [19], and two species are commonly found in the transition areas of the Atlantic Forest and Cerrado, mainly because they represent the vegetation of the State of São Paulo. The species chosen were araticum-de-terra-fria (*Annonaemarginata* (Schlttdl.) H. Rainer) and dairy (*Tabernaemontanacatharinensis* A. DC.). This categorization was performed aiming at greater precision in the visualization and obtaining of the data [20].

## 2.4 Statistical Analysis

The original data of the dependent variables were transformed into a sine arc of  $\sqrt{(x/100)}$ , to normalize the distribution of the deviations [21], then subjected to analysis of variance using the 5% probability F test. For the significant variables, the Tukey test was applied for multiple comparisons of averages [22], in order to detect differences between the fragments. Subsequently, the original data were standardized and subjected to multivariate exploratory analysis, using cluster analysis (Ward's method) and Principal Components (PCA). The results of the multivariate analysis were expressed through tables, dendrogram and biplot [23].

## 3. RESULTS AND DISCUSSION


It was verified that there were significant differences ( $P < .01$ ) between the analyzed fragments for the variables, total frequency of species (F-SPE), F-INS, F-BIR, F-MAM, and ( $P = .05$ ), F-AMP, while for frequencies of occurrence of F-REP and F-TRE no significant differences were found ( $P = .05$ ) as summarized in Table 1.

**Table 1. Summary of variance analyses for the total frequency of species (F-TSP), insects (F-INS), amphibians (F-AMF), birds (F-BIR), mammals (F-MAM), reptiles (F-REP) and trees (F-TRE). Jaboticabal, SP, 2015.**

F.V.	GL	Medium Squares						
		F-TSP	F-INS	F-AMF	F-BIR	F-MAM	F-REP	F-TRE
Fragments	2	144.81**	99.74**	33.88*	57.16**	43.03**	3.66 <sup>ns</sup>	2.54 <sup>ns</sup>
Residue	6	8.80	9.10	6.45	5.16	1.99	7.32	0.63
CV (%)		17.19	32.25	68.30	26.68	32.86	212.13	10.49

\*, \* and <sup>ns</sup> - significant at 1 and 5% probability of error and not significant by the Fischer test, F.V. - sources of variation, GL - degrees of freedom and CV - coefficient of variation.


By analyzing the total frequency of verified species, it was possible to verify that fragment two (FRA-2), expressed superiority of 26.7 e 80.0% in relation to the fragments (FRA-1) it's three (FRA-3), though FRA-2 and FRA-3 have not differed statistically from one another, with averages of 11 and 15. The FRA-1 was superior in 72.7% when compared with FRA-3, with an average frequency of 3 (Figure 2A). The same behavior was observed when the F-INS was analyzed, having recorded averages of 4.3, 5.3 and 0.6 for FRA-1, FRA-2 and FRA-3, respectively, with percent differences of 86.0 and 88.7% when comparing FRA-1 and FRA-2 with FRA-3 (Figure 2B).



**Figure 2. Total species frequency (A), with minimum significant difference (MSD) of 7.43 and frequency of insects (B), with MSD of 7.53, verified in three forest fragments. Mean data of three untransformed sine-arc replicates of the  $\sqrt{(x/100)}$ . Jaboticabal, SP, 2015.**

Forest fragments, given the nature of their classification, express structural differences perceptible to the animals, in order to interfere with their behavior [24]. These authors attribute these behavioral changes, above all, to changes in the natural habitat of the species, while [25] contribute to changes in predation patterns. In a natural remnant fragment, the biological richness is undoubtedly superior to that of anthropized areas [14], reducing the incidence of solar radiation and temperature, increasing the relative air humidity and, thus, favoring the development of several species of fauna and flora [26]. Even in man-altered areas, some species can adapt and take advantage of this situation [27], justifying the frequency of species verified in natural regeneration areas in this work. Among the various species indicative of the quality of ecosystems, insects are often mentioned. For example, beetles [28, 29], caterpillars [30], bee species [31], and ants [32, 33, 6], are often quantified to express the level of environmental disorder based on how often they occur in environments.


Regarding the F-AMP, it was verified that FRA-2 had a higher mean (1.3), although it did not differ significantly from FRA-1 that had an average of 1, whereas FRA-3 expressed average 0, differing from FRA-2 being calculated a percentage difference of 100% between these two fragments (Figure 3A). For the variable F-BIR, this was also found to be superior to FRA-2, with an average 4.6, although this did not differ from FRA-1, with an average 2.6, with FRA-2 higher in 86.9% to FRA-3, where the mean was 0.6 (Figure 3B).



**Figure 3. Frequency of amphibians (A), with MSD of 6.37 and birds (B), with MSD of 5.69, verified in three forest fragments. Mean data of three replicates untransformed in sine-arc  $\sqrt{(x/100)}$ . Jaboticabal, SP, 2015.**

Rapid assessments and biotic integrity indexes, although generally based on a single taxonomic group, are effective methods for assessing biodiversity conservation [15]. These researchers confirm the longing for this research by suggesting that multiparameter-based assessments, for example, the frequency of occurrence of amphibians and birds, provide a more robust assessment of environmentally disturbed forest fragments. Studying the distribution of amphibians, reptiles, birds and mammals, [14] mentions that anthropic habitats are unsuitable for these species, while natural remnants and reforested fragments are potentially habitable, explaining the high frequency of amphibians and birds in FRA-2, followed by intermediate frequencies in FRA-1 and critical values evidenced in FRA-3 in this research. It should be noted that amphibians are one of the most endangered animal classes, mainly because of their sensitivity to environmental changes (for example, habitat destruction, climate change, as well as the reduction of air humidity, or the emergence of new pathogens, such as the quitridio fungus, *Batrachochytrium dendrobatidis*) due to its dependence on water and its permeability of the skin [34]. The distribution results of birds verified by [15] corroborate the findings of this research, certainly due to the characteristics of a particular reduction of the presence of ornithopters in places of intense antropic activity. [24], studying bird species, reported that, although the vast majority of bird species are classified as highly and moderately sensitive to environmental disturbances, there are, although in a smaller number, less sensitive species, justifying the occurrence of birds in the FRA-3 of this study.


No significant difference was recorded between the means 1.0 and 1.6 of the FRA-1 and FRA-2, respectively, when analyzed against the variable F-MAM, however, these two fragments differed significantly from the FRA-3, where there was no presence of mammals, characterizing the superiority of 100% of FRA-1 and FRA-2 in relation to FRA-3 (Figure 4A). For the variables F-REP and F-TRE, no significant differences were found (Figure 4B), which can be justified by the high variation of the original data, reflecting a high coefficient of variation of 212.13% for F-REP absence of normal distribution of data F-REP and F-TRE.



**Figure 4. Frequency of mammals (A), with MSD of 3.54, reptiles and trees (B), with MSDs of 6.78 and 1.99, respectively, verified in three forest fragments. Mean data of three untransformed sine-arc replicates of the  $\sqrt{(x/100)}$ . Jaboticabal, SP, 2015.**

Economic interests, to meet the demands of the growing population, have motivated predatory hunting, animal trafficking, forest deforestation, and expansion of arable land [35]. These researchers add that, fragmented forests tend to harbor fewer mammals compared with intact areas. It should be noted that the distribution dynamics of mammals in fragmented areas is also associated with their size. In fact, [36] report that the population of small rodent mammals can be increased in areas where the frequency of large mammals is reduced. In a complementary sense, [37] explain that changes in the distribution of mammals can be influenced by increasing land occupation for agriculture and livestock, as well as suppression of part of vegetation, alteration of hydrological cycles, burning regime and nutrient cycling in ecosystems. The nonoccurrence of differences between the fragments for the frequency of trees and reptiles can be justified by the fact that local climatic conditions favor the propagation and development of the trees, providing an adequate ecosystem for the occurrence of reptiles in the area of the three fragments studied [38].

On the basis of the Euclidean Distance used to summarize the homogeneity between the experimental units within the groups and heterogeneities between the groups, there were two main groups, the first group being represented by fragment three (FRA-3) and the second group by fragments one (FRA-1) and two (FRA-2), denoting the dissimilarities between the groups based on hierarchical grouping (Figure 5).



**Figure 5. Dendrogram of dissimilarities between three forest fragments (F). Jaboticabal, SP, 2015.**

In a research to compare two multivariate methodologies in the study of similarities between fragments of Atlantic forest, [39] point out that there is dissimilarity between groups of forest fragments, emphasizing that fragment groupings are due to the similarities of their variables, justifying these similarities due to their geographical proximity. [40] also observed that floristic similarity decreased with increasing distance between areas, in agreement with the ideas of [41] and [42], according to which geographical proximity would be the only reliable factor to predict the similarity between areas. These evidences allow us to infer that, due to the high geographic proximity of the fragments investigated in this research, the differences observed are due to the particular characteristics of these fragments, mainly anthropization in fragments of natural regeneration and reforestation.

It is observed in Table 2, that the set of seven categories (variables) analyzed was summarized in two latent variables (constructs), called Principal Components 1 ( $PC_1$ ) and 2 ( $PC_2$ ), which were selected based on eigenvalues, 4.83 and 1.26 because they were  $\geq 1$ , satisfying the criterion of Kaiser-Meyer-Olkin (KMO).


**Table 2. Eigenvalues (AV), relative variance ( $S^2_r$ ) and absolute ( $S^2_a$ ) and variable loads. Jaboticabal, SP, 2015.**

PCs	Variable Loads									
	AV	$S^2_r$ (%)	$S^2_a$ (%)	F- TES	F-INS	F- ANF	F- AVE	F- MAM	F- REP	F- ARV
$PC_1$	4.83	69.07	69.07	-0.98	-0.93	-0.91	-0.96	-0.85	0.09	-0.73
$PC_2$	1.26	18.03	87.11	0.17	0.25	-0.06	0.19	-0.18	0.97	-0.38

*PCs: Principal Components.*

The  $PC_1$  and  $PC_2$  account for 87.1% of the total cumulative variance, with  $PC_1$  accounting for 69.07% of this total, with  $PC_2$  accounting for 18.04%. It is observed that the FRA-3

presented greater dissimilarities compared with the FRA-2, with the FRA-1 occupying intermediate position (Figure 6).



**Fig. 6. Two-dimensional projection (Biplot) of three forest fragments (F) and representatives of fauna and flora in two Principal Components (PC<sub>1</sub> and PC<sub>2</sub>). Jaboticabal, SP, 2015.**

Analyzing forest fragments through Principal Component Analysis, [35] report that the first two components account for 56% of the total variation in mammalian distribution among the sampled sites. These researchers point out that the lower incidence of mammalian species in altered areas can be explained by the hunter's pursuit of animals, especially game animals and those that cause damage to agricultural crops, while trees and nontarget species of hunting tend to be seen more frequently in anthropized areas.

On the basis of the results, the use of Principal Component Analysis is justified, since it provides a structural simplification of the original data. In fact, in the research carried out by [39], 462 dimensions were reduced in 10 Principal Components resulting from linear combinations between original variables. Therefore, these authors report that using the first 10 PCs is as efficient as the use of the 462 initial variables with regard to the explanation of the variance. Thus, the use of two PCs in this research was sufficient enough to explain the variance under study.

On the basis of this information, it is believed that the divergences evidenced between the fragments of this research can be a reflection of the anthropic actions, especially of the illegal hunting that can occur in these places. For [39], each fragment exhibits a species composition that appears to result from a series of factors that varied differently over time and space. Perhaps that is why it is so difficult to establish these areas. This difficulty, however, indicates that each fragment presents a set of its own characteristics, which emphasizes its importance in terms of conservation.



#### 4. CONCLUSION

There are differences in the biodiversity of the forest fragments analyzed, highlighting the areas of natural remnants and reforestation with greater biological diversity, to the detriment of the natural regeneration areas with insufficient biological indicators, denoting adequacy of the first two fragments and inadequacy of the latter with respect to the potential of use as a legal reserve.

Two groups of environments were evidenced according to the potential hierarchy for use as legal reserve, the first group being characterized as inadequate and the second as adequate. Of the seven analyzed variables, six were considered essential to the correct evaluation of the environments.

#### COMPETING INTERESTS

The authors have not declared any conflict of interests.

#### AUTHORS' CONTRIBUTIONS

This work was carried out with the collaboration of all authors. All authors read and approved the final manuscript.

#### REFERENCES

1. Ashby MF. What is a "Sustainable Development"? In: Ashby MF, editor. *Materials and Sustainable Development*. Amsterdã: Butterworth-Heinemann; 2016.
2. Powell LL, Stouffer PC, Wolfe JD, Johnson EI, Hines JE, Nichols JD. Heterogeneous movement of insectivorous Amazonian birds through primary and secondary forest: A case study using multistate models with radiotelemetry data. *Biological Conservation*.2015;188(1):100-108. a
3. Arcilla N, Holbechc LH, O'Donnell S. Severe declines of understory birds follow illegal logging in Upper Guinea forests of Ghana, West Africa. *Biological Conservation*. 2015;188(1):41-49.
4. Cordeiro N, Dampf CJ, Borghesio L, Joho MP, Monoski TJ, Mkongewa VJ. Forest fragmentation in an African biodiversity hotspot impacts mixed-species bird flocks. *Biological Conservation*.2015;188(1):61-71.
5. PavlackyJr DC, Possingham HP, Goldizen AW. Integrating life history traits and forest structure to evaluate the vulnerability of rainforest birds along gradients of deforestation and fragmentation in eastern Australia. *Biological Conservation*.2015;188(1):89-99.
6. Rocha WO, Dorval A, Peres Filho O, Vaez C dos, Ribeiro ES. Formigas (Hymenoptera: Formicidae) bioindicadores de degradaçãoambientalemPoxoréu, MatoGrosso, Brasil. *Floresta e Ambiente*.2015;22(1):88-98. Portuguese.
7. Velazco SJE, Galvão F, Keller HA, Bedrij NA. Florística e fitossociologia de uma floresta estacional semidecidual, reservaprivada de Osununú-Misiones, Argentina. *Floresta e Ambiente*.2015;22(1):1-12. Portuguese.

8. Silvério Neto R, Bento M de C, Menezes SJM da C de, Almeida FS. Caracterização da cobertura florestal de unidades de conservação da Mata Atlântica. *Floresta e Ambiente*.2015;22(1):32-40. Portuguese.
9. Fiorentin LD, Téó SJ, Schneider CR, Costa RH, Batista S. Análise florística e padrão espacial da regeneração natural em área de floresta ombrófila mista na região do Caçador SC. *Floresta e ambiente*.2015;22(1):60-70. Portuguese.
10. Powell LL, Cordeiro NJ, Stratford J. Ecology and conservation of avian insectivores of the rainforest understory: A pantropical perspective. *Biological Conservation*.2015;188(1):1-10. b
11. Vogel HF, Cardoso O, Watzlawick LF, Campos JB. Pesquisas em unidades de conservação urbanas no Paraná: conhecimentos raramente divulgados ou aplicados. *Ambiência - Revista do Setor de Ciências Agrárias e Ambientais*.2015;11(1):75-93. Portuguese.
12. Summers DM, Bryan BA, Nolan M, Hobbs TJ. The costs of reforestation: A spatial model of the costs of establishing environmental and carbon plantings. *Land Use Policy*.2015;44(1):110-121.
13. Cunningham SC, Mac Nally R, Baker PJ, Cavagnaro TR, Beringer J, Thomsonb JR, Thompsonb RM. Balancing the environmental benefits of reforestation in agricultural regions. *Perspectives in Plant Ecology, Evolution and Systematics*.2015;17(4):301-317.
14. Botello F, Sarka S, Sánchez-Cordero V. Impact of habitat loss on distributions of terrestrial vertebrates in a high-biodiversity region in Mexico. *Biological Conservation*.2015;184(1):59-65.
15. Medeiros HR, Bochio GM, Ribeiro MC, Torezan JM, Anjos L. Combining plant and bird data increases the accuracy of an Index of Biotic Integrity to assess conservation levels of tropical forest fragments. *Journal for Nature Conservation*.2015;25(1):1-7.
16. Mussury RM, Scalón S de PQ, Gomes AA, Batista MR, Scalón Filho H. Flutuação populacional da mesofauna em fragmentos de mata na região de Dourados MS. *Ciência e Agrotecnologia, Lavras*.2008;32(2):645-650. Portuguese.
17. Laake P, Fagerland MW. Statistical Inference. In: Laake P, Benestad H, Olsen BR, editors. *Research in Medical and Biological Sciences*. 1st ed. Amsterdã: Academic Press; 2015.
18. Moen K, Middelthon AL. Qualitative Research Methods. In: Laake P, Benestad H, Olsen BR, editors. *Research in Medical and Biological Sciences*. 1st ed. Amsterdã: Academic Press; 2015.
19. Lorenzi H. *Árvores Brasileiras: Manual de identificação e cultivo de plantas arbóreas nativas do Brasil*. Nova Odessa: Instituto Plantarum; 2009. Portuguese.
20. Depoy E, Gitlin LN. Collecting data through measurement in experimental-type research. In: Depoy E, Gitlin LN *Introduction to Research*. 5th ed. Amsterdã: Mosby; 2016.
21. Pimentel-Gomes F. *Curso de estatística experimental*. 14th ed. Piracicaba: Degaspari; 2000. Portuguese.

22. Santos JW, Almeida FAC, Beltrão NEM, Cavalcante FB. Estatística Experimental Aplicada. 2nd ed. Campina Grande: EmbrapaAlgodão/UFCG; 2008.Portuguese.
23. Hair JF, Black WC, Babin BJ, Anderson RE, Tatham RL. Análisemultivariada de dados. Porto Alegre: Bookman; 2009.Portuguese.
24. Stratford JA, Stouffer PC. Forest fragmentation alters microhabitat availability for Neotropical terrestrial insectivorous birds. *Biological Conservation*.2015;188(1):109-115.
25. Visco DM, Sherry TW. Increased abundance, but reduced nest predation in the chestnut-backed antbird in costa rican rainforest fragments: surprising impacts of a pervasive snake species. *Biological Conservation*.2015;188(1):22-31.
26. Pollock HS, Cheviron ZA, Agin TJ, Brawn JDAC. Absence of microclimate selectivity in insectivorous birds of the neotropical forest understory. *Biological Conservation*.2015;188(1):116-125.
27. Boyle WA, Sigel BJ. Ongoing changes in the avifauna of La Selva Biological Station, Costa Rica: Twenty-three years of Christmas Bird Counts. *Biological Conservation*.2015;188(1):11-21.
28. Figueiras BKC, Tabarelli M, Leal IR, Vaz-de-Mello F, Iannuzzi L. Dung beetle persistence in human-modified landscapes: Combining indicator species with anthropogenic land use and fragmentation-related effects. *Ecological Indicators*.2015;55(1):65-73.
29. Salomão RP, Iannuzzi L. Dung beetle (Coleoptera, Scarabaeidae) assemblage of a highly fragmented landscape of Atlantic forest: from small to the largest fragments of northeastern Brazilian region. *RevistaBrasileira de Entomologia*.2015;59(1):126-131.
30. Maguire DY, Bennett EM, James PMA, Buddle CM. Landscape connectivity and insect herbivory: A framework for understanding tradeoffs among ecosystem services. *Global Ecology and Conservation*.2015;4(1):73-84.
31. Pereira SAN, Sousa CS. Levantamento da fauna de abelhas no município de montecarmelo-MG. *Getec*.2015;4(7):11-24.Portuguese.
32. De La Mora A, García-Ballinas JA, Philpott SM. Local, landscape, and diversity drivers of predation services provided by ants in a coffee landscape in Chiapas, Mexico. *Agriculture, Ecosystems and Environment*.2015;201(1):83-91.
33. Komonen A, Övermark E, Hytönen J, Halme P. Tree species influences diversity of ground-dwelling insects in afforested fields. *Forest Ecology and Management*.2015;349(1):12-19.
34. Grosjean S, Cruaud C, Ohler A, Hassanin A, Chuaynkern Y. Improving biodiversity assessment of anuran amphibians using DNA barcoding of tadpoles. Case studies from Southeast Asia. *ComptesRendusBiologies*.2015;338(1):351-361.
35. Meyer NFV, Helen JE, Ricardo M, Frank VL, Yorick L, David RO, Chantal BFV, Andrew DC, Clayton KN, Patrick AJ. "An assessment of the terrestrial mammal communities in

- 436 forests of Central Panama, using camera-trap surveys". *Journal for Nature*  
437 *Conservation*.2015;(6):28-35.
- 438
- 439 36. Galetti M, Bovendorp SR, Guevara R. Defaunation of large mammals leads to an  
440 increase in seed predation in the Atlantic forests. *Global Ecology and*  
441 *Conservation*.2015;3(1):824-830.
- 442
- 443 37. Ripple WJ, Newsome TM, Wolf C, Dirzo R, Everatt KT, Galetti M, Hayward MW, Kerley  
444 GIH, Levi T, Lindsey PA, Macdonald DW, Malhi Y, Painter LE, Sandom CJ, Terborgh J,  
445 Valkenburgh BV. Collapse of the world 's largest herbivores Collapse of the world 's  
446 largest herbivore. *Science Advances*.2015;(5):1-12.
- 447
- 448 38. Paula DM de, Groeneveld J, Huth A. Tropical forest degradation and recovery in  
449 fragmented landscapes — Simulating changes in tree community, forest hydrology and  
450 carbon balance TL - 3. *Global Ecology and Conservation*.2015;(3):664-77.
- 451
- 452 39. Ferreira RLC, Mota AC, Silva JAA, Marangon LC, Santos ES. Comparação de  
453 duas metodologias multivariadas no estudo de similaridade entre fragmentos de  
454 floresta atlântica. *Revista Árvore*.2008;32(3):511-521. Portuguese.
- 455
- 456 40. Jacquemyn H, Butaye J, Dumortier M, Hermy M, Lust N. Effects of age and distance on  
457 the composition of mixed deciduous forest fragments in an agricultural landscape.  
458 *Journal of Vegetation Science*.2011;(12):635-642.
- 459
- 460 41. Condit R. Defining and mapping vegetation types in mega-diverse tropical forests.  
461 *Trends in Ecology and Evolution*.1996;11(1):4-5.
- 462
- 463 42. Cook SA. Diversity of approaches to the study of species richness. *Trends in Ecology*  
464 *and Evolution*.1998;13(9):340-341.