

## Original Research Article

# Anti-aging activity of *Xylaria striata* in *Drosophila melanogaster*

### ABSTRACT

**Aims:** To evaluate the application potential of *Xylaria striata* Pat. in anti-aging field.

**Methodology:** Extracting the fruit body of *Xylaria-X. striata* by ultrasonic-assisted extraction method. In this study, *Drosophila melanogaster* was used as an anti-aging organism material. After feeding with different concentrations of extract, the natural survival time, survival time under oxidative stress and survival time under UV irradiation of *Drosophila-D. melanogaster* were all measured. In addition, the *in vivo* activity of SOD, CAT and MAD, protein concentration and body weight were determined to evaluate the anti-aging effect of ethanol extract from *Xylaria-X. striata*.

**Results:** The results showed that the ethanol extract of *Xylaria-X. striata* could extend the lifespan of *Drosophila-D. melanogaster* under both irradiation and oxidative stress condition. And the ethanol extract could enhance the activity of CAT in *Drosophila-D. melanogaster*, especially at concentration of 50 µM, and the content of MAD in *Drosophila-D. melanogaster* were decreased significantly.

**Conclusion:** This study clarified the anti-aging activity of *Xylaria-X. striata* in *Drosophila-D. melanogaster* and it would provide some theoretical basis for its further development and utilization in anti-aging drugs and health food.

**Keywords:** *Xylaria striata*; anti-aging; *Drosophila melanogaster*; lifespan.

### 1 INTRODUCTION

The higher fungi contain compounds with many novel structure, which have variety and remarkable biological activity<sup>[1-3]</sup>. Because of potential medicinal and economic value, they have drawn an increasing attention and become a highlight research field of science and industry in recent years.

The genus *Xylaria*, a big family of mycomycetes, has great value in application according to a large amount of literature reports. There were many kinds of compounds such as terpenoids, sterols, alkaloids, polyketones, polysaccharides, cyclic peptides and carboxylic acids<sup>[4-6]</sup> had been isolated from the *Xylaria* genus, which exhibited antioxidant, antimicrobial, antitumor, enzyme inhibitory abilities. *Xylaria-X. striata* Pat. 1887, one kinds of summer-living mushroom belong to the family of *Xylaria*, mainly grows on decayed barks and lived roots of broad-leaved woodland. It was used as folk medicine in China<sup>[7]</sup>. After finding its diverse biological activities in the preliminary screening of large numbers of edible and medicinal mushrooms, our research group started a systematic study on *Xylaria striata* from cultivation method to chemical components since 2013<sup>[8-9]</sup>. These research results clarified its effect of anti-animal and plant pathogens<sup>[10]</sup>, anti-tumor<sup>[11]</sup>, and promoting sleep<sup>[12]</sup>.

Furthermore, another mushroom from the same genus named *Podosordaria nigripes* (Klotzsch) P.M.D. Martin (Syn: *Xylaria nigripes* (Klotzsch) Cooke has been widely used to prevent and treat senile diseases in China. It can promote the effect of antidepressants<sup>[14,13]</sup>

**Comment [u1]:** It is very well-marked that this study is acceptable with minor revision and useful for publish in this journal. The discussion section can be developed with literature data.

Note: Please, I want to see manuscript again after the corrections.

In addition to, please add some papers and you can use them in manuscript.

Please see below for papers.

2. Sevindik M. Investigation of Antioxidant/Oxidant Status and Antimicrobial Activities of *Lentinus tigrinus*. *Advances in pharmacological sciences*, 2018. DOI:10.1155/2018/1718025

3. Sevindik M. Investigation of oxidant and antioxidant status of edible mushroom *Clavariadelphus truncatus*. *The Journal of Fungus*. 2018; 9(2):165-168

**Formatted:** Font: Italic

**Field Code Changed**

**Field Code Changed**

**Field Code Changed**

**Field Code Changed**

**Formatted:** Font: Not Italic

**Formatted:** Font: Italic

**Formatted:** Font: Italic

and alleviate depressive symptoms in patients with epilepsy<sup>[14]</sup>. Hence, the main objectives of this study were to evaluate the anti-aging activity of *Xylaria-X. striata* using *Drosophila melanogaster* as the model organism by assessing the lifespan of flies, the activity of SOD and CAT, the content of MDA and protein and its stress resistant ability under H<sub>2</sub>O<sub>2</sub> and UV irradiation treatment.

Formatted: Font: Italic

Field Code Changed

## 2 MATERIALS

### 2.1 Biomaterial

Field Code Changed

*Xylaria-X. striata* was obtained in 2013 from Qingyi town, Mianyang, Sichuan province of China, and identified by professor Xin-sheng He of microbiology, Southwest University of Science and Technology. The voucher specimen was preserved in the Microbiology Laboratory of the same university. *Drosophila-D. melanogaster*, reared under the condition of temperature 25 ± 1 °C, humidity 60%-80%.

Field Code Changed

### 2.2 Reagents

Protein content determination kit, superoxide dismutase (SOD) determination kit, catalase (CAT) determination kit, and malondialdehyde (MDA) content determination kit are all purchased from Nanjing Jiancheng bioengineering institute; Corn flour, agar and yeast extract was purchased from Beijing Aobaoing Bio-Tech Co., Ltd.; Ethyl ether, ethanol, 30% hydrogen peroxide and other reagents were purchased from ChengDu Chon Chemicals Co., Ltd.; All reagents are analytical pure grade and pure water is made in the laboratory.

### 2.3 Equipments

Biochemical incubator (SPX- 80B, Shanghai Kuntian instrument Co., Ltd.), Thermo Scientific Microplate Reader (Multiskan Spectrum, Thermo Fisher), Ultraviolet lamp (LP-GDZJ40W, LONGPRO CO., Ltd.), Ultrasonic cleaner (KQ-200KDE, Kunshan Ultrasonic Instrument Co., Ltd.).

## 3 METHODS

### 3.1 Sample pretreatment

The cultivated fruiting body<sup>[15]</sup> of *Xylaria-X. striata* was dried at 50 °C and ethanol-assisted ultrasonic extraction process was carried out at the ratio of material to liquid (1:30 g:mL) for three times, 10 min of each. Then, the extract was obtained by vacuum distilling of the filtered supernatant at 65 °C and preserved at 4 °C until use.

### 3.2 Diet preparation and Fly husbandry

The wild-type *Drosophila-D. melanogaster* Canton-S (CS) flies, selected as the experiment organism for later assay, were reared at temperature 25 ± 1 °C, humidity 60%-80%. Before the experiment, *Drosophila-D. melanogaster*s were mass reared in 500 mL erlenmeyer flask containing 100 mL of standard cornmeal diet<sup>[16]</sup> (13% cornmeal w/v, 8% sugar w/v, 2.4% Agar w/v, 1.6% yeast extract w/v and 0.4% acetic acid v/v). To avoid overcrowding, 200 flies of each bottle was enough. When the progeny drosophila was going to hatch, all adult flies would be transferred to the new culture bottle. Afterwards, the flies eclosed within 24 h were collected and segregated according to their sex.

Field Code Changed

### 3.3 Longevity assay

Refers to Proshkina *et al*<sup>[17]</sup>, 10 newly eclosed male flies, reared on a standard cornmeal medium with ethanol extract from *Xylaria-X. striata*. The extract was dissolved in water and mixed with the standard diet at final concentrations of 50, 100 and 200 µM. Negative control diets contained only water while positive control contained ascorbic acid at final

concentrations of 100  $\mu\text{M}$ . During the rearing process, diet was replaced with fresh medium every five days and the numbers of alive flies were recorded every two days. The test was stopped until all flies were dead. All the treatments were carried out with 3 replicate. Median lifespan was calculated using the method reported by Kaplan–Meier<sup>[18]</sup> previously.

### 3.4 Oxidative stress resistance

The situation of group dividing is the same as Longevity assay. Newly eclosed male flies were fed on standard diet with the extract from *Xylaria-X. striata* for 25 days, followed by an oxidative challenge of 9%  $\text{H}_2\text{O}_2$  in 6% glucose solution on filter paper strips<sup>[19]</sup>. The number of death was recorded every 4 h until all flies were dead. Median lifespan was calculated as the same as 3.3.

### 3.5 Irradiation resistance

The flies were treated at a constant distance of 10 cm from the ultraviolet lamp and irradiation time was 20 min<sup>[20]</sup>. We strictly controlled the surrounding temperature at 25 $^{\circ}\text{C}$  to decrease heat stroke from the lamp during irradiation. Then, the flies were transferred to vials. Dead flies were counted every day until all flies died. Life span was studied in the same way.

### 3.6 SOD、CAT、MDA、Protein content and weight assays

After 25 days of feeding, flies were transferred into empty tube for 2 hours. After anesthesia, the flies were homogenized in an ice bath at the ratio of weight to normal saline at 1:19 (g:mL). The supernatant was obtained after the homogenate was centrifuged in 2500  $\text{r} \cdot \text{min}^{-1}$  for 20 min at 4 $^{\circ}\text{C}$ . The activity of CAT(catalase) and SOD(superoxide dismutase), the content of MDA(malondialdehyde) and protein were determined by the Thermo Scientific Microplate Reader according to the instruction described by the kit. Flies were anesthetized and weighed every ten flies. For each concentration, 3 replicate were set up.

### 3.7 Statistical Analysis

All the data collected were repeated 3 times, and the data were displayed in the form of means $\pm$ SD. The statistical analysis was performed with SPSS 20.0 software, and the difference analysis between groups was performed by One-way ANOVA analysis. Significant differences were expressed by: \*  $P < 0.05$ ; \*\*  $P < 0.01$ ; \*\*\*  $P < 0.001$ .

## 4 RESULTS AND DISCUSSION

### 4.1 Longevity assay

*Drosophila-D. melanogaster* has been widely used in anti-aging experiments as a model organism because of its short lifespan and easy reproduction<sup>[21]</sup>. The anti-aging ability of the drugs or its toxicity can be reflected at a certain level by comparing the lifespan of *Drosophila-D. melanogaster* before and after feeding the drugs. As shown in Table 1, the ethanol extract of *Xylaria-X. striata* can significantly shorten the mean lifespan of *Drosophila-D. melanogaster*, and with the increasing of the dosage, the more significant the shorten effect is. In addition, the maximum lifespan and half survival time of *Drosophila-D. melanogaster* were also reduced. Therefore, there may be some biologically toxic substances in the ethanol extracts of *Xylaria-X. striata*. It has been reported by Yuan et al.<sup>[11]</sup> that *Xylaria-X. striata* has an inhibitory effect on plant and animal pathogens, which can also prove that it may contain some toxic substances.

Table 1 The effect of *Xylaria-X. striata* on lifespan

| Treatment | Concentration<br>( $\mu\text{M}$ ) | Max $\pm$ SD<br>(day) | H $\pm$ SD<br>(day) | M $\pm$ SD<br>(day) |
|-----------|------------------------------------|-----------------------|---------------------|---------------------|
|-----------|------------------------------------|-----------------------|---------------------|---------------------|

Formatted: Subscript

Formatted: Subscript

|                           |     |          |             |            |
|---------------------------|-----|----------|-------------|------------|
| Control                   | -   | 45.3±1.2 | 35.7±1.6    | 30.6±2.1   |
| <i>Xylaria-X. striata</i> | 50  | 32±0.9*  | 20±2**      | 28.2±0.15  |
|                           | 100 | 41.3±6.2 | 31.3±4.7    | 34.1±3.7   |
|                           | 200 | 38±4.3   | 15.3±1.9*** | 24.2±2.1** |
| Ascorbic acid             | 100 | 48.9±3.2 | 39.0±3.4**  | 37.7±0.7*  |

Note: Max, maximum lifespan, days; M, mean lifespan, days; H, half survival lifespan; \*p < 0.05, \*\*p < 0.01, \*\*\*p < 0.001

Formatted: Font: Italic

## 4.2 Oxidative stress resistance

*In vivo*, hydrogen peroxide reacts with oxygen to produce hydroxyl radicals, which can induce acute oxidative stress damage and shorten the survival time of *Drosophila-D. melanogaster*<sup>[22]</sup>. As shown in Table 2, the ethanol extract of *Xylaria-X. striata* could prolong the maximum lifespan, half survival time and mean lifespan of *Drosophila-D. melanogaster* under oxidative stress. When the concentration of extract was 100 µM, the maximum lifespan and the mean lifespan were obviously extended, and the prolongation of the mean lifespan was extremely significant at the concentration of 200 µM.

Field Code Changed

Table 2 The effect of oxidative stress on lifespan

| Treatment                 | Concentration (µM) | Max±SD (h) | H±SD (h) | M±SD (h)   |
|---------------------------|--------------------|------------|----------|------------|
| Control                   | 0                  | 23.3±3.1   | 16.7±1.1 | 17.2±0.6   |
| <i>Xylaria-X. striata</i> | 50                 | 22±2       | 17±1.3   | 17±0.6     |
|                           | 100                | 29.3±3.3*  | 18±4     | 18.3±0.3*  |
|                           | 200                | 21.3±2.3   | 15.3±1.1 | 18.7±0.3** |
| Ascorbic acid             | 100                | 26±2       | 18.7±2.3 | 18.5±0.5** |

Formatted: Font: Italic

Note: \*p < 0.05, \*\*p < 0.01, \*\*\*p < 0.001

## 4.3 UV irradiation resistance

Ultraviolet radiation can cause changes in some physiological functions and damages the body. Herein, we determined if the *Xylaria-X. striata* could repair UV irradiation damage by observing the lifespan changes of *Drosophila-D. melanogaster* under UV irradiation stress. According to a previous report, the sensitivity of *Drosophila-D. melanogaster* to ultraviolet light was closely related to the melanin content<sup>[23]</sup>. So, this method could also show the effect of the tested agent on melanin production in *Drosophila-D. melanogaster*. As can be shown in Table 3, a low dose of extract could significantly prolong the lifespan of *Drosophila-D. melanogaster* after UV irradiation. Medium, high dose groups could also prolong lifespan, but the effect was inferior to the low dose group. This may be due to the variety of substances in the extract. As the increasing of dose, the increasing biotoxicity resulted in a decrease in lifespan. In addition, we found ascorbic acid can't prolong the lifespan, which indicated that it unable repaired the radiation damage.

Table 3 Effect of UV irradiation on lifespan

| Treatment                 | Concentration (µM) | Max±SD (day) | H±SD (day) | M±SD (day) |
|---------------------------|--------------------|--------------|------------|------------|
| Control                   | 0                  | 12.5±0.7     | 6±1.3      | 5.2±0.4    |
|                           | 50                 | 17±1.4*      | 11.5±0.7*  | 10.4±2.3*  |
| <i>Xylaria-X. striata</i> | 100                | 15±1.4       | 10±2.8     | 8.5±1.5    |
|                           | 200                | 14.5±1.5     | 9.3±2.9    | 9±2.5      |

Formatted: Font: Italic

Ascorbic acid 100 12.5±0.7 9.5±0.7 9.4±0.4

Note: \*p < 0.05, \*\*p < 0.01, \*\*\*p<0.001

#### 4.4 SOD、CAT、MAD, protein content and weight assay

After feeding on the extract, the overall living condition can be indirectly reflected by determining the weight and protein content of *Drosophila-D. melanogaster*. In Table 4, there were no significant changes in the weight and protein content of flies.

Superoxide dismutase (SOD), one kind of free radical enzyme, can scavenge the superoxide anion in body, and prevent the body from being oxidized. Hydrogen peroxide reacts with oxygen can produce hydroxyl radicals, which are harmful to the body. However, hydrogen peroxide decomposition *in vivo* can be catalyzed by catalase (CAT) to reduce the generation of hydroxyl free radicals<sup>[24]</sup>. Hence, SOD and CAT are two kinds of representative enzymes in antioxidant system. The activities of these two enzymes can indirectly evaluate the free radical scavenging ability of the organism<sup>[25]</sup>.

Malondialdehyde (MDA) is a kind of lipid peroxidation product produced by the reaction of free radicals and unsaturated fatty acids. MDA residues in body can be further cross-linked with proteins and peptides to accelerate the aging of the body<sup>[26]</sup>.

As can be seen in Table 4, The ethanol extract of *Xylaria-X. striata* has no positive effect on SOD. But the extract at low dose can promote the activity of CAT to reduce the free radicals. Furthermore, the extract can also significantly decrease the content of MDA which is consistent with the content of free radicals. It is implied that the MDA content can also indirectly reflect the content of free radicals and the degree of lipid peroxidation. So, it is conducted that the ethanol extract of *Xylaria-X. striata* keeps body much away from peroxidation and prevent it from aging too quickly.

Table 4 Effect of *Xylaria-X. striata* on SOD/CAT/MDA

| Treatment                 | Concentration (μM) | Protein content (g/L) | CAT (U/mL)   | MDA (nmol/mL) | SOD (U/mg) | Weight (mg/10 flies) |
|---------------------------|--------------------|-----------------------|--------------|---------------|------------|----------------------|
| Control                   | 0                  | 1.08±0.16             | 61.91±5.92   | 2.65±0.40     | 43.21±8.5  | 6.7±0.76             |
| <i>Xylaria-X. striata</i> | 50                 | 1.32±0.23             | 87.47±11.34* | 0.9±0.27***   | 34.59±6.71 | 6.9±0.32             |
|                           | 100                | 1.37±0.22             | 69.46±6.94   | 1.75±0.11**   | 36.51±6.91 | 6.6±0.83             |
|                           | 200                | 1.3±0.23              | 58.49±10.34  | 1.03±0.26***  | 29.45±2.91 | 6.2±0.26             |
| Ascorbic acid             | 100                | 1.37±0.23             | 92.27±13.23* | 2.65±0.27     | 34.11±6.2  | 5.9±0.07             |

Note: \*p < 0.05, \*\*p < 0.01, \*\*\*p<0.001

#### 5 CONCLUSIONS

From the previous report that *Xylaria-X. striata* can promote pentobarbital-induced sleep by not only increasing the number of falling asleep and prolonging sleeping time but also reducing sleep latency<sup>[27]</sup>, this mushroom had potential to be as a functional food used in the field of geriatrics. Above all, the anti-aging activity of *Xylaria-X. striata* was evaluated by measuring the survival time under various conditions, the activity of SOD/CAT and the content of MAD/protein of *Drosophila-D. melanogaster*. The results showed that, although the ethanol extract of *Xylaria-X. striata* could shorten the lifespan of *Drosophila-D. melanogaster* under natural conditions which indicated its biologically toxic, it could extend the lifespan of flies in the longevity test of two stress models by repair both ultraviolet radiation damage and hydrogen peroxide oxidative stress damage. The results of enzyme activity and MAD content

Formatted: Font: Italic

Field Code Changed

showed that the ethanol extract of *Xylaria X. striata* could not only block the source of free radicals but also eliminate the reaction products of free radicals.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Zhao JD. Chinese fungi(The third volume). 1th ed. Beijing; Science Press, 1998.
2. Sevindik M. Investigation of Antioxidant/Oxidant Status and Antimicrobial Activities of *Lentinus tigrinus*. *Advances in pharmacological sciences*, 2018. DOI:10.1155/2018/1718025
3. Sevindik M. Investigation of oxidant and antioxidant status of edible mushroom *Clavariadelphus truncatus*. *The Journal of Fungus*. 2018; 9(2):165-168
4. Gao C, Luo J, Liu X, Ma L, Yuan XH. Recent advances in the studies of chemical compositions and bioactivities of the genus *Xylaria*. *Mycosystema*. 2016,35(07):767-781. DOI:10.13346/j.mycosystema.150061
5. Macias-Rubalcava M, Sanchez-Fernandez RE. Secondary metabolites of endophytic *Xylaria* species with potential applications in medicine and agriculture. *World Journal Of Microbiology & Biotechnology*. 2017, 33(01):15. DOI: 10.1007/s11274-016-2174-5
6. Song F, Wu SH, Zhai YZ, Xuan QC, Wang T. Secondary Metabolites from the Genus *Xylaria* and Their Bioactivities. *CHEMISTRY & BIODIVERSITY*. 2014, 11(05): 673-694. DOI: 10.1002/cbdv.201200286
7. He XS. Images of mushroom fungus in SiChuan. 1th ed. Bei Jing: Science Press; 2011.
8. Liu X, Gao C, Zhong YT, Yuan XH, He XS, Ma L. Optimization of submerged culture conditions for *Xylaria striata* mycelium. *Journal of Chinese medicinal materials*. 2014, 37(08):1317-21.
9. Lei CW, Yang ZQ, Zeng YP, Zhou Y, Huang Y, He XS, et al. *Xylatriasan A*, a new cytochalasan from the fungus *Xylaria striata*. *Natural product research*. 2018, 32(01):7-13. DOI: 10.1080/14786419.2017.1324959
10. Zhang CJ, Liu X, Zhu P, Liu XL, He XS. Antimicrobial Activities of Ethanol Extracts from Six Species of Higher Fungi against Plant Pathogen. *Journal of Anhui Agricultural Sciences*. 2013, 41(12):5289-5294. DOI: 10.13989/j.cnki.0517-6611.2013.12.108
11. Yuan XH, Zhang CJ, Zheng RL, Xu JR, Liu Z, He XS. Screening of Anti-tumor Activities of Ethanol Extracts from Six Species of Higher Fungi. *Journal of Southwest University of Science and Technology*, 2013, 28(3):95-97.
12. Lie CW. Studies on the chemical constituents of *Xylaria striata* and their bioactivities. *Southwest University of Science and Technology*, 2017.
13. Zheng W, Zhang YF, Zhong HQ, Mai SM, Yang XH, Xiang YT. Wuling Capsule for Major Depressive Disorder: A Meta-analysis of Randomised Controlled Trials. *East Asian archives of psychiatry : official journal of the Hong Kong College of Psychiatrists*. 2016, 26(3): 87-97.
14. Peng WF, Wang X, Hong Z, Zhu GX, Li BM, Li Z, et al. The anti-depression effect of *Xylaria nigripes* in patients with epilepsy: A multicenter randomized

- double-blind study. *Seizure-european Journal of Epilepsy*. 2015, 29: 26-33. DOI: 10.1016/j.seizure.2015.03.014
15. Hou CL, Sun XL, Tan J, Zhang CX, Yang DS, He XS. Study on Culture of *Xylaria striata* Fruit Body. *Food and Fermentation Sciences & Technology*. 2013,49(02): 44-46+102. DOI:1674-506X (2013) 02-0044-0003
16. Chattopadhyay D, Chitnis A, Talekar A, Mulay P, Makkar M, James J, et al. Hormetic efficacy of rutin to promote longevity in *Drosophila melanogaster*. *Biogerontology*. 2017,18(3): 397-411. DOI: 10.1007/s10522-017-9700-1
17. Proshkina E, Lashmanova E, Dobrovolskaya E, Zemskaya N, Kudryavtseva A, Shaposhnikov M, et al. Geroprotective and Radioprotective Activity of Quercetin, (-)-Epicatechin, and Ibuprofen in *Drosophila melanogaster*. *Frontiers in Pharmacology*. 2016, 7: 505. DOI: 10.3389/fphar.2016.00505
18. Kaplan EL, Meier P. Nonparametric estimation from incomplete observations. *Journal of the American Statistical Association*. 1958, 53(282): 457-481.
19. Bayliak MM, Lylyk MP, Gospodaryov DV, Kotsyubynsky VO, Butenko NV, Storey KB, et al. Protective effects of alpha-ketoglutarate against aluminum toxicity in *Drosophila melanogaster*. *Comparative biochemistry and physiology. Toxicology & pharmacology : CBP*. 2019, 217: 40-53. DOI:10.1016/j.cbpc.2018.11.020
20. Rajpurohit S, Schmidt PS. Latitudinal Pigmentation Variation Contradicts Ultraviolet Radiation Exposure: A Case Study in Tropical Indian *Drosophila melanogaster*. *Frontiers in Physiology*. 2019, 10: 84. DOI: 10.3389/fphys.2019.00084
21. Piper MDW, Partridge L. *Drosophila* as a model for ageing. *BIOCHIMICA ET BIOPHYSICA ACTA-MOLECULAR BASIS OF DISEASE*. 2018, 1864(09): 2707-2717. DOI: 10.1016/j.bbdis.2017.09.016
22. Niki E. Lipid peroxidation products as oxidative stress biomarkers. *Biofactors*. 2008, 34: 171-180.
23. Li HJ, Zhang SF, Li JX, Zhao Z. Anti-UV Irradiation Effect of Melanin Derived from Apricot Kernel Skin in *Drosophila*. *Food Science*. 2012,33(21): 285-289.
24. Jiao SP, Chen B, Du PG. Anti-lipid peroxidation effect of *Rosa davuriuca* Pall.fruit. *Journal of Chinese Integrative Medicine*. 2004, 2(5):364-365.
25. Zhao J, Chen C, Ma N, Sun ZO, Xue WC, Wang H. The Anti-Aging Effect of Phloridein in Male *Drosophila melanogaster*. *Acta Nutrimenta Sinica*. 2015, 37(04): 372-375+383. DOI:10.13325/j.cnki.acta.nutr.sin.2015.04.019
26. Zhou Q, Zhu L, Zhang D, Li N, Li Q, Dai P, et al. Oxidative Stress-Related Biomarkers in Postmenopausal Osteoporosis: A Systematic Review and Meta-Analyses. *Disease Markers*. 2016, 2016:7067984. DOI: 10.1155/2016/7067984
27. Lei CW, Yang ZQ, Zeng YP, Zhou Y, Huang Y, He XS, et al. Xylastriasan A, a new cytochalasan from the fungus *Xylaria striata*. *Natural product research*. 2018, 32(1): 7-13. DOI: 10.1080/14786419.2017.1324959