

# Performance of skim goat milk mineral content subjected to the block freeze concentration process

Maria Helena Machado Canella<sup>1</sup>, Eulália Lopes da Silva Barros<sup>2</sup>, Callebe Camelo-Silva<sup>2</sup>, Silvani Verruck<sup>1</sup>, Heitor Daguer<sup>3</sup>, Luciano Molognoni<sup>3</sup> and Elane Schwinden Prudencio<sup>1\*</sup>

<sup>1</sup>Department of Food Science and Technology, Federal University of Santa Catarina (UFSC), Agricultural Science Center, Florianópolis, Santa Catarina, Brazil.

<sup>2</sup>Department of Chemistry and Food Engineering, Federal University of Santa Catarina (UFSC), Technological Center, Florianópolis, Santa Catarina, Brazil.

<sup>3</sup>Ministry of Agriculture, Livestock and Food Supply, National Agricultural and Livestock Laboratory, Advanced Laboratories Section of São José, São José, Santa Catarina, Brazil

## Authors' contributions

*This work was carried out in collaboration between all authors. M H M C and E S P designed this study, reviewed all steps and the data analysis. M H M C, L M and E S P wrote the protocol of the analysis and the first draft. M H M C, E L S B and C C S realized the statistical analysis and managed the literature searches. S V and H D reviewed all steps of this work. All authors read and approved the final manuscript.*

---

## ABSTRACT

The aim of this study was to evaluate the goat milk mineral performance concentrated by block freeze concentration process. Twenty batches of skim goat milk, each one with one liter, were subjected until the third stage of the freeze concentration process. The initial skim goat milk, concentrated, and ice fractions obtained were analyzed by calcium, magnesium, zinc, phosphorus, sodium and potassium content. Results showed that with the freeze concentration stages only phosphorus content not increased ( $P < 0.05$ ), for concentrated and ice fractions. In the first stage of freeze concentration process, the magnesium element showed the higher ( $P < 0.05$ ) efficiency (95 %). However, the higher ( $P < 0.05$ ) concentration factor was determinate to calcium element in the third stage (10000 %). Also, it was observed an increase in the minerals contents evaluated with the increasing of freeze concentration stages of skim goat milk. Based on results obtained in the present study, the concentrated from the first stage showed the best performance of skim goat milk mineral content.

*Keywords: Goat milk concentrated, concentration process, main mineral elements, efficiency of process, concentration factor.*

## 1. INTRODUCTION

Goat milk and its products are important in human nutrition and have become a part of the current trend of healthy eating around the world [1, 2]. Goat milk has high added value because it is a source of nutritional compositional of excellent quality, including the quantity of minerals [3, 4]. Goat milk present some major and minority minerals in larger amounts than cow milk [5-7]. Minerals are fundamental for human health, as they are required for many physiological functions such as tissue growth, regulation of enzyme activities, blood

40 clotting, and to facility of membrane transport of essential nutrients [8, 9]. Besides their  
41 effects on health, minerals influence milk technological traits, casein micelle structure and  
42 aggregation, rennet coagulation time, curd structure, and cheese yield [8, 10-11].

43

44 It is known that most people consume foods that have less than two-thirds of one or more  
45 essential minerals [12]. In addition, because of that, the production of mineral-supplemented  
46 foods **is growing** as an important strategy to prevent mineral deficiencies. Milk and milk  
47 based products are good **materials** for mineral fortification due to their worldwide  
48 consumption by all groups at risk of deficiency, and also because of their high nutritional  
49 value, of the buffering effect on digestion and absorption processes, and of the positive  
50 effects on growth [9]. The concentration of milk may be an alternative to supplementation of  
51 these products. New methods are developed to increase goat milk and its derivate quality.  
52 Also, the development of new added value products has led to increased interest in specific  
53 studies focused on the suitable ways of improve goat milk nutrition, quality, and  
54 consumption.

55

56 The block freeze concentration technology makes it possible to produce food **concentrated**  
57 with high quality by recovering a food solute based on the separation of pure ice crystals  
58 from a freeze-concentrated aqueous phase. When compared with traditional concentration  
59 processes, such as evaporation, freeze concentration not only shows some significant  
60 potential advantages for the production of a concentrated where no vapor/liquid interface  
61 exists but also can protect thermally fragile food compounds [13]. According to Sánchez et  
62 al. [14], the freeze concentration reduces **about** three times the total cost of the process  
63 (including capital, cleaning and energy), when compared to the evaporation or reverse  
64 osmosis processes.

65

66 The freeze concentration has highly promising applications, especially, in the production of  
67 foods and ingredients that have high nutritive value [15]. In this technology, a food liquid  
68 solution is completely frozen and then, the whole frozen solution is thawed, with separation  
69 of concentrated fraction from ice fraction by gravitational thawing. Sometimes the separation  
70 may be carried out assisted by other techniques to enhance separation efficiency [16, 17].  
71 The concentration of solutes retained in the ice formed determines the efficiency of this  
72 process [15]. This technique has been used in concentration of different foods, such as  
73 cheese whey [14, 18], milk [19], skim milk [20], wine [17], fruit juices [13, 21, 22], coffee  
74 extract [23], and tofu whey [24].

75

76 Studies **have been** conducted on the properties of concentrated skim goat milk prepared by  
77 ultrafiltration [11]. However, in the light of our knowledge, there are no reports in the  
78 literature on how mineral performance of skim goat milk is affected by the block freeze  
79 concentration technology. A better understanding of this behavior is necessary to further  
80 understand the use of freeze concentrated milk in production and processing of new dairy  
81 products. Therefore, the aim of the present study was to **concentrate** skim goat milk by block  
82 freeze concentration process and to evaluate the impact of the process on mineral  
83 performance of the concentrated and the ice fractions.

84

85

## 86 **2. MATERIALS AND METHODS**

87

### 88 **2.1 Materials**

89

90 Commercial skim UHT goat milk (Capilat®, CCA Laticínios, Rio de Janeiro, Brazil) was  
91 used as the start material. The skim goat milk composition was  $8.46 \pm 0.01$  g total solids 100

92  $\text{g}^{-1}$ ,  $2.91 \pm 0.05 \text{ g total protein } 100 \text{ g}^{-1}$ ,  $3.93 \pm 0.05 \text{ g lactose } 100 \text{ g}^{-1}$  and  $0.89 \pm 0.03 \text{ g ash}$   
93  $100 \text{ g}^{-1}$ . All reagents were of analytical grade.


94

## 95 2.2. Protocol of the skim goat milk freeze concentration procedure

96

97 The freeze concentration procedure used to concentrate the skim goat milk was carried out  
98 by applying the block freeze concentration technique, according to the process proposed by  
99 [25]. An initial volume of 20 L of skim goat milk was separated into twenty batches of 1 L.  
100 Each 1L of skim goat milk was fractionated in plastic containers and were frozen at  $-20 \pm 2$   
101  $^{\circ}\text{C}$  in a freezer unit (Consul, Biplex CRD41D, São Bernardo do Campo, Brazil). After the  
102 skim goat milk has been completely frozen, 50 % of the initial volume was defrosted at room  
103 temperature ( $20 \pm 2 \text{ }^{\circ}\text{C}$ ), obtaining two fractions, the concentrated goat milk (CG1) and the  
104 ice (I1). The defrosted liquid (CG1) was frozen at  $-20 \pm 2 \text{ }^{\circ}\text{C}$  and used as feed solution in  
105 the second stage. This procedure was repeated until the third stage (Figure 1). After each  
106 stage, a portion of concentrated (CG1, CG2, and CG3), and ice fractions (I1, I2 and I3) was  
107 collected and stored at  $-20 \pm 2 \text{ }^{\circ}\text{C}$  until the analysis.

108



109

110 Fig. 1. Diagram of the skim goat milk block freeze concentration process.

111

## 112 2.3 Mineral fractions analysis

### 113 2.3.1 Calcium, magnesium, and zinc content

114

115 The determination of mineral elements Ca, Mg, and Zn content ( $\text{mg kg}^{-1}$ ) were carried out  
 116 by flame atomic absorption spectrometry (F-AAS) according to Navarro-Alarcón et al. [26],  
 117 with modifications. The spectrometer used was the AAnalyst 200 model (PerkinElmer, Inc.,  
 118 Waltham, MA, EUA) equipped with the background corrector, and the deuterium arc  
 119 illumination, using the Echelle resolution system. Acetylene (purity 99.7 %) was employed as  
 120 fuel gas to heat the atomization system and as compressed gas was used as the  
 121 compressed air. Before the measurement, all samples were calcined at  $520\text{ }^{\circ}\text{C}$ , and the ash  
 122 obtained were treated with hydrochloric acid  $8\text{ mol L}^{-1}$ . The analytical and instrumental  
 123 parameters were adjusted to obtain the best sensitivity for each element (Table 1). For this,  
 124 the samples were diluted with Milli-Q water for interpolation in the linear range of each  
 125 mineral element. Cathode lamps (PerkinElmer, Inc., Waltham, MA, USA) were employed to  
 126 determinate minerals elements. All analyses were carried out in triplicate, and blanks were  
 127 prepared with bidistilled deionized water.

128 **Table 1.** Flame atomic absorption spectrometry (F-AAS) instrumental parameters.

Minerals	Wavelengths (nm)	Linear range ( $\text{mg kg}^{-1}$ )
Ca	422.67	1.00 - 5.00
Mg	285.21	0.10 - 0.30
Zn	213.86	0.10 - 1.50

130

### 131 **2.3.2 Phosphorus content**

132

133 Phosphorus content ( $\text{mg kg}^{-1}$ ) was measured by molecular spectrometry at 420 nm in a  
 134 spectrophotometer UV-Vis, with deuterium lamp (Thermo Fisher Scientific Inc., Waltham,  
 135 MA, USA). The samples were initially calcined ( $520\text{ }^{\circ}\text{C}$ ), and complexed with molybdenum  
 136 phosphoric acid. Samples results were interpolated in calibration curves constructed with  
 137 diacid phosphate of potassium, in the range of 1 to  $20\text{ mg L}^{-1}$ . All analyses were carried out  
 138 in triplicate, and blanks were prepared with bidistilled deionized water.

139

### 140 **2.3.3 Sodium and potassium content**

141

142 The sodium and potassium content ( $\text{mg kg}^{-1}$ ) were determined through the technique of  
 143 atomic emission spectrometry (F-AES), with a flame photometer 910M (Analyser Comércio e  
 144 Indústria Ltda., São Paulo, Brazil) at 589.0 e 710 nm, respectively. For the evaluation of  
 145 these minerals, the samples were calcined at  $520\text{ }^{\circ}\text{C}$ , and treated with nitric acid  $4\text{ mol L}^{-1}$ .  
 146 Sample results were interpolated in calibration curves constructed in the range of 1 to  $10\text{ mg}$   
 147  $\text{L}^{-1}$ . All analyses were carried out in triplicate, and blanks were prepared with bidistilled  
 148 deionized water.

149

## 150 **2.4 Freeze concentration parameters**

151

### 152 **2.4.1 Concentration factor**

153

154 The concentration factor (CF) was calculated in agreement with the method proposed by  
 155 Aider and Ounis [27]. The CF of each freeze concentration stage was determinate as a  
 156 function of the increase of mineral content, using the following Equation 1:

157

$$158 \quad CF (\%) = \frac{MC_n}{MC_0} \times 100 \quad (1)$$

159

160 where  $MC_n$  is the mineral ( $\text{mg kg}^{-1}$ ) content of the concentrated goat milk from each freeze  
 161 concentration stage and  $MC_0$  is the mineral ( $\text{mg kg}^{-1}$ ) content of the initial skim goat milk.

162

163  
164  
165  
166  
167  
168

## 2.4.2 Process efficiency

The process efficiency (*eff*) was calculated based on the increase of mineral content ( $\text{mg kg}^{-1}$ ) in the concentrated goat milk ( $\text{MC}_n$ ) in relation to the mineral content ( $\text{mg kg}^{-1}$ ) remaining in the ice ( $\text{MC}_i$ ) from each freeze concentration stage ( $n$ ), as described in the Equation 2:

$$\text{eff} (\%) = \frac{\text{MC}_n - \text{MC}_i}{\text{MC}_n} \times 100 \quad (2)$$

170  
171  
172

## 2.5 Statistical analysis

Data were expressed as means and standard deviations. Statistical analysis of data was performed using the software STATISTICA 13.3 software (TIBCO Software Inc., Palo Alto, CA). One-way analyses of variance (ANOVA) and Tukey's studentized range (5 % significance) were carried out to test significant differences between the results.

173  
174  
175  
176  
177  
178

## 3. RESULTS AND DISCUSSION

179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199

Goat milk is considered an exceptionally important food because is rich in mineral content. The mineral fractions of skim goat milk, concentrated (CG1, CG2, and CG3), and ice fractions (I1, I2, I3) are shown in Table 2. Generally, the mineral content in the concentrated and ice fraction increased with increase in freeze concentration stages. When verified the concentration of major elements such as Ca, Mg, Na, K and P, it was possible to note that the values of Ca, Mg, Na, K were higher ( $P < 0.05$ ) in all concentrated fractions (CG1, CG2, and CG3), when compared with the initial skim goat milk. Besides that, these minerals contents in CG1, CG2, and CG3 increased ( $P < 0.05$ ) with the increase of the freeze concentration stages. This performance over the freeze concentration stages was expected, because similar behavior was reported in block freeze concentration process of the skim cow milk [20]. The concentration of Ca and Mg were higher than those reported by Moreno-Montoro et al. [11] during the ultrafiltration of skimmed goat milk. Ca and Mg contents are related to casein structure, which is primarily involved in the coagulation process and curd formation, and a higher concentration of Ca to milk decreases rennet clotting time and increases curd firmness [10, 28-30]. The P content showed no difference ( $P > 0.05$ ) between the initial skim goat milk and concentrated fraction (CG1, CG2, and CG3). It was noted a slight progressive increase in relation to Ca, Mg, Na, K, and P contents for the ice fractions of freeze concentration stages. However, I1 and I2 fractions showed lower values ( $P < 0.05$ ) of these minerals when compared with the initial skim goat milk.

200  
201

**Table 2. Mineral contents performance of skim goat milk, concentrated (CG1, CG2, and CG3), and ice (I1, I2, and I3) fractions during block freeze concentration stages.**

Samples	Ca ( $\text{mg kg}^{-1}$ )	Mg ( $\text{mg kg}^{-1}$ )	Na ( $\text{mg kg}^{-1}$ )	K ( $\text{mg kg}^{-1}$ )	P ( $\text{mg kg}^{-1}$ )	Zn ( $\text{mg kg}^{-1}$ )
Skim goat milk	987.48 ± 3.38 <sup>dB</sup>	82.52 ± 1.21 <sup>dB</sup>	676.59 ± 0.32 <sup>dB</sup>	1429.89 ± 108.48 <sup>dB</sup>	476.74 ± 35.62 <sup>aB</sup>	6.97 ± 0.03 <sup>dA</sup>
CG1	1720.38 ± 3.31 <sup>c</sup>	147.36 ± 0.01 <sup>c</sup>	1004.86 ± 1.19 <sup>c</sup>	1842.38 ± 239.03 <sup>c</sup>	508.25 ± 4.55 <sup>a</sup>	10.71 ± 0.01 <sup>b</sup>
I1	122.30 ± 0.01 <sup>D</sup>	7.11 ± 0.12 <sup>D</sup>	316.00 ± 6.00 <sup>D</sup>	316.02 ± 6.50 <sup>D</sup>	242.52 ± 5.96 <sup>D</sup>	0.93 ± 0.01 <sup>D</sup>
CG2	2307.57 ± 6.66 <sup>b</sup>	235.97 ± 1.20 <sup>b</sup>	1258.26 ± 0.56 <sup>b</sup>	2831.08 ± 1.26 <sup>b</sup>	503.99 ± 0.22 <sup>a</sup>	7.94 ± 0.03 <sup>c</sup>
I2	707.34 ± 6.52 <sup>C</sup>	13.97 ± 0.01 <sup>C</sup>	608.27 ± 0.52 <sup>C</sup>	1216.55 ± 1.05 <sup>C</sup>	420.83 ± 0.11 <sup>C</sup>	2.55 ± 0.01 <sup>C</sup>
CG3	10388.28 ± 213.62 <sup>a</sup>	458.99 ± 9.44 <sup>a</sup>	1652.70 ± 33.99 <sup>a</sup>	3305.39 ± 67.97 <sup>a</sup>	522.40 ± 10.74 <sup>a</sup>	17.36 ± 0.36 <sup>a</sup>
I3	1494.75 ± 6.67 <sup>A</sup>	136.45 ± 1.20 <sup>A</sup>	880.67 ± 1.28 <sup>A</sup>	1907.97 ± 204.80 <sup>A</sup>	529.81 ± 0.17 <sup>A</sup>	5.26 ± 0.01 <sup>B</sup>

<sup>a,b,c</sup> Within a column, means ± standard deviations with different superscript lowercase letters denote significant differences ( $P < 0.05$ ) between the skim goat milk and the concentrated fraction of each freeze concentration stage.

<sup>A,B,C</sup> Within a column, means ± standard deviations with different superscript uppercase letters denote significant differences ( $P < 0.05$ ) between the skim goat milk and the ice fraction of each freeze concentration stage. CG1: concentrated fraction of first freeze concentration stage. I1: ice fraction of first freeze concentration stage. CG2: concentrated fraction of second freeze concentration stage. I2: ice fraction of second freeze concentration stage. CG3: concentrated fraction of third freeze concentration stage. I3: ice fraction of third freeze concentration stage.


202

203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225

The Zn content decreased ( $P < 0.05$ ) for the CG2 in comparison with the CG1, and with the skim goat milk. At the third stage, the Zn content increased ( $P < 0.05$ ), showing higher values for the CG3. The initial skim goat milk showed higher ( $P < 0.05$ ) Zn content than all ice fractions. According to Gao et al. [31], and Aider and Ounis [27], freezing of salt solution above its eutectic temperature causes rejection of salt (poorly soluble in ice) to the surrounding medium, creating water with very high salt content brine.

Minerals content of Ca, P, K, Na and Mg were higher than those reported by Balde and Aider [20] during the block freeze concentration of skim cow milk. This performance could be related to the fact the goat milk present some mineral contents, such as Ca, P, K and Mg, in larger amounts than cow milk [5-7].

Regarding mineral efficiency concentration (Fig. 2), overall notable values were achieved. However, the best value was obtained at concentration of Mg with an efficiency of approximately 95 % in the first stage and around 70 % at the third stage. The lowest efficiency was to P concentration with an efficiency of 52 %, 16 %, and 17 % at the first, second, and third stages, respectively. In general, the highest process efficiencies were recorded at the end of the first freeze concentration stages. These results indicate that more minerals were entrapped in the ice fraction at the final stages of freeze concentration process (I2 and I3). This performance was also stated by Aider, de Halleux, and Melnikova [32] for the freeze concentration of skim acidic milk.




226  
227  
228  
229

**Fig.2. Performance of freeze concentration process on the efficiency (eff) of goat milk minerals concentration as a function of freeze concentration stages ( □ initial skim goat milk, ■ stage 1, ▨ stage 2, and ■ stage 3).**

230  
231  
232  
233  
234  
235  
236

In the present study, for all mineral content evaluated, the concentration factor (CF) (Fig. 3) showed an opposite performance than those observed by the mineral efficiency concentration. An increase ( $P < 0.05$ ) of the concentration factor was observed over the freeze concentration stages, reaching a CF of 10000 % for the Ca content in the third freeze concentration stage.



237  
238  
239  
240  
241

**Fig.3. Performance of freeze concentration process on the concentration factor (CF) of goat milk minerals concentration as a function of freeze concentration stages (□ initial skim goat milk, ■ stage 1, ▨stage 2, and ■stage 3).**

242  
243  
244  
245  
246  
247  
248

As observed by Ceballos et al. [5], Yadav, Singh, and Yadav [7], and Campos et al. [33] in the present work it is possible to note that main elements contents of skim goat milk are higher than cow milk. Finally, in a near future, the generate results from the block freeze concentration process of skim goat milk mineral content performance can be used by dairy industries to produce nutritive products with high mineral contents without mineral supplementation, which positively affects the economic and the nutritive value of milk products.

249  
250  
251

#### 4. CONCLUSION

252  
253  
254  
255  
256  
257  
258

The mineral content of skim goat milk was successfully freeze concentrated by applying the block freeze concentration. As the freeze concentration stages increased, Ca, Mg, Na, K, and Zn contents increased in both concentrated and ice fractions. It was possible to concentrated Ca and Mg after three stages, around 10 and 6 times more than the initial skim goat milk, respectively. Indeed, the K, Na and Zn were concentrated after three stages, almost 3 times more than initial skim goat milk, respectively. However, the phosphorus showed no difference of concentrated fraction in the three stages compared with the initial

259 skim goat milk. All mineral content showed high efficiency and concentration factor during  
260 the freeze concentration process. The study concluded that the concentrated from the first  
261 stage was the best. This is because higher efficiencies results were obtained in this stage.

262

## 263 **COMPETING INTERESTS**

264

265 Authors have declared that no competing interests exist.

266

## 267 **REFERENCES**

268

269 1. Hassan FAM, Abbas HM, Abd El-Gawad MAM, Enab AK. (2014). Goats Dairy Products  
270 as a Potentially Functional Food. *Life Sci. J.* 11;648–657.

271 2. Sosnowski M, Rola JG, Osek J. (2016). Alkaline phosphatase activity and microbiological  
272 quality of heat-treated goat milk and cheeses. *Small Rumin. Res.* 136;132–136.

273 3. Olalla M, Ruiz-López M, Navarro M, Artacho R, Cabrera C, Giménez R, et al. (2009).  
274 Nitrogen fractions of Andalusian goat milk compared to similar types of commercial milk.  
275 *Food Chem.* 113;835–838.

276 4. Yangilar F. (2013). As a Potentially Functional Food: Goats' Milk and Products. *J. Food  
277 Nutr. Res.* 1;68–81.

278 5. Ceballos LS, Morales ER, de la Torre Adarve G, Castro JD, Martínez LP, Sampelayo,  
279 MRS, (2009). Composition of goat and cow milk produced under similar conditions and  
280 analyzed by identical methodology. *J. Food Compos. Anal.* 22;322–329.

281 6. Raynal-Ljutovac K, Lagriffoul G, Paccard P, Guillet, I, Chilliard Y. (2008). Composition of  
282 goat and sheep milk products: An update. *Small Rumin. Res.* 79;57–72.

283 7. Yadav AK, Singh J, Yadav SK. (2016). Composition, nutritional and therapeutic values of  
284 goat milk: A review. *Asian J. Dairy Food Res.* 35;96–102.

285 8. Stocco G, Cipolat-Gotet C, Bittante G, Cecchinato A. (2016). Variation of major mineral  
286 contents in Mediterranean buffalo milk and application of Fourier Transform Infrared  
287 spectroscopy for their indirect prediction. *J Dairy Sci.* 99(11);8680-8686.

288 9. Lombardi J, Pellegrino J M, Soazo M, Corrêa A P F, Brandelli A, Risso P. et al. (2018).  
289 Mineral fortification modifies physical and microstructural characteristics of milk gels  
290 coagulated by a bacterial enzymatic pool. *Colloids Surf., B.* 161;296-301.

291 10. Franzoi M, Niero G, Penasa M, Cassandro M, De Marchi M. (2018). Development and  
292 validation of a new method for the quantification of soluble and micellar calcium,  
293 magnesium, and potassium in milk. *J Dairy Sci.* 101(3);1883-1888.

294 11. Moreno-Montoro M, Olalla M, Giménez-Martínez R, Bergillos-Meca T, Ruiz-López MD,  
295 Cabrera-Vique C. et al. (2015). Ultrafiltration of skimmed goat milk increases its nutritional  
296 value by concentrating nonfat solids such as proteins, Ca, P, Mg, and Zn. *J. Dairy Sci.*  
297 98;7628–7634.



- 298 12. Achanta K, Aryana KJ, Boeneke CA. (2007). Fat free plain set yogurts fortified with  
299 various minerals. *LWT-Food Sci Technol.* 40(3);424-429.
- 300 13. Petzold G., Moreno,J., Lastra P., Rojas K., Orellana P. (2015). Block freeze  
301 concentration assisted by centrifugation applied to blueberry and pineapple juices. *Innov.*  
302 *Food Sci. Emerg. Technol.* 30;192–197.
- 303 14. Sánchez J., Hernández E., Auleda J.M., Raventós M., 2011. Freeze concentration of  
304 whey in a falling-film based pilot plant: Process and characterization. *J. Food Eng.* 103;147–  
305 155.
- 306 15. Aider M, de Halleux D. (2009). Cryoconcentration technology in the bio-food industry:  
307 Principles and applications. *LWT - Food Sci. Technol.* 42;679–685.
- 308 16. Orellana-Palma P, Petzold G, Pierre L, Pensaben JM. (2017). Protection of polyphenols  
309 in blueberry juice by vacuum-assisted block freeze concentration. *Food Chem. Toxicol.*  
310 109;1093–1102.
- 311 17. Petzold G, Orellana P, Moreno J, Cerda E, Parra P. (2016). Vacuum-assisted block  
312 freeze concentration applied to wine. *Innov. Food Sci. Emerg. Technol.* 36;330–335.
- 313 18. Canella, MHM, Muñoz IB, Pinto SS, De Liz GR, Müller CMO, Amboni RDMC. et al.  
314 (2018). Use of Concentrated Whey by Freeze Concentration Process to Obtain a Symbiotic  
315 Fermented Lactic Beverage. *Adv. J. Food Sci. Technol.* 14;56–68.
- 316 19. Muñoz IB, Canella, MHM, Verruck S, Müller CMO, De Liz GR, Amboni RDMC. et al.  
317 (2017). Potential of Milk Freeze Concentration for the Production of Functional Fresh  
318 Cheeses. *Adv. J. Food Sci. Technol.* 13;196–209.
- 319 20. Balde A, Aider M. (2016). Impact of cryoconcentration on casein micelle size distribution,  
320 micelles inter-distance, and flow behavior of skim milk during refrigerated storage. *Innov.*  
321 *Food Sci. Emerg. Technol.* 34;68–76.
- 322 21. Hernández E, Raventós M, Auleda JM, Ibarz A. (2009). Concentration of apple and pear  
323 juices in a multi-plate freeze concentrator. *Innov. Food Sci. Emerg. Technol.* 10;348–355.
- 324 22. Miyawaki O, Omote C, Gunathilake M, Ishisaki K, Miwa S, Tagami A. et al. (2016).  
325 Integrated system of progressive freeze-concentration combined with partial ice-melting for  
326 yield improvement. *J. Food Eng.* 184;38–43.
- 327 23. Moreno FL, Quintanilla-Carvajal MX, Sotelo LI, Osorio C, Raventós M, Hernández E. et  
328 al. (2015). Volatile compounds, sensory quality and ice morphology in falling-film and block  
329 freeze concentration of coffee extract. *J. Food Eng.* 166;64–71.
- 330 24. Belén F, Benedetti S, Sánchez J, Hernández E, Auleda JM, Prudêncio ES. et al. (2013).  
331 Behavior of functional compounds during freeze concentration of tofu whey. *J. Food Eng.*  
332 116;681–688.
- 333 25. Aider M, de Halleux D, Akbache A. (2007). Whey cryoconcentration and impact on its  
334 composition. *J. Food Eng.* 82;92–102.
- 335 26. Navarro-Alarcón M, Cabrera-Vique C, Ruiz-López MD, Olalla M, Artacho R, Giménez, R.  
336 (2011). Levels of Se, Zn, Mg and Ca in commercial goat and cow milk fermented products:

- 337 Relationship with their chemical composition and probiotic starter culture. Food Chem.  
338 129;1126–1131.
- 339 27. Aider M, Ounis WB. (2012). Skim milk cryoconcentration as affected by the thawing  
340 mode: Gravitational vs. microwave-assisted. Int. J. Food Sci. Technol. 47;195–202.
- 341 28. Landfeld A, Novotná P, Houška M. (2002). Influence of the Amount of Rennet, Calcium  
342 Chloride Addition, Temperature, and High-Pressure Treatment on the Course of Milk  
343 Coagulation. Czech J. Food Sci. 20(6);237–244.
- 344 29. Malacarne M, Franceschi P, Formaggioni P, Sandri S, Mariani P, Summer A. (2014).  
345 Influence of micellar calcium and phosphorus on rennet coagulation properties of cows milk.  
346 J. Dairy Res. 81;129–136.
- 347 30. Visentin G, Penasa M, Gottardo P, Cassandro M, De Marchi M. (2016). Predictive  
348 ability of mid-infrared spectroscopy for major mineral composition and coagulation traits of  
349 bovine milk by using the uninformative variable selection algorithm. J. Dairy  
350 Sci. 99(10);8137-8145.
- 351 31. Gao W, Smith DW, Segó DC. (2004). Treatment of pulp mill and oil sands industrial  
352 wastewaters by the partial spray freezing process. Water Res. 38;579–584.
- 353 32. Aider M, de Halleux D, Melnikova I. (2009). Skim acidic milk whey cryoconcentration and  
354 assessment of its functional properties: Impact of processing conditions. Innov. Food Sci.  
355 Emerg. Technol. 10;334–341.
- 356 33. Campos MS, López-Aliaga I, Alférez MJM, Nestares T, Barrionuevo M. (2003). Effects of  
357 goats' or cows' milks on nutritive utilization of calcium and phosphorus in rats with intestinal  
358 resection. Br. J. Nutr. 90;61-67.