

Effect of zinc and boron application on yield of **brinjal** (*Solanum melongena* L.) through On Farm Trial in Bharuch District of Gujarat

D. J. Modi*, L. M. Patil, H. M. Vasava and M. M. Patel

BAIF Krishi Vigyan Kendra- Bharuch -393 130, Gujarat

ABSTRACT

The field experiment was carried out as on farm trial in the year 2016-17 during Rabi season to know effect of zinc and boron on yield of **brinjal**. Total 10 farmers were selected from three tribal blocks namely Netrang, Zaghadia and Valia of Bharuch District. Farmers are not getting good fruit yield of brinjal. The production of brinjal crop continues to be quite low. The objectives were to find out the nutritional requirement of the crop and for making fertilizer recommendation based on soil test for **brinjal** crop. Soil samples were collected from the selected field. The collected samples were analyzed to evaluate the fertility condition of field. After analysis of soil result is deficient of Zinc and Boron. The fruit yield of brinjal crop can be increased by the application of Zinc and Boron in soil as well as foliar spray. The plant height, average fruit weight, number of fruit per plant, yield per hacter, gross returns, net returns and benefit cost ratio recorded was highest in soil application of zinc and boron as compared **farmer's** practices.

Key Words: **On farm trial (OFT)**, Zinc, Boron, Brinjal, Yield and Economics

1. INTRODUCTION

Brinjal (*Solanum melongena* L.), or egg plant is one of the most common, popular and major vegetable crop grown in India and other parts of the world. The brinjal is of much important in the warm areas of Far East, being grown extensively in India and other Asian countries like Bangladesh, Pakistan, and Philippines. Other major brinjal producing countries are China, Turkey, Japan, Egypt, Indonesia, Iraq, Italy, Syria and Spain. The cultivated brinjal is of Indian origin and has been in cultivation from long time (Thompson and Kelly, 1957). Brinjal (*Solanum melongena* L.) belongs to the family Solanaceae.

Micronutrients like, **copper** and zinc also play a positive role for increasing fruit as well as seed yield in brinjal. Micronutrients are those elements that are essential for plant growth, but which was required in smaller amount. If these elements are not available sufficiently, plants will suffer from physiological stresses caused by inefficiency of several enzymatic systems and other related metabolic functions. Various responses were observed in growth and yield in crops species and in cultivars to trace elements deficiency (Fageria, 2009).

But brinjal grower farmers do not have the suitable techniques to increase the fruit yield of brinjal. Due to lack of awareness regarding soil trial and balanced fertilizer application farmers are facing low crop yield. By conducting farmer's field visit, it was observed that one of the important factors for low productivity of brinjal was not applying micronutrients, wilting problem and infestation of shoot & fruit borer. It was with this objective in view, the present study was initiated.

Corresponding author Email ID: modihort1983@yahoo.com

2. MATERIAL AND METHODOLGY

The field experiment was conducted at the ten farmers field under supervision of Krishi Vigyan Kendra, Bharuch, Gujarat during (2016-17) Rabi season, in five villages of three blocks Netrang, Zaghadia and Valia. Ten innovative and receptive progressive farmers from five villages were selected for conducting the trial to ensure their active participation. Investigation on soil test crop response correlation for zinc and boron in order to evolve soil test based fertilizer recommendation for Brinjal crop. These farmers were selected for the On Farm Trial of zinc and boron soil application and foliar spray on their area and production. The demonstration of improved technology was taken in area of 0.4 ha of each farmer. The trial plot of farmer was equally divided for three treatments. The soil and foliar application of zinc and boron along with recommended doses of fertilizer (100:50:50 kg NPK /ha) was compared against farmer practices. The soil application of zinc (10 kg per acre) and boron (4 kg per acre) was done along with basal dose, however the foliar spray of 0.5% zinc and 0.2% boron was made at the flowering and fruiting stage. Nitrogen was applied in at basal and at flowering stage. At the time of experiment parameters observed were like plant height, average fruit weight (gm), number of fruits per plant and total yield of each experimental plot as well as farmer's practices.

The primary data on output of brinjal yield were collected from the selected brinjal farmers plots, qualitative data was converted in to quantitative form and expressed in term of percent increased yield was calculated by the using formula. The soil of the plot was sandy medium black in texture having good fertility properly leveled and well drained. Analysis of soil sampled from the experimental land of farmer's field showed pH, electrical conductivity (dS/m), organic carbon (%), available Nitrogen (kg/ha), available phosphorus (kg/ha) and available potash (kg/ha), available zinc and boron (Table 1). The yield data were collected from demonstrated and farmer's practices.

$$\text{Percentage increased yield} = \frac{\text{Demonstration yield} - \text{Local check yield}}{\text{Local Check yield}} \times 100$$

Benefit cost ratio calculated by the formula given below

$$\text{Benefit cost ratio} = \frac{\text{Gross Return}}{\text{Cost of cultivation}} \times 100$$

Table 1: Soil test results of plots before conducting on farm trial.

Treatment	pH	Electrical conductivity (dS/m)	Organic Carbon (%)	Available Nitrogen (kg/ha)	Available phosphorus (kg/ha)	Available Potash (kg/ha)	Available Zinc (ppm)	Available Boron (ppm)
Field -1	6.54	1.05	0.75	270.51	38.90	160.80	0.10	0.38
Field -2	6.81	1.10	0.75	224.67	37.40	155.10	0.49	0.53
Field -3	6.75	1.01	0.44	320.52	41.19	282.50	0.55	0.39
Field -4	6.87	1.03	0.71	275.31	35.56	145.23	0.48	0.43
Field -5	7.30	1.07	0.64	186.90	83.48	144.34	0.45	0.32
Field -6	7.21	1.01	1.01	235.21	32.13	87.09	0.41	0.45
Field -7	6.93	1.05	0.83	295.75	134.69	180.60	0.44	0.51
Field -8	6.68	1.00	0.62	203.21	38.96	86.41	0.52	0.40
Field -9	7.12	1.04	0.68	370.36	75.48	141.30	0.35	0.48

Field -10	7.24	1.03	0.83	219.40	69.20	246.70	0.41	0.37
-----------	------	------	------	--------	-------	--------	------	------

3. RESULT AND DISCUSSION

The differences between on farm trial and farmer's practice are shown in table 2. The farmers practices include are no seed treatment, imbalance fertilizer application, no use of micronutrient and no adoption of plant protection measures. All these collectively resulted in low yield of brinjal in the district. However, in on farm trial plots, all the improved package of practices of brinjal was demonstrated.

Table 2: Details of brinjal growing under on farm trial and existing farmer's practices.





Operation	Existing practices	Improved Practices of Demonstrated
Variety Use	Surti Gulabi	Surti Gulabi
Seed treatment	No seed treatment	Seed treatment with Biofertilizer (20ml/kg seed) & <i>Trichoderma viride</i> 3gm/kg seed.
Sapling root treatment	No sapling root treatment	Sapling root treatment with Imidacloprid 5ml/10 liter water for 30 minutes
Spacing	80 cm x 60 cm	90 cm X 90 cm
Fertilizer application & Soil trial	Imbalance application of fertilizer & no soil test	Application of recommendation dose of fertilizer: FYM 10 t/ha and 100:50:50 kg NPK /ha and application of zinc and boron.
Plant protection measure	Non adoption of IPM practices	Spraying of Imidacloprid 5ml/10 liter water for sucking pest & Carbendazim 20 gm/10 liter water for fungus diseases.

3.1 PERFORMANCE OF ON FARM TRIAL

Soil application along with recommended doses of fertilizer significant effect on the yield, plant height, average fruit weight and number of fruits per plant attributes of brinjal in both the years (Table 3). Soil application of **zinc & boron** in addition to recommended doses of fertilizer recorded significantly higher plant height (52.39 cm), average fruit weight (167.49 g), highest number of fruits per plant (11.02) and maximum yield per hectare (33.73 tonne) compared to farmer's practices (fig. 1-4). This was followed by treatment recommended doses of fertilizer along with foliar spray application of zinc and boron. The plant height (51.17 cm), average fruit weight (164.8 g), highest number of fruits per plant (9.52) and yield per hectare (29.14 tonne) was recorded in recommended doses of fertilizer along with foliar spray application of zinc and boron. These results were in conformity with those of Pandav *et al.* (2016), Sliman *et al.* (1999) and Acharya *et al.* (2015).

Table 3: Effect of different treatments of soil application & foliar sprays of micronutrients on growth and yield parameters of brinjal

Treatment	Plant Height (cm)	Average fruit weight (gm)	No. of fruit per plant	Yield per ha (tonne)
Farmer's Practices	49.21	162.37	7.84	23.81
Recommended doses of fertilizer (RDF) + Foliar Spray	51.17	164.81	9.52	29.14
Recommended doses of fertilizer (RDF) + Soil Application	52.39	167.49	11.02	33.73
C.D.5 %	1.73	3.95	0.59	2.40
CV	2.91	2.06	5.39	7.15



3.2 ECONOMIC RETURN

The inputs and outputs prices of commodities prevailed during the study of demonstration were taken for calculating net return and benefit: cost ratio (Table 4). The cost of cultivation is comparatively more in recommended dose of fertilizers along with soil application of zinc and boron (Rs 49320 per ha) as well as in foliar application of zinc and boron (Rs 48500 per ha) as compared to farmer's practices (Rs 41600 per ha). The cultivation of brinjal under recommended doses of fertilizer along with soil application of zinc and boron gave higher net return of Rs. 119180/- per hectare followed by recommended doses of fertilizer along with foliar spray of zinc and boron Rs. 97200/- per ha as compared to farmer's practices. The benefit cost ratio of brinjal cultivation under soil application of zinc and boron practices were 3.41 against 2.86 under farmer's practice. This may be due to higher yield obtained under improved technologies compared to farmer's practice.

Table 4: Economic Impact of soil application & foliar sprays of micronutrients of brinjal

Economics of demonstration (Rs/ha)								Economics of farmer practices (Rs/ha)			
Recommended doses of fertilizer + Soil Application				Recommended doses of fertilizer + Foliar Spray							
Gross cost	Gross Income (Rs./ha)	Net Return (Rs./ha)	BCR	Gross cost	Gross Income (Rs./ha)	Net Return (Rs./ha)	BCR	Gross cost	Gross Income (Rs./ha)	Net Return (Rs./ha)	BCR
49320	168500	119180	3.41	48500	145700	97200	3.0	41600	119050	77450	2.86

4. CONCLUSION

The soil application of zinc and boron in addition to recommended doses of NPK can be effective practice to deal with low productivity of the brinjal due to zinc and boron deficiency. Farmers were very much satisfied with average fruit weight and no of fruits per plants & yield and higher benefit cost ratio that was better in soil application as compare to routine practices. However horizontal spread of recommended and improved technologies may be achieved by the successful implementation of results of on farm trials (OFT) and various extensions activities like training programmes, Kisan gosthi, and farmers meeting etc.

REFERENCES

1. Acharya U., Sharma N., Rimal K., Venkatesan T., Saraswathi K. and Subramanian S.. Response on growth, yield and quality parameters of Multiplier Onion (*Allium cepa* L. var. aggregatum Don.) var. CO (On) 5 with different doses and method of zinc application. Int J Agron and Agril Res 2015; 6(4): 59-67.
2. Dube B.K., Pratima S., Chatterjee C. and Sinha P. Effect of zinc on the yield and quality of tomato. Indian J. Hort., 2003; 60(1): 59-63.
3. Fageria N. K. The use of nutrients in crop plants. Boca Raton, 2009 Fl.: CRC (*In Press*).
4. Kiran J. Effect of fertilizer, bio-fertilizer and micronutrients on seed yield and quality of brinjal (*Solanum melongena* L.). Thesis submitted to Department of seed science and technology, College of Agriculture, Dharwad, (India) 2006 pp. 1- 2.
5. Pandav A. K., Nalla M. K., Aslam, T., Rana M. K. and Bommesh, J. C. Effect of foliar application of micronutrients on growth and yield parameters in eggplant cv. HLB 12. Environment & Ecology. 2016; 35 (3): 1745— 1748.
6. Sharma S. P. and Brar J. S. Nutritional requirements of brinjal (*Solanum melongena* L.). Punjab Agricultural University, Bathinda-151 001, India, Agric. Rev. 2008; 29(2): 79-88.
7. Sliman Z. T, Abdelhakim M. A. and Omran A. A. Response of onion to foliar application of some micronutrients. Egyptian J of Agr Res 1999; 77(3): 983-993.
8. Suganiya, S. and Kumuthini H. D. Effect of boron on flower and fruit set and yield of raton brinjal crop. International Journal of Scientific Research and Innovative Technology. 2015; 2 (1):135-141.
9. Thompson B. C. and Kelly W. C. Vegetable crops McGraw - Hill, New York, 1957; 611pp.