

2 **Growing media quality and plug cell volume**  
3 **would be interactive abiotic stresses for**  
4 ***Impatiens walleriana* pot yield**

5  
6  
7  
8 **ABSTRACT**

9 Higher bedding plant yields per unit greenhouse area was reaching through two grower's  
10 currently decision-making: plug cell volume during nursery and growing media quality for  
11 both nursery and pot cycle. With the goal of maximizing bedding plant yield to identify the  
12 main limiting factor at the pot stage, we evaluated *I. walleriana* yield to the end of the pot  
13 growth stage when four different pre-transplant cell volume and four pre or post-transplant  
14 growing media with different physical properties were used. The hypothesis tested was that  
only one of the potentially negative stress source (pre-transplant cell volume or growing  
medium quality) is the main responsible for decreasing biomass accumulation at the post-  
transplant pot growing cycle. The experimental design was a randomised factorial with three  
blocks of five single-pot replications of each treatment combination (plug cell volume ×  
growing medium × pre- and post-transplant). The main result was that, in *I. walleriana*  
seedlings, the combining abiotic stresses imposed by both the growing medium quality and  
nursery plug cell volume defined biomass accumulation (on a fresh and dry base), leaf area  
expanded and photo assimilates partitioned as opposed to a previous report, which indicate  
that that growing media quality would be a more limited factor than plug cell volume for *I.*  
*walleriana* seedlings during nursery.

15 **1. INTRODUCTION**

16  
17 Bedding plant industry has been exponentially expanded around the world according to  
18 significant costs decrease. The last has been related to higher bedding pot plant yield per unit  
19 greenhouse area through two grower currently decision-making: plug cell volume during nursery  
20 and growing media quality for both nursery and pot cycle. Commercial profits have been related  
21 to a decrease in plug cell volume [1] and the use of lower expensive growing media [2,3,4].  
22 However, these business choices imply that plants will suffer different root restriction stresses  
23 during most growing cycle.

24  
25 Usually, the 'root restriction syndrome' has been defined as a physical stress imposed on a root  
26 system when plants are grown in small containers, which leads to a pronounced decrease in  
27 both root and shoot growth at the transplant stage. The pre- and post-transplant effects of the  
28 container volume during nursery [5,6] have been extensively studied by our laboratory and we  
29 found that growth restrictions would be closely related to endogenous cytokinin synthesis by  
30 roots. A limited plug cell volume gives a vertical root restriction when root apical meristem  
31 comes to the bottom of the cell or the pot. At this moment, both primary root growth and root  
32 branching decrease [7] and cytokinins, the main endogenous hormone synthesized by root  
33 apical meristems would decrease as well [8,9].  
34

35 On the other hand, when a growing media quality decreases, it changes their physical  
36 properties [3,10], which generally results in decreased pore sizes, and would be taken into  
37 account as an abiotic stress [4]. As pore size decreases, total porosity and air-filled porosity  
38 decrease as well. Bailey-Serres and Colmer [11] and Voesenek and Sasidharan [12] have  
39 indicated that a lack of oxygen inhibits respiration, decrease metabolic plant adaptations to cope  
40 with the hypoxic and anoxic conditions and resulting energy deficits, as well as change  
41 anatomical and morphological adaptations to improve internal O<sub>2</sub> supply. These metabolic  
42 changes would give the same effects on endogenous cytokinin synthesis as plug cell volume  
43 [13,14].

44  
45 With the goal of maximizing bedding plant yield to identify the main limiting factor it is  
46 imperative. Previous reports on ornamentals [15,16,17] and vegetables [18,19,20,21] have  
47 shown that nursery growth has a significant effect on post-transplant biomass accumulation.  
48 The precise effects of combined plug cell volume and growing media quality on nursery have  
49 been recently indicated as well [22,23,24]. However, the simultaneously post-transplant  
50 interactions between these two stresses source during both nursery and pot growth is lacking.

51  
52 *Impatiens walleriana* (Hook.f.) is a commercially important year-round garden crop for  
53 landscape, and the first best-selling bedding plants in both developed and undeveloped  
54 countries. Most *Impatiens* genotype produce a compact green foliage and covers itself with  
55 extremely uniform growth habit and bright blooms. *Impatiens* F<sub>1</sub> genotypes prefer partial  
56 sun/shade (8-25 mol photons m<sup>-2</sup> day<sup>-1</sup>) [25]. Dry mass and flowering increase from 14 to 28°C  
57 [26]. Plants only grow well with 100% evapotranspiration [27]. *I walleriana* has been included  
58 in most research from our laboratory for the last decade.

59  
60 The aim of this work was to evaluate *I. walleriana* yield to the end of the pot growth stage when  
61 four different pre-transplant cell volume and four pre or post-transplant growing media with  
62 different physical properties were used. The hypothesis tested was that only one of the  
63 potentially negative stress source (pre-transplant cell volume or growing medium quality) is the  
64 main responsible for decreasing biomass accumulation at the post-transplant pot growing cycle.

## 65 66 67 **2. MATERIALS AND METHODOLOGY**

### 68 **2.1 Plant Material**

69  
70 Experiment were carried out under a greenhouse at the Faculty of Agronomy, University of  
71 Buenos Aires, Argentina (34° 35' 59"S, 58° 22' 23"W) from October 10<sup>th</sup> 2012 to December 9<sup>th</sup>  
72 2013 and repeated from October 16<sup>th</sup> 2013 to December 15<sup>th</sup> 2014.

73  
74 *I. walleriana* 'Xtreme White' seeds (Goldsmith Inc., NY, USA) were germinated and grown in  
75 50-, 128-, 288- and 512-cell plug tray<sup>-1</sup> (55.70, 17.37, 6.18 and 2.50 cm<sup>3</sup> cell<sup>-1</sup> respectively) in  
76 four different pre-transplant growing media as follows:

- 77  
78 1) Klasmann 411® medium (Klasmann-Deilmann, GmbH, Germany): Canadian  
79 *Sphagnum* peat moss-perlite-vermiculite (70/20/10 v/v/v) (**K**)
- 80 2) *Sphagnum maguellanicum*-perlite (80/20 v/v) (**S**)
- 81 3) River waste-perlite (80-20 v/v) (**R**)
- 82 4) *Sphagnum maguellanicum*-river waste-perlite (40-40-20, v/v/v) (**SR**).
- 83

84 When seedlings reached the transplant stage, they were transplanted into 1,200 cm<sup>3</sup> pots filled  
85 with a post-transplant Klasmann 411® medium (Klasmann-Deilmann, GmbH, Germany). At the  
86 same time, plants grown at the pre-transplant stage in a Klasmann 411® medium (Klasmann-  
87 Deilmann, GmbH, Germany) were transplanted to 1,200 cm<sup>3</sup> pots filled with the four different pre-  
88 transplant growing media tested, given 32 combinations of plug cell volume-growing media.

### 89 90 91 **2.2 Cultivation and Meteorological Data**

92

Plants were irrigated daily at saturation with high quality tap water using intermittent overhead mist. Growing media was weekly fertilized with 1.0: 0.5: 1.0: 0.5 (v/v/v/v) N: P: K: Ca through the overhead irrigation water (Stage 2: 50 mg L<sup>-1</sup> N; Stage 3-4: 100 mg L<sup>-1</sup> N; pot: 150 mg L<sup>-1</sup> N).

Daily mean temperatures (22.26 to 25.06°C) and daily photosynthetic active radiation (4.24 to 5.03 mol photons m<sup>-2</sup> day<sup>-1</sup>) for the experiments were recorded with a HOBO sensor (H08-004-02) (Onset Computer Corporation, MA, USA) connected to a HOBO H8 data logger. The plants were arranged at a density of 25 plants m<sup>-2</sup>, which avoided mutual shading.

### 2.3 Sampling and Growth Evaluations

Samples of each substrate were collected at the beginning of the pot experiments (before transplant to 1,200-cm<sup>3</sup> pots) and total porosity, air-filled porosity, bulk density and container capacity were determined according to Fonteno [28]. Data are indicated in Table 1 and show significant physical properties differences in of the growing media tested.

**Table 1: Physical properties for the growing media tested. K: [Canadian *Sphagnum* peat (70%) + Perlite (20%) + Vermiculite (10%)], S: [*Sphagnum maguellanicum* (80%) + Perlite (20%)], R: [River waste (80%) + Perlite (20%)], SR: [*Sphagnum maguellanicum* (40%) + River waste (40%) + perlite (20%)]. Standard errors are indicated.**

Growing media	Total porosity (%)	Air-filled porosity (%)	Bulk density (g dm <sup>-3</sup> )	Container capacity (%)
F	60.00 ± 0.55	12.93 ± 0.98	0.21 ± 0.04	36.89 ± 1.46
S	70.67 ± 0.67	29.67 ± 2.15	0.15 ± 0.01	48.00 ± 0.38
R	72.67 ± 0.18	44.60 ± 0.95	0.18 ± 0.01	50.22 ± 0.44
SR	67.53 ± 0.64	23.27 ± 2.43	0.21 ± 0.01	42.67 ± 0.38

Plants were harvested at the transplant stage and at 15, 30, 45, and 60 days after transplanting. Roots were washed and root, stem and leaf fresh weights (FW) were recorded. Dry weights (DW) were obtained after drying roots, stems and leaves to constant weight at 80°C for 96 hours. The number of leaves was recorded, and each leaf area was determined using the ImageJ® (Image Processing and Analysis in Java) software. The number of stems and nodes has been recorded as well.

The relative rate of leaf area expansion (RLAE) was calculated as the slope of the regression of the natural logarithm (ln) of total leaf area versus time (in days). The rate of leaf appearance (RLA) was calculated as the slope of the number of fully expanded leaves versus time (in weeks). The specific leaf area on a FW basis (SLA) and leaf weight rate (LWR) were calculated as the ratio between the area of the new individual leaf and leaf FW and the ratio between the leaf DW and the total plant DW respectively at the end of the experiments. The relative growth rate (RGR) was calculated as the slope of the regression of the ln of whole plant DW versus time (in days).

The mean net assimilation rate (NAR) and leaf area ratio (LAR) were calculated according to Potter and Jones [29] as follows:

$$NAR = \frac{k_w W_0 e^{k_w t}}{A_0 e^{k_a t}}$$

$$LAR = \frac{k_w}{NAR}$$

where W<sub>0</sub>: extrapolated value of total DW (g) at time zero; k<sub>w</sub>: RGR (day<sup>-1</sup>); A<sub>0</sub>: extrapolated value of leaf area (cm<sup>2</sup>) at time zero; k<sub>a</sub>: RLAE (day<sup>-1</sup>); t: time (days) at the midpoint of the experimental period and e: base of the ln.

143 The allometric coefficients between root and shoot were calculated as the slope ( $\beta$ ) of the  
144 straight-line regression of the ln of the root DW versus the ln of the shoot DW. The Root: Shoot  
145 ratio (at the end of the experiment) was performed as well.

146  
147

## 148 2.4 Statistical Analysis

149

150 The experimental design was a randomised factorial with three blocks of five single-pot  
151 replications of each treatment combination (plug cell volume  $\times$  growing medium  $\times$  pre- and post-  
152 transplant). Since there were no significant differences between the two experiments, they were  
153 considered together ( $n = 30$ ). Data were subjected to three-way analysis of variance (ANOVA).  
154 STATISTICA 8 (StatSoft) software was used and the assumptions of the ANOVA were checked.  
155 Means were separated by Tukey's tests ( $P \leq 0.05$ ). Slopes from straight-line regressions of  
156 RLA, RLAE, RGR and allometric values were tested using the SMATR package [30].


157  
158

## 159 3. RESULTS


### 160 3.1. Biomass accumulation

161

162 Total fresh weight at the end of the pot growth cycle (60 days from transplant) was higher in  
163 plants from 50-plug tray<sup>-1</sup> and decreased according cell numbers increased in all growing media  
164 tested. Anyway, growing media significantly changed post- transplant biomass accumulation on  
165 a FW base as during the pre- as the post-transplant stage, but especially during nursery. The  
166 higher FW was found in plants grown in R- and S-growing media (Fig. 1A). When the mean  
167 aerial FW was plotted against the mean root FW (Fig. 1B), a positive correlation was found ( $r^2 =$   
168 0.661;  $P < 0.001$ ).



169




170  
 171 **Figure 1. Total fresh weight at the end of the experiment (60 days from transplanting) for**  
 172 ***Impatiens walleriana* plants grown in four plug cell volumes (50-, 128-, 288-, and 512-cell**  
 173 **tray<sup>-1</sup>) at the pre-transplant stage and four growing media at the pre- or post-transplant**  
 174 **stage. Bars indicate standard errors and vertical line indicate least significant differences**  
 175 **(LSD). Panel B. Relationships between shoots and roots FW according to four plug cell**  
 176 **volumes (50-, 128-, 288-, and 512-cell tray<sup>-1</sup>) at the pre-transplant stage and four growing**  
 177 **media at the pre- (full symbols) or post-transplant stage (empty symbols). For substrate**  
 178 **abbreviations see Table 1. F: ♦-◇; R: ■-□; S: ●-○; SR: ▲-△. The straight-line regression**  
 179 **was: Shoot FW = 6.65 Root FW + 6.73 ( $r^2 = 0.661$ ). The probability of the slope being zero**  
 180 **was  $P < 0.001$ .**

181  
 182


### 183 3.2. Leaf area expansion

184

185 Total leaf area at the end of the experiment was once again higher in plants from 50-plug cells  
 186 and in those grown at different growing media during the pot grow stages (Fig. 2A). Although  
 187 there were significant differences in individual leaf area according to different pre-transplant cell  
 188 volume and pre- or post-transplant growing media, changes were smaller than in total leaf area  
 189 (Fig. 2B).



190



191  
192 **Figure 2. Total (A) and individual (B) leaf area at the end of the experiments (60 days from**  
193 **transplanting) for *Impatiens walleriana* plants from four plug cell volumes (50-, 128-, 288-,**  
194 **and 512-cell tray<sup>-1</sup>) at the pre-transplant stage and four growing media at the pre- or post-**  
195 **transplant stage. Bars indicate standard errors and vertical line indicate least significant**  
196 **differences (LSD). For substrate abbreviations see Table 1.**

197  
198 **Table 2: Changes in RLA, RLAE and SLA for *Impatiens walleriana* plants from four plug**  
199 **cell volumes (50-, 128-, 288-, and 512-cell tray<sup>-1</sup>) at the pre-transplant stage and four**  
200 **growing media at the pre- or post-transplant stage. Different lower-case letters indicate**  
201 **significant differences (P < 0.05) between pre-transplant plug cell volumes, while different**  
202 **capital letters indicate significant differences (P < 0.05) between pre- and post- growing**  
203 **media. For substrate abbreviations see Table 1. The probability of the RLA and RLAE**  
204 **slopes being zero was P < 0.001.**


205

	RLA (leaves week <sup>-1</sup> plant <sup>-1</sup> )		RLAE (cm <sup>2</sup> cm <sup>-2</sup> day <sup>-1</sup> )		SLA (cm <sup>2</sup> g <sup>-1</sup> )	
	Pre-	Post-	Pre-	Post-	Pre-	Post-
<b>F</b>						
50	0.512 <sup>aB</sup>	0.972 <sup>aA</sup>	0.0286 <sup>aB</sup>	0.0635 <sup>aA</sup>	188.43 <sup>CA</sup>	195.89 <sup>bA</sup>
128	0.449 <sup>bB</sup>	0.861 <sup>bA</sup>	0.0282 <sup>aB</sup>	0.0622 <sup>aA</sup>	230.92 <sup>aA</sup>	197.49 <sup>bB</sup>
288	0.340 <sup>cB</sup>	0.744 <sup>cA</sup>	0.0231 <sup>bB</sup>	0.0616 <sup>aA</sup>	238.04 <sup>aA</sup>	197.35 <sup>bB</sup>
512	0.291 <sup>dB</sup>	0.648 <sup>dA</sup>	0.0223 <sup>bB</sup>	0.0508 <sup>bA</sup>	215.16 <sup>bA</sup>	215.78 <sup>aA</sup>
<b>S</b>						
50	0.859 <sup>aA</sup>	0.627 <sup>aB</sup>	0.0363 <sup>aB</sup>	0.0567 <sup>aA</sup>	156.39 <sup>CB</sup>	204.58 <sup>dA</sup>
128	0.843 <sup>aA</sup>	0.624 <sup>aB</sup>	0.0371 <sup>aB</sup>	0.0552 <sup>aA</sup>	185.15 <sup>bB</sup>	226.07 <sup>CA</sup>
288	0.838 <sup>aA</sup>	0.547 <sup>bB</sup>	0.0317 <sup>bB</sup>	0.0543 <sup>aA</sup>	220.20 <sup>aA</sup>	224.42 <sup>bA</sup>
512	0.604 <sup>bA</sup>	0.518 <sup>bB</sup>	0.0280 <sup>CB</sup>	0.0441 <sup>bA</sup>	200.77 <sup>aA</sup>	294.62 <sup>aA</sup>
<b>R</b>						
50	1.000 <sup>aA</sup>	0.950 <sup>aB</sup>	0.0367 <sup>aB</sup>	0.0575 <sup>aA</sup>	179.64 <sup>bB</sup>	200.44 <sup>aA</sup>
128	0.902 <sup>bA</sup>	0.752 <sup>bB</sup>	0.0369 <sup>aB</sup>	0.0509 <sup>bA</sup>	184.61 <sup>aA</sup>	198.96 <sup>aA</sup>
288	0.788 <sup>cA</sup>	0.666 <sup>CB</sup>	0.0340 <sup>bB</sup>	0.0505 <sup>bA</sup>	186.52 <sup>aA</sup>	201.48 <sup>aA</sup>
512	0.764 <sup>cA</sup>	0.619 <sup>dB</sup>	0.0280 <sup>CB</sup>	0.0481 <sup>CA</sup>	183.03 <sup>aA</sup>	201.23 <sup>aA</sup>
<b>SR</b>						
50	0.949 <sup>aA</sup>	0.987 <sup>aA</sup>	0.0400 <sup>aB</sup>	0.0586 <sup>aA</sup>	176.86 <sup>bB</sup>	220.88 <sup>bA</sup>
128	0.704 <sup>bA</sup>	0.745 <sup>bA</sup>	0.0352 <sup>bB</sup>	0.0569 <sup>bA</sup>	179.18 <sup>bB</sup>	227.05 <sup>bA</sup>
288	0.549 <sup>CB</sup>	0.741 <sup>bA</sup>	0.0296 <sup>CB</sup>	0.0531 <sup>CA</sup>	187.80 <sup>aB</sup>	225.63 <sup>bA</sup>
512	0.543 <sup>CB</sup>	0.671 <sup>CA</sup>	0.0258 <sup>DB</sup>	0.0440 <sup>DA</sup>	189.70 <sup>aB</sup>	240.21 <sup>aA</sup>


206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227

The higher plug cell volume the higher RLA and RLAE but the lower SLA (leaves has a higher thickness). The response to a change in growing media quality at the pre- or post-transplant changed according pre-transplant plug cell volume and growing media tested in the same way that total leaf area (Table 2).

Stem numbers per plant showed significant differences in plants grown at nursery in different both plug cell volume and growing media; they were positively stimulated by a change in post-transplant growing media in plants from K-growing media. An inverse or no significant result was found when S-, R- or SR-growing media were used at the post-transplant stage (Fig. 3A). A similar response from the number node per plant was found (Fig. 3B).



228



229

230 Fig. 3. Stem (A) and node (B) number at the end of the experiments (60 days from  
 231 transplanting) for *Impatiens walleriana* plants from four plug cell volumes (50-, 128-, 288-,  
 232 and 512-cell tray<sup>-1</sup>) at the pre-transplant stage and four growing media at the pre- or post-  
 233 transplant stage. Bars indicate standard errors and vertical line indicate least significant  
 234 differences (LSD). For substrate abbreviations see Table 1.

235

236

### 237 3.3. Dry weight accumulation

238

239 Due there is no DW significant differences between treatments (data not shown), the traditional  
 240 growth analysis approach would be performed. During the experiments, RGR values were  
 241 significantly different from plants grown in different plug cell volume although data from post-  
 242 transplant growing media were higher than those from plants grown at the same growing media  
 243 at the pre-transplant stage. When RGR was separating from their 'physiological' (NAR) and  
 244 'morphological' (LAR) components, NAR decreased according plug cell volume decrease with  
 245 significant differences between growing media tested and time (lesser pre-transplanted plants  
 246 than post-transplanted ones). Quite opposite responses were found for LAR (Table 3).

247

248

249 **Table 3: Changes in RGR, NAR and LAR for *Impatiens walleriana* plants from four plug  
 250 cell volumes (50-, 128-, 288-, and 512-cell tray<sup>-1</sup>) at the pre-transplant stage and four  
 251 growing media at the pre- or post-transplant stage. Different lower-case letters indicate  
 252 significant differences (P < 0.05) between pre-transplant plug cell volumes, while different  
 253 capital letters indicate significant differences (P < 0.05) between pre- and post- growing  
 254 media. For substrate abbreviations see Table 1. The probability of the RGR slope being  
 255 zero was P < 0.001.**

256


257

	RGR (g g <sup>-1</sup> día <sup>-1</sup> )		NAR (g cm <sup>-2</sup> día <sup>-1</sup> ) x 10 <sup>-5</sup>		LAR (cm <sup>2</sup> g <sup>-1</sup> )	
	Pre-	Post-	Pre-	Post-	Pre-	Post-
<b>F</b>						
50	0.030 <sup>aB</sup>	0.059 <sup>aA</sup>	6.11 <sup>aB</sup>	18.30 <sup>aA</sup>	49.10 <sup>dA</sup>	32.24 <sup>cB</sup>
128	0.031 <sup>aB</sup>	0.059 <sup>aA</sup>	4.17 <sup>bB</sup>	10.27 <sup>bA</sup>	74.34 <sup>bA</sup>	57.45 <sup>bB</sup>
288	0.028 <sup>bB</sup>	0.058 <sup>aA</sup>	4.46 <sup>bB</sup>	10.15 <sup>bA</sup>	62.78 <sup>cA</sup>	57.14 <sup>bA</sup>
512	0.026 <sup>bB</sup>	0.052 <sup>aA</sup>	1.88 <sup>cB</sup>	7.46 <sup>cA</sup>	138.30 <sup>aA</sup>	69.71 <sup>aB</sup>
<b>S</b>						
50	0.038 <sup>aB</sup>	0.059 <sup>aA</sup>	5.95 <sup>aB</sup>	15.20 <sup>aA</sup>	63.87 <sup>cA</sup>	38.82 <sup>cB</sup>
128	0.041 <sup>aB</sup>	0.052 <sup>aA</sup>	4.96 <sup>bB</sup>	8.48 <sup>bA</sup>	82.66 <sup>aA</sup>	61.32 <sup>bB</sup>
288	0.035 <sup>bB</sup>	0.047 <sup>bA</sup>	4.84 <sup>bB</sup>	7.41 <sup>cA</sup>	72.31 <sup>bA</sup>	63.43 <sup>bB</sup>
512	0.035 <sup>bB</sup>	0.045 <sup>bA</sup>	4.24 <sup>cB</sup>	5.65 <sup>dA</sup>	82.55 <sup>aA</sup>	79.65 <sup>aA</sup>
<b>R</b>						
50	0.044 <sup>aA</sup>	0.051 <sup>aA</sup>	6.39 <sup>aB</sup>	13.44 <sup>aA</sup>	68.86 <sup>dA</sup>	37.95 <sup>cB</sup>
128	0.043 <sup>aA</sup>	0.054 <sup>aA</sup>	5.12 <sup>bB</sup>	6.69 <sup>bA</sup>	83.98 <sup>aA</sup>	80.72 <sup>bA</sup>
288	0.041 <sup>bA</sup>	0.048 <sup>aA</sup>	5.23 <sup>bA</sup>	5.11 <sup>cA</sup>	78.39 <sup>bB</sup>	93.93 <sup>aA</sup>
512	0.038 <sup>cB</sup>	0.046 <sup>aA</sup>	5.40 <sup>bA</sup>	4.73 <sup>dB</sup>	70.37 <sup>cB</sup>	97.25 <sup>aA</sup>
<b>SR</b>						
50	0.042 <sup>aB</sup>	0.053 <sup>aA</sup>	8.16 <sup>aB</sup>	14.34 <sup>aA</sup>	51.47 <sup>cA</sup>	36.96 <sup>cB</sup>
128	0.037 <sup>bB</sup>	0.053 <sup>aA</sup>	6.91 <sup>bB</sup>	12.86 <sup>bA</sup>	53.55 <sup>cA</sup>	41.21 <sup>cB</sup>
288	0.036 <sup>bB</sup>	0.052 <sup>aA</sup>	5.86 <sup>cA</sup>	5.54 <sup>cA</sup>	61.43 <sup>bB</sup>	93.86 <sup>bA</sup>
512	0.030 <sup>cB</sup>	0.048 <sup>aA</sup>	3.16 <sup>dA</sup>	2.08 <sup>dB</sup>	94.94 <sup>aB</sup>	230.77 <sup>aA</sup>

258

259 When plotting the data from all treatments, we found a close direct relationship ( $r^2 = 0.656$  and  
 260  $0.635$ ) for pre- and post-transplant values respectively between RGR and NAR (Fig. 4A) and an  
 261 inverse relationship between RGR and LAR ( $r^2 = 0.191$  and  $0.328$ ) (Fig. 4B).





263

264


265 **Figure 4: The net assimilation rate (NAR) (A) and the leaf area ratio (LAR) (B) related to**266 **the relative growth rate (RGR) for *Impatiens walleriana* plants from four plug cell volumes**267 **(50-, 128-, 288-, and 512-cell tray<sup>-1</sup>) at the pre-transplant stage and four growing media at**268 **the pre- or post-transplant stage. The probability of the NAR and LAR slopes being zero**269 **was  $P < 0.001$ . For substrate abbreviations see Table 1. F: ●-○; R: ■-□; S: ◆-◇; SR: ▲-△.**270 **The straight-line regressions were:  $NAR_{Pre} = 189.25 RGR - 1.67$  ( $r^2 = 0.656$ );  $NAR_{Post} =$** 271  **$739.27 RGR - 29.99$  ( $r^2 = 0.635$ );  $LAR_{Pre} = -1,596.20 RGR + 132.07$  ( $r^2 = 0.191$ ) and  $LAR_{Post} = -$** 272  **$5,572.40 RGR + 368.89$  ( $r^2 = 0.328$ ).**

273


274 Positive relationships between RLAE ( $r^2 = 0.645$   $P < 0.001$ ) (Fig. 5A), RLA ( $r^2 = 0.627$   $P < 0.001$ ) (Fig. 5B),275 RGR ( $r^2 = 0.665$   $P < 0.001$ ) (Fig. 5C), NAR ( $r^2 = 0.602$   $P < 0.001$ ) (Fig. 5D),

276 and root DW were found. The higher values were those from plants grown in different growing

277 media at the post-transplant stage.



278



279  
 280 Fig. 5. Relationship between RLAE (A), RLA (B), RGR (C), NAR (D) and root dry weight (RDW)  
 281 for *Impatiens walleriana* plants from four plug cell volumes (50-, 128-, 288-, and 512-cell  
 282 tray<sup>-1</sup>) at the pre-transplant stage and four growing media at the pre- (full symbols) or  
 283 post-transplant stage (empty symbols). The straight-line regressions were: RLAE = 0.067  
 284 RDW + 0.023 ( $r^2 = 0.645$  P < 0.001), RLA = 0.735 root DW + 0.44 ( $r^2 = 0.627$  P < 0.001), RGR =  
 285 0.036 RDW + 0.03 ( $r^2 = 0.665$  P < 0.001), NAR = 11.86 RDW + 2.69 ( $r^2 = 0.602$  P < 0.001). The  
 286 probability of the slopes being zero was P < 0.001. F:  $\blacklozenge$ - $\diamond$ ; R:  $\blacksquare$ - $\square$ ; S:  $\bullet$ - $\circ$ ; SR:  $\blacktriangle$ - $\triangle$ .

287

288

289

290

### 291 3.4. Photo assimilates partitioning

292

293 The higher plug cell volume the lower  $\beta$  coefficient for root: shoot allometries for all growing  
 294 media at the pre-transplant, which showed a higher photo assimilates partitioning to roots. The  
 295 same  $\beta$  response pattern was found during the post-transplant but absolute values were even  
 296 lower than for the pre-transplant growing media. An increase in root: shoot ratio according plug  
 297 cell volume decrease were found as well with significant differences between growing media  
 298 (Table 4).

299

300 Table 4: Changes in allometric relationships between roots and shoots for *Impatiens*  
 301 *walleriana* plants from four plug cell volumes (50-, 128-, 288-, and 512-cell tray<sup>-1</sup>) at the  
 302 pre-transplant stage and four growing media at the pre- or post-transplant stage.  
 303 Different lowercase letters indicate significant differences (P < 0.05) between pre-  
 304 transplant plug cell volumes, while different capital letters indicate significant differences  
 305 (P < 0.05) between pre- and post- growing media. For substrate abbreviations see Table  
 306 1. The probability of the slopes being zero was P < 0.001.

307

	Roots versus Shoots		Root: Shoot	
	Pre- $\beta$	Post- $\beta$	Pre-	Post-
F				
50	1.031 <sup>dA</sup>	0.889 <sup>cB</sup>	0.215 <sup>bA</sup>	0.147 <sup>bB</sup>
128	1.106 <sup>cA</sup>	0.911 <sup>bB</sup>	0.218 <sup>bA</sup>	0.163 <sup>bB</sup>
288	1.160 <sup>bA</sup>	0.987 <sup>aB</sup>	0.233 <sup>aB</sup>	0.298 <sup>aA</sup>
512	1.188 <sup>aA</sup>	0.965 <sup>aB</sup>	0.243 <sup>aB</sup>	0.317 <sup>aA</sup>
S				
50	1.004 <sup>cA</sup>	0.773 <sup>dB</sup>	0.163 <sup>cB</sup>	0.212 <sup>cA</sup>
128	1.117 <sup>bA</sup>	0.841 <sup>cB</sup>	0.173 <sup>cB</sup>	0.255 <sup>bA</sup>
288	1.117 <sup>bA</sup>	0.900 <sup>bB</sup>	0.207 <sup>bB</sup>	0.270 <sup>bA</sup>
512	1.165 <sup>aA</sup>	1.157 <sup>aB</sup>	0.264 <sup>aB</sup>	0.311 <sup>aA</sup>

R				
50	1.032 <sup>bA</sup>	0.795 <sup>cB</sup>	0.360 <sup>cA</sup>	0.178 <sup>cB</sup>
128	1.040 <sup>bA</sup>	0.623 <sup>dB</sup>	0.458 <sup>bA</sup>	0.181 <sup>cB</sup>
288	1.182 <sup>aA</sup>	0.905 <sup>aB</sup>	0.460 <sup>bA</sup>	0.223 <sup>bB</sup>
512	1.164 <sup>aA</sup>	0.871 <sup>bB</sup>	0.556 <sup>aA</sup>	0.298 <sup>aB</sup>
SR				
50	1.030 <sup>bA</sup>	0.853 <sup>cB</sup>	0.155 <sup>cB</sup>	0.185 <sup>bA</sup>
128	1.037 <sup>bA</sup>	0.961 <sup>bB</sup>	0.165 <sup>bB</sup>	0.195 <sup>bA</sup>
288	1.065 <sup>aA</sup>	0.981 <sup>bB</sup>	0.188 <sup>bB</sup>	0.208 <sup>bA</sup>
512	1.086 <sup>aA</sup>	1.002 <sup>aB</sup>	0.227 <sup>aB</sup>	0.274 <sup>aA</sup>

308

309

310

#### 4. DISCUSSION

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

In a recent previous report [24], we have found that, in *I. walleriana* seedlings, the abiotic stress imposed by the growing medium quality during nursery had a higher effect on biomass accumulation (on both fresh and dry base), leaf area expansion and photo assimilates partitioning than plug cell volume and constitute an interactive process associated with cytokinin synthesis. However, this novelty approach did not exclude the plug cell volume involvement as a limiting abiotic stress source during other parts of the *I. walleriana* growth cycle. In this context we have evaluated pot biomass accumulation of this bedding plant to four growing media at both nursery and pot stage in plants propagated in four different plug cell volumes.

A 'root restriction' syndrome related to either plug cell volume or growing media quality is an exogenous signal, which let plants to sense the volume space and decrease or increase root system accordingly [31,32]. Having in mind that **root is** a major source of cytokinins [33], which control the source of biomass accumulation such as the shoot apical meristem [34], the similarity between *I. walleriana* plants grown in the best growing conditions and exogenous cytokinin-sprayed plants [23,24,35], it is not unexpected. Our results from Fig. 5, which shown a positive relationships between the most growing parameters performed (RLAE, RLA, RGE, NAR) according to a root dry weight increases are in agreement with this previous reports. On the other hand, a positive relationship between **shoots and roots** fresh weight (there were not significant differences between fresh and dry weight) was found (Fig. 1B) in agreement with previous reports [4,15,17,21,22,23,24].

Although growing media performed through a high quality *Sphagnum sp.* peat base (Klasman® commercial growing media) has been indicated as the best for pot plants [1], *I. walleriana* previous results indicated that better growing media for this bedding plant can be found [36].

Results showed significant fresh weight changes at the end of the experiments (Fig. 1A) according a decrease in plug cell volume and a change in growing media quality, which are in agreement to RGR changes (Table 3). Methodology usually used to describe changes in biomass accumulation on both fresh and dry weight included: (i) stems appearance; (ii) leaf area expansion; (iii) photosynthetic capacity and; (iv) photo assimilate partitioning. In ornamental bedding plants, additional traits such as tolerance to biotic and abiotic stresses and aesthetics must be included [37].

Stem branching is an important aspect to consider in ornamental bedding plants because it **takes** part as the biomass accumulation as the aesthetic appearance. Results from Fig. 3 indicated that both shoot number and node number decreased according to plug cell volume decrease, which indicates a desirable commercial ideotype with a lower branching and compact growth habit. However, growing media quality **changes** the impact **on** the response to different pre-transplant plug cell volume.

Aesthetically, total leaf area is the main trait related to plant quality for commercial acceptance of ornamentals and it determines the time of plant sale and at the same time, leaves are the plant organs responsible to light interception. In physiological terms, it implies to expand leaf

355 area at the higher growth rate which included both individual leaf sizes and leaf numbers. Data  
356 from Fig. 2A shown that the total leaf area, with minor effects on individual leaf area (Fig. 2B)  
357 was significantly affect by both plug cell volume and growing media quality. Three growth  
358 parameters can be used to characterize leaf area development: (i) RLA, which is an estimator of  
359 leaf initiation and plastochron length, (ii) RLAE, which let to quantify leaf expansion and, (iii)  
360 SLA, which characterize leaf thickness. Data from Table 2 showed that a decrease in plug cell  
361 volume and a change in growing media quality decreased RLA and RLAE while increased SLA.  
362 These results implies that the changes in total leaf area are mainly related to the meristematic  
363 shoot apex capacity to initiate and to expand leaf primordia [38]. Both processes are mediated  
364 by the down regulation of *KNOTTED* and *WUSCHEL* genes [39] associated to a high cytokinin:  
365 low gibberellin ratio [40]. On the other hand, the lower SLA the higher leaf thickness, which it is  
366 a pre-requisite for a high photosynthetic rate [41]. In this way, Gandolfo et al. [9] found positive  
367 relationships between leaf thickness, intercellular spaces and NAR in *I. walleriana* root-  
368 restricted plants. When the mesophyll thickness of the leaf is increased, the maximum  
369 photosynthetic rate increased as well. This probably explains the strong relationship between  
370 NAR and mesophyll thickness.

371

372 Variation in RGR has the result of two key traits: the 'physiological component' NAR and the  
373 'morphological component' LAR. RGR, which ultimate quantify biomass accumulation, is  
374 greatly influenced by photosynthetic efficiency. Although the higher the plug cell volume the  
375 higher the RGR and NAR, growing media quality at the post-transplant stage increased both  
376 growth parameters (Table 3). Shipley [42] indicated that, in general, NAR was the best general  
377 predictor of variation in RGR, in agreement with our results from Fig. 4. Root restrictions often  
378 depresses photosynthetic capacity [44] and decreased energy synthesis [45]. The positive  
379 relationships between NAR and RGR (Figure 4A) are in agreement with Shi et al. [44,45].

380

381 Root-restricted plants change photo assimilates partition as a response to abiotic stresses  
382 (Table 4). At the end of the experiments, the higher root restriction the higher root: shoot ratio.  
383 Root: shoot allometries let to explain these results because showed a higher photo assimilates  
384 partitioning to roots (lower  $\beta$  coefficients) during the greater part of the experiments in root-  
385 limited treatments.

386

387 As opposed to a previous report [23], which indicate that that growing media quality would be a  
388 more limited factor than plug cell volume for *I. walleriana* seedlings during nursery, our results  
389 showed that both abiotic stresses would be interactive restricting technological factors during the  
390 post-transplant pot stage.

391

392

## 393 5. CONCLUSIONS

394

395 The effect of an abiotic stress and the relationships between multiples stress sources is not the  
396 same according to the plant growth stage. In this context, different *I. walleriana* growth  
397 responses to both plug cell volume and growing media found in our experiments would not be  
398 unexpected results but to extend to other ornamental bedding plants and to perform a  
399 commercial suggestion much more research must be required.

400

401

402

403

## 404 REFERENCES

405

- 406 1. Di Benedetto A. Root restriction and posttransplant effects for bedding pot plants. In:  
407 Ornamental Plants: Types, Cultivation and Nutrition. (J.C. Aquino, ed.), Nova Science  
408 Publishers, Inc. NY, USA. 2011;47-79.
- 409 2. Blok C, Verhagen JBG. Trends in rooting media in Dutch horticulture during the period  
410 2001–2005: The new growing media project. Acta Hort. 2009;819:47-58.
- 411 3. Di Benedetto A, Pagani A. Difficulties and possibilities of alternative substrates for  
412 ornamental bedding plants: An ecophysiological approach. In: Draguhn C, Ciarimboli N,  
413 editors. Peat: Formation, Uses and Biological Effects, Nova Science Publishers, Inc.  
414 NY, USA; 2012.

- 415 4. Pagani A, Molinari J, Lavado RS, Di Benedetto A. Behavior of *Impatiens wallerana*  
416 Hook. f in alternative pot substrates: mechanisms involved and research perspectives.  
417 J. Plant Nutr. 2015;38:2185-2203.
- 418 5. Di Benedetto A, Klasman R. The effect of plug cell volume on the post-transplant  
419 growth for *Impatiens walleriana* pot plant. Eur. J. Hort. Sci. 2004;69(2):82-86.
- 420 6. Di Benedetto A, Klasman R. The effect of plug cell volume, paclobutrazol height control  
421 and the transplant stage on the post-transplant growth for *Impatiens wallerana* pot  
422 plant. Eur. J. Hort. Sci. 2007;72(5):193-197.
- 423 7. Di Benedetto A, Molinari J, Rattin J. The effect of transplant in sweet maize (*Zea mays*  
424 L.). II. Container root restriction. Int. J. Agric. Res. 2006;1 (6):555-563.
- 425 8. Di Benedetto A, Pagani A. Dry weight accumulation in the *Impatiens walleriana* pot  
426 plant in responses to different pretransplant plug cell volume. Europ. J. Hort. Sci.  
427 2013;78(2):76-85.
- 428 9. Gandolfo E, De Lojo E, Gómez E, Pagani A, Molinari J, Di Benedetto A. Anatomical  
429 changes involved in the response of *Impatiens wallerana* to different pretransplant plug  
430 cell volumes and BAP sprays. Eur. J. Hort. Sci. 2014;79(4):226-232.
- 431 10. Thibaud J, Mc. Loughlin T, Pagani A, Lavado R, Di Benedetto A. Alternative substrates  
432 and fertilization routine relationships for bedding pot plants: *Impatiens wallerana*. Eur. J.  
433 Hort. Sci. 2012;77(4):182-191.
- 434 11. Bailey-Serres J, Chang R. Sensing and signalling in response to oxygen deprivation in  
435 plants and other organisms. Ann. Bot. 2005;96(4):507-518.
- 436 12. Voesenek LACJ, Sasidharan R. Ethylene-and oxygen signalling—drive plant survival  
437 during flooding. Plant Biol. 2013;15(3):426-435.
- 438 13. Gandolfo E, Hakim E, Geraci J, Feuring V, Giardina E, Di Benedetto A. Responses of  
439 pansy (*Viola wittrockiana* Gams.) to the quality of the growing media. Amer. J. Exp.  
440 Agric. 2016;12(3):1-10.
- 441 14. Hakim G, Gandolfo E, Giardina E, Di Benedetto A. The effect of the pretransplant pot  
442 media quality on pansy garden performance. Int. J. Plant Soil Sci. 2017;19(4):1-12.
- 443 15. Molinari J, Pagani A, Buyatti M, Giardina E, Di Benedetto A. *Impatiens* 'New Guinea'  
444 (*Impatiens hawkeri* Bull) propagation effects on the pre-transplant biomass  
445 accumulation. Int. J. Adv. Agric., 2018;3(9):1-15.
- 446 16. Molinari J, Buyatti M, Pagani A, Giardina E, Di Benedetto A. New Guinea *Impatiens*  
447 (*Impatiens hawkeri* Bull) hormonal effects on the post-transplant biomass accumulation.  
448 J. Adv. Agric. 2019;10:1629-1649.
- 449 17. Piotti M, Molinari J, Pagani A, Giardina E, Di Benedetto A. Root restriction effects on  
450 *Zantedeschia aethiopica*. Int. J. Plant Soil Sci. 2018;23(2):1-15.
- 451 18. Coro M, Araki A, Rattin J, Miravé P, Di Benedetto A. Lettuce and celery responses to  
452 both BAP and PBZ related to the plug cell volume. Amer. J. Exp. Agric.  
453 2014;4(10):1103-1119.
- 454 19. Di Matteo J, Rattin J, Di Benedetto A. Increase of spinach growth through the use of  
455 larger plug cell volume and an exogenous BAP spray. Amer. J. Exp. Agric.  
456 2015;6(6):372-383.
- 457 20. Rattin J, Pico Estrada O, Giardina E, Di Benedetto A. Nursery pre- and posttransplant  
458 effects on tomato (*Solanum lycopersicum* L.) growth and yield. J. Exp. Agric. Int.  
459 2017;18(5):1-14.
- 460 21. Geraci J, Di Matteo J, Feuring V, Giardina E, Di Benedetto A. Exogenous BAP spray on  
461 biomass accumulation in spinach. J. Exp. Agric. Int. 2018;25(6):1- 17.
- 462 22. Williams T, Sala A, Feuring V, Giardina E, Di Benedetto A. Relationships between plug  
463 cell size and quality substrate in the bedding pot plant *Impatiens wallerana* (Hook.f.).  
464 Int. J. Plant Soil Sci. 2016;12(2): 1-12.
- 465 23. DeLojo J, Gandolfo E, Gómez D, Feuring V, Monti S, Giardina E, Boschi C, Di  
466 Benedetto A. Root restriction effects on the bedding pot plant *Impatiens walleriana*. J.  
467 Exp. Agric. Int. 2017;15(4):1-16.
- 468 24. De Lojo J, Gandolfo E, Boschi C, Giardina E, Di Benedetto A. Pot cell volume, growing  
469 media quality and benzyl aminopurine (BAP) spray effects for nursery growth of  
470 *Impatiens walleriana*. International J. Plant Soil Sci. 2019;27(6):1-13.
- 471 25. Faust JE, Holcombe V, Rajapakse NC, Layne DR. The effect of daily light integral on  
472 bedding plant growth and flowering. Hort Science. 2005;40(3):645-649.
- 473 26. Pramuk LA, Runkle ES. Modeling growth and development of Celosia and Impatiens in  
474 response to temperature and photosynthetic daily light integral. J. Amer. Soc. Hort. Sci.

- 475 2005;130(6):813-818.
- 476 27. Henson DY, Newman SE, Hartley DE. Performance of selected herbaceous annual  
477 ornamentals grown at decreasing levels of irrigation. HortScience. 2006; 41(6):1481-  
478 1486.
- 479 28. Fonteno WC. Growing Media Types and Physical/Chemical Properties. In: Water,  
480 Media and Nutrition for Greenhouse Crops. A Grower's Guide. (Reed, D.W., Ed.), Ball  
481 Publishing, Batavia, Illinois, U.S.A.; 1996.
- 482 29. Potter J, Jones JW. Leaf area partitioning as an important factor in growth. Plant  
483 Physiol. 1977;59(1):10-14.
- 484 30. Warton DI, Duursma RA, Falster DS, Taskinen S. SMATR 3-an R package for  
485 estimation and inference about allometric lines. Methods Ecol. Evol. 2012;3(2):257-259.
- 486 31. Puig J, Pauluzzi G, Guiderdoni E, Gantet P. Regulation of shoot and root development  
487 through mutual signaling. Mol. Plant. 2012;5(5):974-983.
- 488 32. Chen BJ, Daring HJ, Vermeulen PJ, Kroon H, Poorter H, Anten NP. Corrections for  
489 rooting volume and plant size reveal negative effects of neighbour presence on root  
490 allocation in pea. Functional Ecol. 2015;29(11):1383-1391.
- 491 33. Yruela I. Plant development regulation: Overview and perspectives. J. Plant Physiol.  
492 2015;182(5): 62-78.
- 493 34. Bartoli CG, Casalengué CA, Simontacchi M, Marquez-Garcia B, Foyer CH. Interactions  
494 between hormone and redox signalling pathways in the control of growth and cross  
495 tolerance to stress. Environ. Exp. Bot. 2013;94:73-88.
- 496 35. Zwack PJ, Rashotte A.M. Interactions between cytokinin signalling and abiotic stress  
497 responses. J. Exp. Bot. 2015; 66(16):4863-4871.
- 498 36. Di Benedetto A, Petracchi JC, Marcella G, Montaron P, Chavez P. Evaluation of  
499 alternative substrates for bedding plants. Int. J. Agric. Res. 2006;1(6):545-554.
- 500 37. Santagostini P, Demotes-Mainard S, Huché-Thélier L, Leduc N, Bertheloot J, Guérin V,  
501 Bourbeillon J, Sakr S, Boumazad R. Assessment of the visual quality of ornamental  
502 plants: Comparison of three methodologies in the case of the rose bush. Scientia Hortic.  
503 2014;168:17-26.
- 504 38. Perilli S, Moubayidin L, Sabatini S. The molecular basis of cytokinin function. Curr. Op.  
505 Plant Biol. 2010;13(1):21-26.
- 506 39. Holt, A. L.; Van Haperen, J.M.; Groot, E.P.; Laux, T. Signaling in shoot and flower  
507 meristems of *Arabidopsis thaliana*. Current Opinion in Plant Biology, 2014, 17, 96-102.
- 508 40. Hay A, Tsiantis M. KNOX genes: Versatile regulators of plant development and  
509 diversity. Develop. 2010;137(19):3153-3165.
- 510 41. Milla R, Reich PB, Niinemets Ü, Castro-Díez P. Environmental and developmental  
511 controls on specific leaf area are little modified by leaf allometry. Funct. Ecol.  
512 2008;22:565-576.
- 513 42. Shipley B. Net assimilation rate, specific leaf area and leaf mass ratio: Which is most  
514 closely correlated with relative growth rate? A meta-analysis. Funct. Ecol. 2006;  
515 20(4):565-574.
- 516 43. Shi K, Ding XT, Dong DK, Zhou YH, Yu JQ. Root restriction-induced limitation to  
517 photosynthesis in tomato (*Lycopersicon esculentum* Mill.) leaves. Scientia Hortic.  
518 2008;117(3):197-202.
- 519 44. Shi K, Fu LJ, Dong DK, Zhou YH, Yu JQ. Decreased energy synthesis is partially  
520 compensated by a switch to sucrose synthase pathway of sucrose degradation in  
521 restricted root of tomato plants. Plant Physiol. Biochem. 2008;46(12):1040-1044.  
522