

SCREENING FOR HYPERTENSION AND DIABETES IN AN UNDERSERVED POPULATION THROUGH COMMUNITY OUTREACH; A CASE OF RURAL COMMUNITY IN ENUGU STATE, NIGERIA

ABSTRACT

Introduction

Hypertension and Diabetes are the commonest co-morbidity of each other and are among the principal cause of the burden of non-communicable diseases in developing countries. It is important to identify patients with these conditions early in the disease process. This study was to determine the prevalence of elevated Blood Pressure (BP) and elevated Fasting Blood Sugar (FBS) as well as relate it to the characteristics of the study participants in a rural community in Enugu State, Nigeria.

Methods

Community based cross-sectional study in form of outreach was done. The study was conducted over 1 week period among participants aged 18 years and above. Proforma was used in collecting information on characteristics of participants including age, sex and occupation. Measurements of BP, FBS and BMI were done. Chi square test and Binary Logistic Regression were used for analysis.

Results

Majority of participants were aged > 45 years 127(56.7%), and females 139(62.1%), Mean(SD) 46.89((21.84) Elevated BP 55(24.6%), elevated FBS 42(18.8%), both elevated BP and FBS 13(5.8%). higher proportion of those aged > 45 years had elevated BP 51(92.7%) and elevated FBS 37(88.1%). More Females had elevated BP 35(63.6%) and elevated FBS 28(66.7%). Predictors were; age >45 years for elevated BP (AOR 18.4; 95% CI 5.7-59.5) and for have elevated FBS (AOR 8.9; 95% CI 3.0-26.5).

32 **Conclusion**

33 Prevalence of raised BP and FBS as well as co-morbid condition was high. It was more among
34 females and older age. Age was a predictor of both raised BP and FBS. This calls for
35 interventional programmes that will assist in limiting the increasing burden of the diseases in
36 rural communities

37

38 **Keywords;** Raised Blood Pressure, raised Blood sugar, Screening, Outreach, rural community

39

40 **INTRODUCTION**

41 Non-communicable diseases (NCD), essentially cardiovascular diseases like Hypertension,
42 Diabetes, Cancer and chronic respiratory diseases, are responsible for about 68% (38 million) of
43 the 56 million deaths that occurred globally during 2012.¹ Practically, 80% of these NCD deaths
44 (29 million) occurred in low and middle-income countries (LMICs).¹ In addition, the African
45 region of the world is experiencing a double epidemic of both communicable and non-
46 communicable diseases. It is reported that in Sub-Saharan Africa, the menace of NCDs could
47 surpass that of communicable diseases in the nearest future.²

48 Hypertension and diabetes are among the important diseases responsible the high burden of non-
49 communicable diseases in developing countries.² Both have significantly contributed to medical
50 morbidity and mortality. Hypertension is the commonest co-morbidity of diabetes and diabetes is
51 the commonest co-morbidity of hypertension. They have common genetic predisposition, share
52 risk factors, and have similar environmental influences as causative factors, and are interrelated.
53 They bring about an enormous financial stress on individuals, families, communities and the
54 health system of any country.³ Worldwide, Diabetes is responsible for about USD 727 billion
55 dollars in health expenditure and 12% of total expenditure on health of adults in 2017.⁴ Fuelling
56 the crisis is the upsurge in unhealthy lifestyle such as tobacco and alcohol consumption, the

57 reduction of physical activity and the changes in dietary intake due to westernization of our
58 culture.²

59 In 2012, report according to The World Health Statistics was that one in three adults worldwide
60 has an elevated Blood Pressure (BP) and one in 10 adults have diabetes.⁵ The raised BP is
61 responsible for about 50 percent of all deaths from stroke and heart disease.⁵. The percentage of
62 deaths attributed to raised blood glucose in those aged 20–69 years in LMICs was 60.5% in men
63 and 45.6% in women.² There was doubling in the prevalence of diabetes between 1980 and 2014.
64 Approximately 425 million adults (20-79 years) were living with diabetes in 2017 and is
65 projected that by 2045 this will rise to 629 million.⁴ The International Diabetic Federation (IDF)
66 statistics on diabetes reported that Nigeria has the largest population of people living with
67 diabetes and impaired fasting glucose in Africa.⁶ Hypertension is also the commonest
68 cardiovascular disease reported in the country.⁷ Findings from studies done in Nigeria
69 documented that the prevalence of diabetes varies across different regions of the country with
70 range from 2.2 - 9.8%.⁸⁻¹¹ In line with this, other similar studies reported varying prevalence
71 rates of hypertension in various regions of the country.^{8,12,13}

72 To reduce the prevalence and consequences of hypertension and diabetes a complimentary
73 mixture of population-wide and individual interventions is required. Allowing persons to present
74 at health facilities is not yielding desired result as majority of persons do not know that that they
75 have these silent killers. Likewise, there might be an enormous burden of hypertension and
76 diabetes mellitus among underserved rural dwellers in the country. It is important to identify
77 patients with these conditions early in the disease process. Periodic outreaches will contribute to
78 early detection of persons with such conditions. Prompt and effective referral following such
79 detection help reduce morbidity and complications associated with the diseases. This study was
80 to determine the prevalence of elevated Blood Pressure (BP) and elevated Fasting Blood Sugar

81 (FBS) among rural residents of a community in Enugu State Nigeria well as to find out the risk
82 factors associated with these conditions.

83 **METHODS**

84 **3.1. Study Area**

85 This was at Abor, a rural community in Udi Local government of Enugu state, South East
86 Nigeria. The community is about 30km to Enugu Metropolis They engage in farming, trading as
87 well as civil/public services. There are no well established health facilities even though there is a
88 primary health care facility.

89 **Study Design and population**

90 A Community based cross-sectional study in form of outreach was done. The study was
91 conducted over 1 week period. All participants aged 18 years and above who gave informed
92 consent were included in the study

93 **Sampling Technique and Sample Size Determination**

94 All participants who meet the inclusion criteria were recruited consecutively as they present for
95 screening at the outreach venue throughout the period of study. A total of 224 patients were
96 studied.

97 **Data collection tools and method**

- 98 1. Proforma was designed and used in collecting information on characteristics of
99 participants including age, sex, occupation, recording BP, FBS and BMI
- 100 2. Measurements of BP, FBS and BMI were done using standard tools and observing
101 standard procedure

102 **Blood pressure:** The BP was measured in the sitting position with an appropriate sized cuff
103 encircling the left arm held at the level of the heart. This was measured using the OMRON Arm-
104 type fully Automatic Digital Blood Pressure Monitor, Model BP - 103H. Raised BP was defined
105 using Joint National Committee on Hypertension (JNC) 7 classification as systolic BP \geq 140
106 mmHg and/or diastolic BP \geq 90 mmHg

107 **Blood sugar:** One microliter (1 μ L) of whole blood was collected and tested for blood glucose
108 level using the Accu-chek active test strip and glucometer (Roche Diagnostics GmbH,
109 Mannheim, Germany). Aseptic conditions were maintained throughout the procedure. Diabetes
110 was defined as a fasting blood glucose $>$ 110 mg/dl (6.1mmol/l)

111 **Weight and height:** the weight and height of the respondents were measured using a
112 standardized Stadiometer. Weight was measured to the nearest 0.5kg with the subject standing
113 motionless on the calibrated scale without footwear. Height was measured with the subject
114 standing in an erect position and head positioned so that the top of the external auditory meatus
115 was level with the inferior margin of the bony orbit. The BMI of the subjects was calculated as
116 weight in kilograms divided by height in meters squared.

117 **Data Collection and analysis**

118 Patient information were recorded by trained health workers to ensure accuracy of data.
119 Measurements of BP and FBS were done by qualified medical doctors. IBM Statistical Package
120 for Social Sciences Version 21 was used for data entry, editing and analysis. Results were
121 presented in tables. Mean, Standard deviation, proportion and percentages were used as summary
122 measures where appropriate. Chi square test was used to establish associations between

123 characteristics of participants with BP and FBS status. Binary Logistic Regression was done for
124 variables significant variables (age). Level of confidence was at $p < 0.05$.

125

126 **Ethical consideration**

127 The Health Research and Ethics committee of University of Nigeria Teaching Hospital,
128 Enugu gave ethical clearance. Permission was equally obtained from traditional rulers of
129 constituents communities. Informed consent was obtained from participant,. They were
130 ensured of voluntary participation and confidentiality of their information.

131

132 **Action taken**

133 Participants found to have elevated BP and/or FBS during the screening were provided education
134 on appropriate lifestyle and dietary modifications, such as salt and fatty reduction as well as need
135 for improved physical activity where not adequate. They were also instructed and referred to
136 tertiary health care facilities.

137 **RESULTS**

138 **Table 1: Characteristics of participants**

Variables	Frequency	Percent(%)
Age (Yrs)		
≤ 45	97	43.3
>45	127	56.7
<i>Mean(SD)</i>	<i>46.89(21.84)</i>	
Gender		
Female	139	62.1
Male	85	37.9
Occupation		
Civil/public servant	63	28.1

Trading	30	13.4
Farming	40	17.9
Skilled worker	18	8.0
Unemployed/student	73	32.6
BMI		
<18.5	66	29.5
18.5-24.9	99	44.2
25-29.9	57	25.4
≥30	2	.9

139
140 Table 1 shows that majority of participants were aged > 45 years 127(56.7%) with mean age of
141 46.89 SD of 21.84 were females 139(62.1%), unemployed/students 73(32.6%) followed by
142 Civil/public servants 63(28.1%) and 99(44.2%) had BMI of 18.5-24.9 mg/m²

143
144 **Table 2: Screening status of participants**

Variables	Frequency	Percent(%)
Blood pressure (BP)		
Normal	169	75.4
Elevated	55	24.6
Fasting Blood Sugar (FBS)		
Normal	182	81.3
Elevated	42	18.8
Both BP and FBS		
No	211	94.2
Elevated	13	5.8

145
146
147 Table 2 shows that 55(24.6%) of participants had elevated Blood Pressure, 42(18.8%) had
148 elevated Fasting Blood Sugar while 13(5.8%) both have elevated BP and FBS.

149 **Table 3: Blood pressure and Fasting Blood Sugar disaggregated by Characteristics of**
 150 **participants**

Variables	Blood pressure		Fasting Blood Sugar	
	Normal Freq(%)	Elevated Freq(%)	Normal Freq(%)	Elevated Freq(%)
Age (Yrs)				
≤ 45	93(55.0)	4(7.3)	92(50.5)	5(11.9)
>45	76(45.0)	51(92.7)	90(49.5)	37(88.1)
Gender				
Female	104(61.5)	35(63.6)	111(61.0)	28(66.7)
Male	65(38.5)	20(36.4)	71(39.0)	14(33.3)
Occupation				
Civil/public servant	41(24.3)	22(40.0)	44(24.2)	19(45.2)
Trading	26(15.4)	4(7.3)	24(13.2)	6(14.3)
Farming	26(15.4)	14(25.5)	33(18.1)	7(16.7)
Skilled worker	14(8.3)	4(7.3)	17(9.3)	1(2.4)
Unemployed/student	62(36.7)	11(20.0)	64(35.2)	9(21.4)
BMI				
<18.5	54(32.0)	12(21.8)	56(30.8)	10(23.8)
18.5-24.9	77(45.6)	22(40.0)	81(44.5)	18(42.9)
25-29.9	37(21.9)	20(36.4)	43(23.6)	14(33.3)
≥30	1(0.6)	1(1.8)	2(1.1)	0(0.0)

151
 152 Table 3 shows that higher proportion of those aged > 45 years had elevated Blood pressure
 153 51(92.7%) and elevated FBS 37(88.1%). More Females had elevated Blood pressure 35(63.6%)
 154 and elevated FBS 28(66.7%).. More Civil/public servants had elevated Blood pressure
 155 22(40.0%) and elevated FBS 19(45.2%). Higher proportion of those that had BMI of 18.5-
 156 24.9 18.5-24.9 mg/m² had elevated Blood pressure 22(40.0%) and elevated FBS 18(42.9%).

157 **Table 4: Relationship of Characteristics of participants with Blood Pressure and Fasting**
 158 **Blood Sugar**

BLOOD PRESSURE				
	Normal	Elevated	χ ² (p value)	AOR (95% CI of AOR)

Variables	Freq(%)	Freq(%)		
Age (Yrs)				
≤ 45	93(95.9)	4(4.1)	38.547(<0.001)	1
>45	76(59.8)	51(40.2)		18.36(5.66-59.54)
Gender				
Female	104(74.8)	35(25.2)	0.078(0.781)	NA
Male	65(76.5)	20(23.5)		
Occupation				
Civil/public servant	41(65.1)	22(34.9)		
Trading	26(86.7)	4(13.3)	11.648(0.020)	NA
Farming	26(65.0)	14(35.0)		
Skilled worker	14(77.8)	4(22.2)		
Unemployed/student	62(84.9)	11(15.1)		
BMI				
<18.5	54(81.8)	12(18.2)		
18.5-24.9	77(77.8)	22(22.2)	5.850(0.119)	NA
25-29.9	37(64.9)	20(35.1)		
≥30	1(50.0)	1(50.0)		
FASTING BLOOD SUGAR				
	Normal	Elevated		
Age (Yrs)				
≤ 45	92(94.8)	5(5.2)	20.757(<0.001)	1
>45	90(70.9)	37(29.1)		8.92(3.00-26.52)
Gender				
Female	111(79.9)	28(20.1)	0.467(0.494)	NA
Male	71(83.5)	14(16.5)		
Occupation				
Civil/public servant	44(69.8)	19(30.2)		
Trading	24(80.0)	6(20.0)	9.487(0.050)	NA
Farming	33(82.5)	7(17.5)		
Skilled worker	17(94.4)	1(5.6)		
Unemployed/student	64(87.7)	9(12.3)		
BMI				
<18.5	56(84.8)	10(15.8)		
18.5-24.9	81(81.8)	18(18.2)	FT(0.454)	NA
25-29.9	43(75.4)	14(24.6)		
≥30	2(100.0)	0(0.0)		

159 Table 4 shows that there were statistically significant association of blood pressure with age ($\chi^2 =$
160 38.547, $p < 0.001$), Occupation ($\chi^2 = 11.648$, $p = 0.020$). However there were no statistically
161 significant association of blood pressure with gender ($\chi^2 = 0.078$, $p = 0.781$). and BMI ($\chi^2 = 5.850$,

162 p = 0.119). Also, there were statistically significant association of Fasting blood sugar with age
163 ($\chi^2 = 20.757$, p < 0.001). However there were no statistically significant association of blood
164 pressure with gender ($\chi^2 = 0.467$, p = 0.494), Occupation ($\chi^2 = 9.487$, p = 0.050) and BMI (FT, p
165 = 0.454).

166 Those aged >45 years were about 18 times (AOR 18.4; 95% CI 5.7-59.5 likely to have elevated
167 BP than those aged \leq 45 years. Also those aged >45 years were about 9 times (AOR 8.9; 95% CI
168 3.0-26.5 likely to have elevated BP than those aged \leq 45 years.

169 DISCUSSION

170 Hypertension and Diabetes are the commonest co-morbidity of each other. They have common
171 genetic predisposition, share risk factors, and have similar environmental influences as causative
172 factors, and are interrelated¹⁴ Both elevated Blood Pressure and elevated Fasting Blood Sugar
173 occur more with advancing age. Result from this study reported that majority of participants
174 were aged > 45 years and females. This is expected as most rural areas are inhabited by retired
175 workers and older persons due urban migration in the country for greener pasture. Majority being
176 females can be partly explained by the better health seeking behavior of females compared
177 to males.

178 It was also noted that generally, 24.6% of participants had elevated Blood Pressure, 18.8% had
179 elevated Fasting Blood Sugar and 5.8% had both elevated BP and FBS. The reported prevalence
180 for elevated Blood Pressure was lower than the 42.0%, 44.5% and 46.4% reported respectively
181 different studies in South Eastern Nigeria.¹⁵⁻¹⁷ It is similar to reports from other studies.^{12,13} The
182 observed differences may be due differences in sampling technique and location of the study as
183 most of those previous studies were in urban areas whose life style is different from rural

184 communities. However, these finding is revealing as it shows that hypertension, DM and co
185 morbid condition are of high prevalence in rural communities in Nigeria. This calls for
186 interventional programmes including; **persistent** health education, enlightenment campaigns and
187 community surveillance programmes to **aid reduce** this **growing burden of the diseases in** rural
188 communities.

189 Based on Gender, 25.2% females and 23.5% males had elevated Blood Pressure, 20.1% females
190 and 16.5% males had elevated Fasting Blood Sugar, 5.0% females and 7.1% males had both
191 elevated BP and FBS. More Females equally had elevated Blood pressure 63.6% and elevated
192 FBS 66.7%. The higher prevalence among females were also documented in previous similar
193 studies.^{12,20-23}. However, there was discordance with many other previous reports. A study
194 involving review of studies on hypertension over five decades **reported a similar prevalence in**
195 **men and women with range in prevalence of 8% to 46.45%.**¹³ A Meta analysis of the prevalence
196 of hypertension from population based studies in south western Nigeria **reported a higher**
197 **prevalence in men than women with prevalence ranging from 12.4% to 34.8%.**¹² Another study
198 documented prevalence of HTN of 22% (25.9% in males and 20% in females). Similarly other
199 studies had similar findings of higher prevalence among males.¹⁸⁻²¹ The findings from this study
200 can partly be explained by fact that women are generally more likely than men to say they are
201 unwell.

202 Findings also show that those aged >45 years were about 18 times likely to have elevated BP
203 than those aged ≤ 45 years as well as about 9 times likely to have elevated BP than those aged ≤
204 45 years. Also Higher proportion of those aged > 45 years had elevated Blood pressure 92.7%
205 and elevated FBS 88.1%. A study done in Mali documented that OR increases with age from
206 2.06 (30–44 years) to 7.25 (60 and more).²² This is similar to finding in other studies in Africa.²³⁻

207 ²⁵ In Ibadan South West Nigeria, a study revealed that hypertension was significantly associated
208 with being in age groups 30-49 years (OR 2.258, 95% CI: 1.311 - 3.884), ≥ 50 years (OR 7.145,
209 95% CI: 3.644 - 14.011).²⁶ In the United States, the estimated percentage of people having
210 diagnosed or undiagnosed diabetes was increasing with age. In the age group of 20-44 years,
211 about 3.7% people had diabetes; in the age group 45-64 years about 13.7%; and age group of \geq
212 65 years about 26.9% had diabetes.²⁷ The study done in Bali showed that the prevalence of raised
213 blood sugar and DM were more than twice higher in the elderly than in the younger age group.²⁸
214 A study done in China documented that Fasting and random blood glucose level rose by 0.15
215 mmol/L, while 2-hour post-prandial blood glucose level rose by 0.26 mmol/L per decade-
216 increase in age.²⁹ Several reports have documented that age is the strongest risk factor for CVD
217 like Hypertension. In the United States, CVD was the leading cause of death for persons 65 years
218 of age and over in 2007. It was responsible for 28% of deaths in this age group.³⁰

219 These findings from current study could possibly be as a result of participants' occupation and
220 residence. Almost all rural dwellers engage in minor farming even if is around their houses
221 Since a high proportion of participants were farmers and traders, trekking long distances to the
222 farm or the farm work itself constituted increased physical activity. Sedentary lifestyle as a a key
223 modifiable factor for both diseases may be rare in these rural communities Age of participants
224 may equally be a factor as these conditions are more with advancing age which is where majority
225 of respondents belong to. The implication of this finding is that since most persons with these
226 conditions do not know that they have the disease, there is the likelihood that they die suddenly
227 with their relatives ascribing it to supernatural things. Then for others that may present at health
228 facility, they are likely to come when complications of their condition has occurred.

229 Recently the numbers of elders are on the increase worldwide with sharp rise in the developing
230 countries. This has impacted on the prevalence of metabolic diseases (impaired fasting glycemia,
231 DM) and Cardio vascular Diseases including Hypertension. This may be as a result of their age,
232 process of aging itself or remotely through several other age-related risk factors. Some of such
233 documented factors include; central obesity, mitochondrial dysfunction, lipid metabolisms
234 disorders, inflammation, β -cell dysfunction, insulin resistance and metabolic syndrome.^{28,31}

235 CONCLUSION

236 Prevalence of raised BP and FBS as well as co-morbid condition was high and in line with
237 findings from other studies. It was more among females and older age. Age was a predictor of
238 both raised BP and FBS. It shows that hypertension, DM and co morbid condition are emerging
239 danger even in rural communities in Nigeria. This calls for interventional programmes including
240 mouth outreaches, persistent health education, enlightenment campaigns and community
241 surveillance programmes to aid reduce this growing burden of the diseases in rural communities.

242 CONFLICT OF INTEREST

243 Authors declare no conflict of interest

244 REFERENCES

- 245
- 246 1. World Health Organization World health report: Global status report on
247 noncommunicable disease 2014. Available
248 from: <http://www.who.int/nmh/publications/ncd-status-report-2014/en/> Accessed on
249 12/2/2019
 - 250 2. Kengne AP, Awah PK, Fezeu L, Mbanya JC. The burden of high blood pressure and
251 related risk factors in urban sub-Saharan Africa: Evidences from Douala in
252 Cameroon. Afr Health Sci. 2007;7(1):38–44
253
254

- 255 3. Isara AR, Okundia PO. The burden of hypertension and diabetes mellitus in rural
256 communities in southern Nigeria. Pan African Medical Journal. 2015;20:103
257
- 258 4. International Diabetes Federation. Diabetes Fact Sheet. 2018. Available at
259 www.idf.org/webdata/. Accessed on 12/2/2019
260
- 261 5. World Health Organization. World health statistics 2012 report. WHO, Geneva, 2012.
262
- 263 6. International Diabetes Federation. Diabetes Fact Sheet. Available at
264 www.idf.org/webdata/docs/background_info_AFR.pdf. Accessed on 12/2/2019
265
- 266 7. Mukadas AO, Misbau U. Incidence and patterns of cardiovascular diseases in north
267 western Nigeria. Niger Med J. 2009; 50(3): 55 - 57
268
- 269 8. Akinkugbe OO Non communicable diseases in Nigeria: national survey (final report) on
270 hypertension, coronary heart disease, diabetes mellitus, G6PD deficiency and anaemia.
271 National Expert Committee on Non-Communicable Disease. Federal Ministry of Health
272 and Social Services, Lagos,
273
- 274 9. Chinenye S, Ogbera AO. Socio-cultural aspects of diabetes mellitus in Nigeria. J Soc
275 Health Diabetes. 2013; 1(1): 15 - 21
276
- 277 10. Omorogiuwa A, Oaikhena GA, Okioya P, Akubueze D, Owobu E, Enahoro I et al
278 Diabetes mellitus: prevalence amongst University staff in southern Nigeria and attitude
279 towards routine glycemc/glycosuric checkup. Int J Biomed & Hlth Sci. 2010; 6(1): 25 -
280 29
281
- 282 11. Nyenwe EA, Odia OJ, Ihekwaba AE, Ojule A, Babatunde S. Type 2 diabetes in adult
283 Nigerians: a study of its prevalence in Port Harcourt, Nigeria. Diabetes Res Clin Pract.
284 2003; 62: 177 - 185
- 285 12. Ekwunife OI, Aguwa CN. A meta analysis of prevalence rate of hypertension in Nigerian
286 populations. Journal of Public Health and Epidemiology. 2011; 3(13): 604 - 607.
287
- 288 13. Ogah OS, Okpechi I, Chukwuonye II, Akinyemi JO, Onwubere BJC, Falase AO, Stewart
289 S, Sliwa K. Blood pressure, prevalence of hypertension and hypertension related
290 complications in Nigerian Africans: A review. World J Cardiol. 2012; 4(12): 327 - 340.
291
- 292 14. Buchan J, Couper I, Tangcharoensathien V, et al. Early implementation of WHO
293 recommendations for the retention of health workers in remote and rural areas. Bull
294 World Health Organ. 2013;91(11):834–840.
295

- 296 15. Ulasi II, Ijoma CK, Onwubere BJ, Arodiwe E, Onodogo O, Okafor C. High prevalence
297 and low awareness of hypertension in a market population in Enugu, Nigeria. *Int J*
298 *Hypertens.* 2011; 869675.
299
- 300 16. Ahaneku GI, Osuji CU, Anisiuba BC, Ikeh VO, Oguejiofor OC, Ahaneku JE. Evaluation
301 of blood pressure and indices of obesity in a typical rural community in eastern Nigeria.
302 *Ann Afr Med.* 2011; 10(2): 120 - 126.
303
- 304 17. Onwubere BJ, Ejim EC, Okafor CI, Emehel A, Mbah AU et al. Pattern of blood pressure
305 indices among residents of a rural community in South East Nigeria. *Int J Hypertens.*
306 2011; 2011: 621074.
307
- 308 18. Sahraki R, Mirshekari M, Sahraki H, Mohammadi AR, Sahraki M, Khazaei Feizabad E.
309 Hypertension Among 30+ Year-Old People in Zahedan (Southeast of Iran). *Shiraz E Med*
310 *J.* 2011;12(3):129–34
311
- 312 19. Sarry El-Din A, Erfan M, Kandeel W, Kamal S, El Banna R, Fouad W. Prevalence of
313 pre-hypertension and hypertension in a sample of egyptian adults and its relation to
314 obesity. *Aust J Basic Appl Sci.* 2012;6(13):481–9.
315
- 316 20. Shapo L, Pomerleau J, McKee M. Epidemiology of hypertension and associated
317 cardiovascular risk factors in a country in transition: a population based survey in Tirana
318 City, Albania. *J Epidemiol Community Health.* 2003;57(9):734–9.
319
- 320 21. Manandhar K, Koju R, Sinha NP, Humagain S. Prevalence and associated risk factors of
321 hypertension among people aged 50 years and more in Banepa Municipality,
322 Nepal. *Kathmandu Univ Med J (KUMJ)* 2012;10(39):35–8.
323
- 324 22. Oumar Bâ, Youssouf Camara, Ichaka Menta, et al., “Hypertension and Associated
325 Factors in Rural and Urban Areas Mali: Data from the STEP 2013 Survey,” *International*
326 *Journal of Hypertension*, vol. 2018, Article ID 6959165, 7 pages,
327 2018. <https://doi.org/10.1155/2018/6959165>.
328
- 329 23. Awoke A, Awoke T, Alemu S, Megabiaw B. Prevalence and associated factors of
330 hypertension among adults in Gondar, Northwest Ethiopia: a community based cross-
331 sectional study. *BMC Cardiovascular Disorders.* 2012;12:113
332
- 333 24. Abebe SM, Berhane Y, Worku A, Getachew A. Prevalence and associated factors of
334 hypertension: a cross-sectional community based study in Northwest Ethiopia. *PLoS*
335 *One.* 2015;10(4):e0125210
336

- 337 25. Gebreselassie KZ, Padyab M. Epidemiology of hypertension stages in two countries in
338 sub-sahara africa: factors associated with hypertension stages. *International Journal of*
339 *Hypertension*, vol. 2015, 12 pages, 2015. <http://dx.doi.org/10.1155/2015/959256>
340
- 341 26. Ajayi IO, Sowemimo IO, Akpa OM, Ossai NE. Prevalence of hypertension and
342 associated factors among residents of Ibadan-North Local Government Area of Nigeria.
343 *Nig J Cardiol* 2016;13:67-75
344
- 345 27. Centers for Disease Control and Prevention. National diabetes fact sheet: national
346 estimates and general information on diabetes and prediabetes in the United States, 2011.
347 Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control
348 and Prevention
349
- 350 28. Suastika K, Dwipayana P, Saraswati IMR, Kuswardhani T, Astika N, Putrawan IB,
351 Matsumoto K, Kajiwaru N, Taniguchi H. Relationship between age and metabolic
352 disorder in the population of Bali. *J Clin Gerontol Geriatrics* 2011, 30: 1-6.
353
- 354 29. Ko GT, Wai HP, Tang JS. Effects of age on plasma glucose levels in non-diabetic Hong
355 Kong Chinese. *Croat Med J.* 2006;47(5):709-13.
356
- 357 30. National Center for Health Statistics. Health, United States, 2010: with special feature on
358 death and dying. Washington DC: US Government Printing Office. Available from:
359 <http://www.cdc.gov/nchs/data/hs/hs10.pdf>. 2011 Accessed on 12/2/2019
360
- 361 31. Suastika K, Dwipayana P, Semadi MS, Kuswardhani TA. Age is an Important Risk
362 Factor for Type 2 Diabetes Mellitus and Cardiovascular Diseases. Available at
363 <http://dx.doi.org/10.5772/52397> Accessed on 12/2/2019
364
365

366