

Phaeophytin and Triterpenoids from *Brachystelma togoense* Schltr, a Nigerian Medicinal Herb

ABSTRACT

The medicinal herb *Brachystelma togoense* Schltr (Apocynaceae) is used traditionally for treatment of ailments. The secondary metabolites, phaeophytin *a*, α -amyirin and lupeol were isolated from the CH₂Cl₂ and MeOH extracts of *Brachystelma togoense*. The structures were elucidated using ¹H, ¹³C and 2D NMR. These phytochemicals have previously being reported to have various biological activities such as anti-inflammatory, anti-fungal and anti-cancer. The presence of phaeophytin *a*, α -amyirin and lupeol in *Brachystelma togoense* justified the use of the plant for medicinal purpose in Nigeria.

Keywords: Secondary metabolites; phaeophytin *a*; α -amyirin; lupeol; *Brachystelma togoense* Schltr

1. INTRODUCTION

Brachystelma was first described by Robert Brown in 1822. The genus *Brachystelma* R. Br. (Apocynaceae: Asclepiadoideae) is represented by about 100-120 species (1). It is an erect perennial herb, growing up to 30 cm high. The genus *Brachystelma* is chiefly distributed in South Africa, South-East Asia and Australasia (2). A total of 18 species are known in India (3) and out of them, 3 species in Maharashtra. *Brachystelma* is found from Ghana to Nigeria, in lowlands to montane areas(4). The raw tuber is said to be edible (4). Many of the tuberous *Brachystelma* are known to be used medicinally for the treatment of headache, stomachache

26 and colds in children(5). *Brachystelma togoense* has being medicinally used for the treatment
27 of dysentery, cough and cold, wounds, stomach ache, typhoid and erectile dysfunction.

28 **2. MATERIAL AND METHOD**

29 **2.1 Collection**

30 The aerial parts of *Brachystelma togoense* was collected during April 2018 from the
31 Ugbokolo forest in Okpokwu, which is the local government area of Benue State-Nigeria.
32 The plant was collect around 10:00 and stored in a plastic container before it was air-dried.
33 The collected specimen was positively identified by Mr. Namadi Sanusi, a botanist at Ahmadu
34 Bello University, Zaria as *Brachystelma togoense*. A specimen (no. 25856) had been retained
35 at the Department of Biological Sciences, Ahmadu Bello University, Zaria-Nigeria (Figure
36 1).

37 **2.2 Extraction and isolation**

38 The air-dried *B. togoense* was manually reduced to powder using mortar and pestil. Exactly
39 1000 g of the powdered plant material was extracted on a shaker at room temperature using
40 100 % dichloromethane (CH_2Cl_2) for 72 h. The extracts were concentrated using a rotary
41 evaporator at 40° C resulting in a brown gum-like texture (32 g). The same procedure was
42 used for methanol (MeOH) which yielded a brown gum-like texture (36 g). The CH_2Cl_2 and
43 MeOH extracts were separated by flash chromatography (Biotage system) over silica gel
44 using three solvents. Firstly, a hexane/ CH_2Cl_2 gradient starting with 100 % hexane and
45 gradually increasing the polarity to 100 % CH_2Cl_2 . Secondly, CH_2Cl_2 /EtOH/Ac from a 100
46 % CH_2Cl_2 to 50 % EtOH/Ac and to 100 % EtOH/Ac to yield various fractions (fr. 1-100).
47 Fr.20 was spotted on the TLC plate using 100 % CH_2Cl_2 and appeared a pure compound **1**
48 (51.0 mg). The same procedure was repeated for the MeOH extract yielding compounds **2**
49 (32.0 mg) and **3** (28.0 mg) which were spotted as pure compounds using CH_2Cl_2 /EtOH/Ac
50 (7:3) from fr.30.

2.2 General experimental procedure

NMR spectra were recorded in CDCl_3 on a 400MHz or 500 MHz Bruker AVANCE III NMR instrument at room temperature. HREIMS were recorded on an Agilent Technologies 6550 iFunnel Q-TOF LC/MS with samples dissolved in CH_2Cl_2 . Infrared spectra were recorded using a Perkin-Elmer (2000 FTIR) spectrometer on NaCl plates.

3. Results and Discussion

The following following compounds phaeophytin *a* (51.0 mg; 0.16 %), α -amyrin (32.0 mg; 0.10 %) and lupeol (28.0 mg; 0.09 %) were isolated from *Brachystelma togoense* using flash chromatography (biotage system). These compounds (Figure 2) were elucidated based on comparison of previous data (6–8).

Phaeophytin-*a* was isolated as a dark green solid from the CH_2Cl_2 extract of the aerial parts of *B. togoense* that was previously described (6). The IR spectrum showed absorbance bands for vinyl proton (3056 cm^{-1}) and sp^3CH ($2987, 2932\text{ cm}^{-1}$) and carbonyl (1736 cm^{-1}) groups. A molecular ion could not be seen in the HRMS spectrometer despite repeated attempts.

From the ^1H and ^{13}C NMR spectra, it was evident that phaeophytin-*a* belonged to the phaeophytin class. This was particularly evident by the downfield shifts at δ_{H} 9.32 s, 9.48 s and 8.56 s which could be assigned as H-5, H-10 and H-20 respectively. The deshielded methyl groups proton resonances occurred at δ_{H} 3.19 (3H-2'), δ_{H} 3.3 (3H-7') and δ_{H} 3.38 (3H-12') and a methoxy group proton resonance occurred at δ_{H} 3.89 (3H-13⁴). The presence of a C-20 phytol tail was evident from the presence of four methyl protons (δ_{H} 0.80 d, $J = 7.3$, δ_{H} 0.82 d, $J = 7.3$, δ_{H} 0.79 s, δ_{H} 1.61 s) and ester carbonyl resonance at δ_{C} 173.8 (C-13³). A comparison of the NMR data of phaeophytin-*a* against literature values for phaeophytin *a* showed the enabled assignment of a keto group carbon resonances at δ_{C} 189.9 to C-13¹ (6,9).

74 Amyrin (α) was isolated as a brown solid from the CH_2Cl_2 extract of the aerial parts
75 of *B. togoense*, which had been isolated previously from the methanol extract of *Sacoglottis*
76 *uchi* (7). The IR spectrum showed absorbance bands for hydroxyl (3055 cm^{-1}) and $\text{sp}^3\text{ CH}$
77 (2987 cm^{-1}) in conjugation and unsymmetrical ethylenic double bond (1733 cm^{-1}) and
78 olefinic carbon (1422 cm^{-1}) groups.

79 The molecular ion was not observed in the HRMS spectrum, however 30 carbons could be
80 counted in the ^{13}C NMR spectrum, indicating the compound was a triterpenoid.

81 The ^1H and ^{13}C NMR spectra (spectrum 3.2 and 3.3) showed the presence of one
82 trisubstituted double bond. A hydroxyl group was placed on C-3 confirmed by the C-3 (δ_{C}
83 79.3) resonance correlating with both the 3H-23 (δ_{H} 0.99 s), 3H-24 (δ_{H} 0.78 s) and H-5 (δ_{H}
84 0.73 d, $J = 11.5$) resonances. A further singlet (δ_{H} 0.79, 0.93, 0.99, 0.78 and 1.24) and two
85 doublet (δ_{H} 0.86 d, $J = 6.2$ and δ_{H} 0.95 d, $J = 6.2$) methyl group proton resonances were
86 present and the typical 12-olaenene double bond (δ_{H} 5.25, δ_{C} 126.1, δ_{C} 138.2) was seen. A
87 comparison against literature data (7) confirmed that this compound was α -amyrin which has
88 been isolated previously from the stem bark of *Sacoglottis uchi* (Humiriaceae)(7).

89 The configuration of the hydroxyl group at C-3 was confirmed as β by the coupling constant
90 of H-3 ($J = 5.1, 11.3$). The configurations at the chiral centres were confirmed using the
91 NOESY spectrum.

92 Lupeol was isolated as a brown solid from the MeOH extract of the aerial parts of *B.*
93 *togoense* which had been isolated previously from the hexane extract of *Magnolia salicifolia*
94 (10) as well as synthesised (8). The IR spectrum showed an absorbance band for hydroxyl
95 (3363 cm^{-1}). The molecular ion was no seen in the HRMS spectrum, however 30 carbons
96 could be counted in the ^{13}C NMR spectrum indicating the compound was a triterpenoid.

The NMR spectra of lupeol showed the presence of an *iso*-propenyl group typical of the lupene-type of pentacyclic triterpenoids. Coupled ²H-29 methylene protons (δ_{H} 4.69 d, $J = 2.1$, δ_{H} 4.57 d, $J = 2.4$) and ¹³C NMR resonances (δ_{C} 105.9, δ_{C} 151.2, δ_{C} 19.5) could be assigned to two H-29 and C-29, C-20 and C-30 respectively (11).

Lupeol was identified as the known 3 β -hydroxylup-20(29)-ene, commonly referred to as lupeol. A literature search revealed that the ¹³C NMR chemical shifts similar to those of lupeol had been reported for lupeol. The configurations at the chiral centres were confirmed using the NOESY spectrum

Previously, pheophytin *a* has been reported to possess antimicrobial activity against *Candida albicans* (ATCC 90028) and *C. albicans* (ATCC 76615) (12) as well as antioxidant activity (13). Amyrin (α) has been reported to exhibit antimicrobial activity against *Escherichia coli*, *Pseudomonas aeruginosa*, *C. albicans*, *Staphylococcus aureus* and *Trichophyton mentagrophytes* (14). Antiprotozoal, anti-inflammatory, antitumor and antimicrobial activity had been reported for lupeol (15).

Conclusion

Phaeophytin *a*, α -amyrin and lupeol are reported here for the first time from *B. togoense*. This was also the first report of the phytochemical quantification in *B. togoense* in Nigeria. However, these secondary metabolites, i.e phaeophytin *a*, α -amyrin and lupeol were reported previously to show various biological activities. Therefore, the results of chemical compound analysis of *B. togoense* justified the ethnomedicinal uses of this plant in Nigeria.

Acknowledgments

The author wishes to thank the Natural Product Research Group, University of Surrey, UK for the opportunity to carry out my research work using their laboratory, Chemicals and Instruments.

Competing Interests

Authors have declared that no competing interests exist.



Figure 1: *Brachystelma togoense* in its natural habitat (16)




Fig.2: Structures of isolated compounds **1-3** from *B.togoense* schtlr

1. Phaeophytin *a*

2. α -Amyrin

3. Lupeol

References

1. Bruyns P V. Three New Species of *Brachystelma* (Apocynaceae, Asclepiadoideae, Ceropegieae) from South Tropical and Southern Africa. Vol. 19. SPIE; 2009. 5 p.
2. Ollerton J, Masinde S, Meve U, Picker M, Whittington A. Fly pollination in *Ceropegia* (Apocynaceae: Asclepiadoideae): biogeographic and phylogenetic perspectives. *Annals of Botany*. 2009;103(9):1501–14.
3. Britto SJ, Bruyns P V. Three new species of *Brachystelma* from Tamil Nadu, India. *Haseltonia*. 2016;(22):48–54.
4. Kew Royal Botanical Gardens. Electronic Plant Information Centre (ePIC) [Internet]. 2019 [cited 2019 Feb 7]. Available from: <http://epic.kew.org/index.htm>
5. Rajaram MS, Rathod J, Dilip V. I S S N 2278 – 4357 Pharmacognostical Studies on the Tuber of *Brachystelma Edulis* Coll . and Helmsl . an Endemic To Peninsular , India . 2014;3(6):1958–65.
6. Schwikkard SL, Mulholland DA, Hutchings A. Phaeophytins from *Tapura fischeri*. *Phytochemistry*. 1998;49(8):2391–4.
7. Abreu VG da C, Corrêa GM, Lagos IA dos S, Silva RR, Alcântara AF de C. Pentacyclic triterpenes and steroids from the stem bark of *uchi* (*Sacoglottis uchi*, Humiriaceae). *Acta Amazonica*. 2013;43:525–8.
8. Cmoch P, Korda A, Rárová L, Oklešťková J, Strnad M, Luboradzki R, et al. Synthesis and structure–activity relationship study of cytotoxic lupane-type 3β-O-monodesmosidic saponins with an extended C-28 side chain. *Tetrahedron* [Internet]. 2014;70(17):2717–30. Available from: <http://www.sciencedirect.com/science/article/pii/S0040402014003123>

- 157 9. Matsuo A, Ono K, Hamasaki K, Nozaki H. Phaeophytins from a cell suspension
158 culture of the liverwort *Plagiochila ovalifolia*. *Phytochemistry* [Internet].
159 1996;42(2):427–30. Available from:
160 <http://www.sciencedirect.com/science/article/pii/0031942296000179>
- 161 10. Silva ATM e, Magalhães CG, Duarte LP, Mussel W da N, Ruiz ALTG, Shiozawa L, et
162 al. Lupeol and its esters: NMR, powder XRD data and in vitro evaluation of cancer
163 cell growth. *Brazilian Journal of Pharmaceutical Sciences* [Internet]. 2018 Feb 1 [cited
164 2019 Feb 19];53(3). Available from:
165 [http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1984-](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1984-82502017000300621&lng=en&tlng=en)
166 [82502017000300621&lng=en&tlng=en](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1984-82502017000300621&lng=en&tlng=en)
- 167 11. Silva ATM e, Magalhães CG, Duarte LP, Mussel W da N, Ruiz ALTG, Shiozawa L, et
168 al. Lupeol and its esters: NMR, powder XRD data and in vitro evaluation of cancer
169 cell growth. *Brazilian Journal of Pharmaceutical Sciences*. 2018 Feb;53(3).
- 170 12. Gomes RA, Teles YCF, Pereira F de O, Rodrigues LA de S, Lima E de O, Agra M de
171 F, et al. Phytoconstituents from *Sidastrum micranthum* (A. St.-Hil.) Fryxell
172 (Malvaceae) and antimicrobial activity of pheophytin a. *Brazilian Journal of*
173 *Pharmaceutical Sciences* [Internet]. 2015;51:861–7. Available from:
174 [http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1984-](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1984-82502015000400861&nrm=iso)
175 [82502015000400861&nrm=iso](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S1984-82502015000400861&nrm=iso)
- 176 13. Kusmita L, Puspitaningrum I, Limantara L. Identification, isolation and antioxidant
177 activity of pheophytin from green tea (*Camellia sinensis* (L.) Kuntze). *Procedia*
178 *Chemistry* [Internet]. 2015;14:232–8. Available from:
179 <https://dx.doi.org/10.1016/j.proche.2015.03.033>
- 180 14. Ragasa CY, Puno MRA, Sengson JMAP, Shen C-C, Rideout JA, Raga DD. Bioactive
181 triterpenes from *Diospyros blancoi*. 2009;23(13):1252–8. Available from:

182 <https://dx.doi.org/10.1080/14786410902951054>

183 15. Gallo M, Miranda bullet, Sarachine J. Biological activities of lupeol. In: International
184 Journal of Biomedical and Pharmaceutical Sciences. 2009. p. 46–66.

185 16. Erpenbach A. West African Plants - A Photo Guide - *Brachystelma togoense* Schltr.
186 [Internet]. 2009 [cited 2019 Feb 18]. Available from:

187 http://www.westafricanplants.senckenberg.de/root/index.php?page_id=14&id=4246

188
189
UNDER PEER REVIEW