

Effect of Tapioca Levels on Production of Swine**ABSTRACT**

There is little definitive information available regarding tapioca's effect on the swine performance and meat quality. Thus, this study was carried out. Thirty-six cross-bred [(Landrace × Yorkshire) × Duroc] growing-finishing swine with their average initial BW of 26.5±2.1 kg was used in this study. The animals were fed with control (no addition of tapioca), treatment 1 (T1 – 10% tapioca) and treatment 2 (T2 – 20% tapioca) for different periods (tapioca as-fed basis). The experimental period lasted for 98d. Carcass characteristics, physicochemical properties, meat composition and sensory test were not significantly different among treatments except for the carcass weight which was increased ($p<0.05$) in the tapioca diet groups. Swine fed with tapioca-replaced diet has no detrimental effects on growth performance or meat quality. Instead, it significantly increased the carcass weight. Therefore, we conclude that tapioca replacement of 20% can aid as alternative feed ingredient of energy source in improving carcass weight for growing-finishing swine.

Keywords: Swine production; growth performance; swine diet; tapioca.

1. INTRODUCTION

Livestock producers are continually looking for new ingredients to include in diets to fulfill specific consumers' demands. Although conventional grains are the most widely used high energy feedstuff, unconventional carbohydrates often provide an alternative. Moreover, a concentrated carbohydrate source provided in a diet with high starch composition may improve the growth rate and carcass traits of pigs [1]. One of these is tapioca, which is a source of starch (62.0%) and has a nutritional value that allows for the partial replacement of cereal grain; this might maximize efficiency for the expected characteristics [2]. Tsudir *et al.* [3] reported that tapioca has dry matter (DM), crude protein (CP), ether extract (EE), crude fibre (CF), nitrogen free extract (NFE) and total ash (TA) of 94.20, 3.30, 0.60, 2.70, 91.10 and 2.30 percentages, respectively. Also, the energy content (ME) of tapioca root in pig was somewhat similar to maize [4, 5, 6]. Tapioca has been used as a livestock feed in some of the countries. It has been included at large scale (multi-millions of tons of feed, annually) without causing (health, production, or meat-quality) problems. However, there is little definitive information available regarding its effect on swine meat characteristics. Zinn and DePeters [7] previously reported that tapioca pellets can be used to replace up to 30% of dry matter intake

35 in growing to finishing diets without adversely affecting the average daily gains of feedlot
36 cattle. Moreover, a 10-25% inclusion level of tapioca feed ingredient in the swine diet was
37 recommended by Moehn *et al.* [2]. However, there is little definitive information available
38 regarding its effect on swine meat characteristics. Thus, the objective of this study was to
39 determine the effect of tapioca as feed ingredient in the diet of growing-finishing swine for growth
40 performance and carcass quality in swine.

41

42 2. MATERIALS AND METHODS

43 2.1 Animals, diets and study design

44 A total of 36 male swine [(Landrace × Yorkshire) × Duroc] with average live weight of
45 26.53±2.10 kg at the beginning of the experiment and 114.13±3.16 kg at the time of slaughter
46 were used in this experiment. Twelve swine were used in each treatment and control group
47 which was represented by three pens with four swine in each pen. The swine were provided
48 balanced diet at 5.5% of BW/d and supplied fresh water throughout the experiment. The
49 feeding was done in phases; grower (20-50kg), early finisher (50-80kg) and late finisher (80-
50 120kg), and tapioca levels were provided at 0% (Control), 10% (T1) and 20% (T2) (Table 1).
51 The composition of the diets and their calculated chemical compositions were prepared in
52 accordance with the National Research Council (NRC) guideline [8]. The animals used in this
53 experiment were cared for in accordance with the guidelines established by National Institute
54 of Animal Science (NIAS), Korea. The research protocol including the procedures for the care
55 and treatment of the animals was reviewed and approved by the Animal Care Committee at
56 the NIAS, Korea.

57

58

59

60

61

62

63

64

65

66

67

68 **Table 1. Ingredient composition** and nutrient content of the experimental diets for
 69 **growing-finishing swine** at different stages (as fed-basis)

Live weight (kg)	Grower (20 ~ 50)			Early finisher (50 ~ 80)			Late finisher (80 ~ 120)		
	Control	Tapioca	Tapioca	Control	Tapioca	Tapioca	Control	Tapioca	Tapioca
		10%	20%		10%	20%		10%	20%
Ingredients, %									
Soybean meal	19.07	22.73	25.66	11.99	16.54	19.18	3.94	9.84	12.49
Corn	68.76	50.73	45.91	76.23	54.65	49.98	68.31	56.28	50.19
Palm meal	-	-	-	-	-	-	-	2.50	5.00
Tapioca	-	10.00	20.00	-	10.00	20.00	-	10.00	20.00
Lupine seed	6.48	-	-	6.36	-	-	8.06	-	-
Wheat grain	-	8.85	1.91	3.3	11.82	5.04	11.63	13.67	5.34
Salt	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30	0.30
Methionine	-	-	0.02	-	-	-	-	0.03	0.06
Lysine	0.19	0.17	0.13	0.14	0.10	0.07	0.17	0.11	0.11
Limestone	0.84	0.84	0.60	0.82	0.77	0.54	0.86	0.86	0.44
Molasses	2.47	2.96	3.00	0.32	3.00	3.00	4.00	3.68	4.00
Dicalcium phosphate	0.77	0.57	0.81	0.54	0.33	0.57	0.23	0.11	0.52
Soybean oil	1.12	3.00	1.81	-	2.49	1.32	2.50	2.62	1.55
Total	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00
Nutrient Content*									
DM, %	89.64	89.65	89.64	89.67	89.66	89.68	89.68	89.68	89.69
CP, %	16.16	15.90	16.00	13.80	13.80	13.80	11.50	11.70	11.50
DE, kcal/kg	3,450	3,450	3,450	3,400	3,400	3,400	3,400	3,400	3,400
CF, %	4.00	4.00	4.00	4.00	4.00	4.00	4.30	4.30	4.30
Ca, %	0.60	0.60	0.60	0.50	0.50	0.50	0.45	0.45	0.45
P, %	0.50	0.50	0.50	0.45	0.45	0.45	0.40	0.40	0.40
Lysine, %	0.95	0.95	0.95	0.75	0.75	0.75	0.60	0.60	0.60
Methionine, %	0.25	0.25	0.25	0.23	0.22	0.22	0.19	0.21	0.21

70 * Calculated values

71 Vit.-Min. premix provided 3.5g per kg of diet containing 1,600,000 IU of vit. A, 300,000 IU

72 of vit D₃, 800 IU of vit E, 132mg of vit K₃, 1,000mg of vit B₂, 1,200 mg of vit. B₁₂, 2,000mg of
73 niacin, 60mg of folic acid, 35,000mg of choline chloride, 800mg of pantothenic calcium,
74 9,000mg of Zn, 12,000mg of Mn, 4,000mg of Fe, 500mg of Cu, 6,000mg of I, and 100mg of Co.

75

76 The experiment was conducted at the Animal Environment Division research farm, NIAS,
77 Suwon, South Korea. According to protocol and management, the swine house had a fully slatted
78 floor pens and an automatic temperature and humidity controller. The average temperature
79 and relative humidity of the house during the experimental period were $20.0 \pm 0.59^{\circ}\text{C}$ and
80 $60.0 \pm 2.8\%$ (mean \pm SD), respectively. The slurry was removed from the pit using a typical
81 gravity drain waste system during experimental periods. The swine were provided with *ad*
82 *libitum* access (5.5% of BW/d) to un-pelleted (except tapioca as pellet) balanced growing
83 swine feed (mash) in 3 daily meals of equal amounts, administered at 0800, 1600, and 2400 h,
84 with a stainless-steel feeder and fresh water supplied by a nipple waterer throughout the
85 experiment. The study was conducted for 14 weeks of experimental period with 7d dietary
86 adaptation. Growth performance such as body weight changes, feed intake and feed
87 conversion ratio were also measured. In addition, carcass characteristics, physicochemical
88 properties, meat composition, color properties and sensory test of pork *longissimus dorsi*
89 muscle at 14th weeks of age were also determined. Three replicates for each of the parameters
90 were used and their averaged data were considered the representative value.

91 2.2 Measurements for growth performance

92 The body weights of the swine were recorded every two weeks from the initial day to the final
93 day of the experiment to calculate the body weight gain (BWG). The feed intake of the swine
94 was recorded every two weeks by offering a weighed quantity of feed and weighing the
95 residual. The feed conversion ratio (FCR) was expressed as gain (G): feed intake (F) of swine.

96 2.3 Meat quality evaluation

97 When the swine reached the average live weight of 114.13 ± 3.16 kg, three swine per pen
98 were randomly selected and transported to a commercial abattoir. They were slaughtered after
99 electrical stunning on the following day and hot carcass weight was measured so that the
100 dressing percentage could be calculated. The dressing percentage for an individual animal was
101 defined as the hot carcass weight divided by the live weight. The carcass and meat quality
102 measurements (obtained from the left side of the carcass) included *longissimus* muscle area,
103 rib eye area, and meat quality grade [9]. Approximately 24 h after slaughter, pH and

104 temperature were determined from the right side of the carcass in the center of the *longissimus*
105 muscle between the 3rd and 4th ribs. A 2.54-cm section of the 9th-rib chop was then removed,
106 and cooking loss and shearing force values were determined as described previously by
107 Kauffman *et al.* [10] and Bee *et al.* [11], respectively. The carcasses were stored in a deep
108 freezer (-18°C) for chemical body analyses. Laboratory analyses of the pork samples were
109 conducted two months after sampling. The samples were thawed at room temperature (20°C),
110 ground, homogenized, and analyzed in triplicate. The preparation of the carcasses for
111 chemical body analyses was conducted by the method developed by Kotarbińska [12]. Meat
112 moisture and ash contents were determined according to AOAC guidelines [13]. Crude
113 protein content in the samples was obtained via the Kjeldahl method [13]. Crude fats were
114 extracted by the Bligh and Dyer method [14] with a chloroform/methanol mixture. Color
115 measurements were taken using a colorimeter (Minolta CM 3500m, Japan). The color
116 readings including lightness (L), redness (a) and yellowness (b) were taken from a
117 *longissimus* section (from the 8th to 10th ribs). The equipment was standardized using a white
118 color standard.

119 **2.4 Sensory evaluation**

120 For the sensory evaluation, meat samples were cooked in an electric grill with double pans
121 (Nova EMG-533, 1,400 W, Evergreen, Korea) to an internal temperature of 75°C. The meat
122 samples (2 × 4 × 1.5 cm) were placed in randomly coded white dishes and served with
123 drinking water. Fifty panel members from the NIAS did the sensory evaluation on the meat. A
124 5-point hedonic scale ranging from 1 (dislike very much) to 5 (like very much) was used to
125 evaluate product attributes (juiciness, tenderness and flavor) in accordance with the guidelines
126 established by Arambawela *et al.* [15].

127 **2.5 Statistical analysis**

128 All the data collected were subjected to one-way ANOVA procedures in a completely
129 randomized design using the general linear model (GLM) procedures (SAS Inst. Inc., Cary,
130 NC) [16]. The growth performance, carcass traits, and pork quality data were compared and
131 significant differences among means of treatment and control groups were separated using
132 Duncan's multiple range (comparison) tests. Variability in the data was expressed as the
133 pooled mean values and standard error (SE) or standard error of the mean (SEM) via the
134 MEANS procedure. The threshold for significance was $p < 0.05$ for all measured variables.

135

136

137

138

139 3. RESULTS AND DISCUSSIONS

140 3.1 Growth performance

141 Several grain sources for swine are available in the market. In spite of that, livestock
142 producers are mostly concerned with choosing carbohydrate-source products are the energy
143 value and cost of the grains. Tapioca is one of these alternative carbohydrate-sources which
144 are more economical. Having somewhat similar energy content (ME) of tapioca root and
145 maize [4, 5, 6] explains unaffected digestible energy (DE) with 3,450 kcal/kg and 3,400
146 kcal/kg in grower and finisher feed formulation as well as other parameters available when we
147 replaced with tapioca in the feed (Table 1).

148 The effects of the experimental dietary treatments on the growth performance, including
149 weight gain, feed intake and feed conversion ratio of the swine are provided in Table 2. The
150 animals remained healthy throughout the duration of the experimental periods and no
151 differences in feed and water intakes were observed between the control and the tapioca-
152 replaced groups. Growth performance was not significantly affected by the treatments. This
153 indicates that replacing corn with tapioca will not affect the growth performance but rather
154 will help the livestock producer in reducing feed cost.

155 **Table 2. Effects of dietary tapioca on the growth performance at 14th weeks of swine¹**

Parameters	Control	Tapioca		SEM ⁴
		10%	20%	
Body weight, kg				
IBW ²	26.5	26.3	26.8	2.10
FBW ³	112.5	115.0	116.9	3.16
Average Daily Gain, kg	0.84	0.87	0.88	0.02
Average Feed Intake, kg/d	2.61	2.73	2.74	0.54
Average Daily Water Intake, L/pig/d ⁵	5.48	5.53	5.59	5.53
Feed conversion ratio	3.11	3.14	3.11	0.09

156 Values presented as Mean; ¹ Individual pig was the experimental unit (n = 12); ² IBW - initial
157 body weight; ³ FBW - final body weight; ⁴ SEM – standard error mean; ⁵ Average daily water
158 intakes during the entire experiment including adaptation and collection periods.

159

160 Although there was not significant difference; between the treatments, swine receiving diets
161 with tapioca tended to show a higher growth performance compared to the control. There was
162 a trend of decreasing final body weight (116.9, 115.0 and 112.5 kg), average daily gain (0.88,
163 0.87 and 0.84 kg), and average feed intake (2.74, 2.73, and 2.61 kg/d) for T2, T1 and control
164 respectively. However, in the study reported by Tsudir *et al.* [3], significantly higher ADG
165 was observed in 50% level of tapioca replacement in feed. The result of our study was
166 different with the result obtained by Tsudir *et al.* [3] due to higher tapioca level was replaced.
167 On the other hand, there was an increase in the intake of feed during the whole experimental
168 period when the grain was replaced with tapioca at different levels which was comparable to
169 the result obtained by Tsudir *et al.* [3]. This indicates that the diet containing tapioca has a
170 high palatability which made it readily accepted by the swine and thus increases in feed intake.

171 3.2 Carcass characteristics and meat quality

172 Indices of carcass quality including carcass characteristics, physicochemical properties, and
173 meat composition are shown in Table 3. The carcass characteristics, (rib eye area, dressing
174 percentage, and meat quality grade), physical properties (shear force, cooking loss, pH,
175 temperature and water holding capacity (WHC)), and meat composition (moisture, fat, protein
176 and ash) were not significantly different except for the carcass weight ($p<0.05$). Moreover,
177 carcass weight showed increasing trend from lowest to highest level of tapioca
178 supplementation ($p<0.05$) with the Control, T1 and T2 recording 85.08, 88.17 and 88.75,
179 respectively. The reason for the increase in carcass weight is unclear. However, Schumacher
180 *et al.* [17] stated that carbohydrates (sucrose) improved carcass weights. Although they
181 employed different carbohydrate ingredients in that study, our results on tapioca replacement
182 were generally consistent with theirs. Even though it was not significant, the increased feed
183 intake and final body weight might be the reason for the significant increase in carcass weight
184 of tapioca-based treatments.

185

186

187

188

189

190

191

192

193

194

195 **Table 3. Effects of dietary tapioca on carcass characteristics, physicochemical properties,**
 196 **and meat composition of pork *longissimus dorsi* muscle at 14th weeks of age**

Parameters	Control	Tapioca		SEM ¹
		10%	20%	
Carcass characteristics				
Rib eye area, cm ²	49.62	50.34	49.53	1.57
Carcass weight, kg	85.08 ^b	88.17 ^a	88.75 ^a	1.54
Dressing percentage, 24-h	73.11	73.95	73.98	0.11
Meat quality grade	1.17	1.17	1.08	0.10
Physicochemical properties of the sirloin				
Shearing force, kg/0.5inch ²	3.89	4.00	3.84	0.08
Oven dry or cooking loss, %	33.44	33.21	32.88	0.40
pH, 24-h	5.58	5.60	5.58	0.02
Temperature, °C, 24-h	3.99	4.01	4.04	0.03
Water holding- capacity, %	53.91	53.27	53.59	0.41
Meat composition, %				
Moisture	72.72	73.22	72.77	0.33
Fat	3.37	3.34	3.38	0.40
Protein	22.32	22.53	22.29	0.15
Ash	0.96	0.99	0.98	0.01

197 ^{a,b} Mean; Means in the same row with different superscripts are significantly different

198 ($p < 0.05$); ¹SEM – standard error of the mean.

199

200 Comparable results were also obtained in physicochemical properties of sirloin and meat
 201 composition of tapioca-replaced and non-replaced treatments (Table 3). The result was in
 202 concordance with Wang *et al.* [18] research wherein meat quality was also not affected by the
 203 treatments of tapioca. The results might be due to comparable CP and DE content of the feed
 204 formulations. As Goerl *et al.* [19] and Witte *et al.* [20] stated, formulating diets based on CP
 205 and energy had no effects on physicochemical properties of muscle such as pH and WBC.
 206 Thus, tapioca-supplementation did not significantly affect the physicochemical properties and
 207 meat composition.

208 **Table 4. Effects of dietary tapioca on organoleptic test of pork *longissimus dorsi* muscle**
 209 **at 14th weeks of age**

Parameters	Control	Tapioca		SEM ¹
		10%	20%	
Color properties of the sirloin				
L	55.17	55.39	55.10	0.76
CIE	a	7.93	7.71	0.26
	b	2.70	2.88	0.31
Hunter	L	48.08	48.30	0.77
	a	6.70	6.52	0.23
	b	2.17	2.32	0.25
Sensory test of pork				
Juiciness	4.53	4.53	4.53	0.15
Tenderness	4.51	4.67	4.53	0.18
Flavor	4.78	4.68	4.68	0.11

210 CIE= International Commission on Illumination; L= lightness; a= Redness; b= Yellowness;
 211 ¹SEM – standard error mean.

212 The color properties (L=lightness, a=redness and b=yellowness) and sensory test (juiciness,
 213 tenderness, flavor) of pork *longissimus dorsi* muscle at 14th week of age are shown in Table 4.
 214 Results were unaffected by the different dietary treatments ($p>0.05$) which is similar to the
 215 results of Goerl *et al.* [19] and Witte *et al.* [20] where the color and sensory properties were
 216 also not affected by their dietary treatments. The results of the present study were also in
 217 concordant with the results of Beech *et al.* [21] and Fernandez *et al.* [22] where no effect was
 218 detected on pork quality carbohydrate (sugar) was added to the diet. This may be due to the
 219 fact that tapioca, which is a type of starch has no strange smell or high fat levels that can
 220 influence carcass characteristics. McKean [23] stated that the desired effect of the tapioca was to
 221 improve weight gain and feed efficiency by improving gut digestion and reducing pathogenic
 222 organism loads.

223

224 4. CONCLUSION

225 The uses of 20% tapioca as feed ingredient improved carcass weight of swine. Thus, tapioca
 226 can be an alternative feed ingredient in growing-finishing swine without any detrimental
 227 effects on growth performance and meat quality.

228 **Ethical disclaimer:**

229 Animal Ethics committee Reference Number is not available with me, but ethical committee's suggestions for
230 management is included. If you suggest, then I may omit. The institute mainly followed EU Ethics.

231 Korea also followed Welfare Quality Assessment Protocol for Pigs (Sows and Piglets, Growing and Finishing
232 Pigs) from

233 [\[https://www.google.co.kr/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&cad=rja&uact=8&ved=0CBsQFjAA&url=http%3A%2F%2Fwww.welfarequalitynetwork.net%2Fdownloadattachment%2F45627%2F21651%2FPig%2520Protocol.pdf&ei=XyXzU6LMFcG48gXyjYC4Dg&usg=AFQjCNG5G_tOnriEP5ZLdOhmEFnXoQ8g5Q\]](https://www.google.co.kr/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&cad=rja&uact=8&ved=0CBsQFjAA&url=http%3A%2F%2Fwww.welfarequalitynetwork.net%2Fdownloadattachment%2F45627%2F21651%2FPig%2520Protocol.pdf&ei=XyXzU6LMFcG48gXyjYC4Dg&usg=AFQjCNG5G_tOnriEP5ZLdOhmEFnXoQ8g5Q)

236

237 REFERENCES

- 238 1. Camp LK, Southern LL, Bidner TD. Effect of carbohydrate source on growth performance,
239 carcass traits, and meat quality of growing-finishing pigs. *J. Anim. Sci.* 2003; 81:2488–2495.
- 240 2. Moehn S, Price JD, Zijlstra RT, Sauer WC, Clark OG, Ball RO, Atakora JKA, Feddes JJ,
241 Zhang Y, Leonard JJ, Edeogu I, Morin B. Climate Change and Managed Ecosystems. Diet
242 Manipulation to Control Odor and Gas Emissions from Swine Production. CRC Press;
243 2006.
- 244 3. Tsudir I, Savino N, Das KC. Effect of replacement of maize with tapioca (*Manihot*
245 *esculenta*) root meal on the performance of growing crossbred pigs. *Livest. Res. Rural*
246 *Dev.* 2012; 24(11):197.
- 247 4. FAO. Feasibility of using cassava versus maize or wheat. In: Cassava for livestock feed in
248 sub-Saharan Africa; 2004.
- 249 5. Muller Z, Chou KC, Nah KC, Tan TK. Study of nutritive value of tapioca in economic
250 rations for growing/finishing pigs in the tropics. Pig & Poultry Research & Training
251 Institute, Singapore (Pigs). UNDP/SF Project SIN 67/505. 1972; R-672:1-35.
- 252 6. Wu JF. Energy value of Cassava for young swine. *J. Anim. Sci.* 1991; 69:1349-1353.
- 253 7. Zinn RA, DePeters EJ. Comparative feeding value of tapioca pellets for feedlot cattle. *J.*
254 *Anim. Sci.* 1991; 69(12):4726-4733.
- 255 8. NRC (National Research Council). Nutrient requirements of swine. 10th revised ed.
256 National Academic Science, Washington, DC, USA; 1998.
- 257 9. NPPC (National Pork Producers Council). Procedures to Evaluate Market Hogs. 3rd
258 ed. Natl. Pork Prod. Council., Des. Moines, IA.; 1991.
- 259 10. Kauffman RG, Eikelenboom G, van der Wal PG, Merkus G, Zaar M. The use of filter
260 paper to estimate drip loss of porcine musculature. *Meat Sci.* 1986; 18(3):191-200.

- 261 11. Bee G, Calderini M, Biolley C, Guex G, Herzog W, Lindemann MD. Changes in the
262 histochemical properties and meat quality traits of porcine muscles during the growing-
263 finishing period as affected by feed restriction, slaughter age, or slaughter weight. *J. Anim.*
264 *Sci.* 2007; 85(4):1030-1045.
- 265 12. Kotarbińska M. The chemical composition of the body in growing pigs. *Rocz. Nauk rol.*
266 1971; B-93-1:129–135.
- 267 13. Cunniff PA. *Official Methods of Analysis of AOAC International*. 6th ed. Association of
268 Official Analysis Chemists, Arlington, USA; 1998.
- 269 14. Bligh EG, Dyer WJ. A rapid method of total lipid extraction and purification. *Can. J.*
270 *Biochem. Physiol.* 1959; 37(8):911-917.
- 271 15. Arambawela WJ, Nielsen HE, Danielsen V, Eggum BO. Effect of replacing barley with
272 tapioca meal at two different levels of feeding on the growth and health of early weaned
273 pigs. *Livest. Prod. Sci.* 1975; 2(3):281-288.
- 274 16. SAS. *Statistical Analysis Systems for Windows*. Release 9.1. SAS Institute Inc., Cary, N.
275 C., USA; 2002.
- 276 17. Schumacher E, Elliott R, McMeniman NP, Griffiths I. Evaluation of raw
277 sugar as an energy source for growing/fattening pigs. 1986; Pages 359-362 in 9th Animal
278 Science Congress of the Asian-Australasian Association of Animal Production Societies.
279 ASAP.
- 280 18. Wang JP, Hong SM, Yan L, Yoo JS, Lee JH, Jang HD, Kim HJ, Kim IH. Effects of single
281 or carbohydrases cocktail in low-nutrient-density diets on growth performance, nutrient
282 digestibility, blood characteristics, and carcass traits in growing-finishing pigs. *Livest. Sci.*
283 2009; 126(1-3):215-220.
- 284 19. Goerl KF, Eilert SJ, Mandigo RW, Chen HY, Miller PS. Pork characteristics as affected
285 by two populations of swine and six crude protein levels. *J. Anim. Sci.*
286 1995; 73(12):3621-3626.
- 287 20. Witte DP, Ellis M, McKeith FK, Wilson ER. Effect of dietary lysine level and
288 environmental temperature during the finishing phase on the intramuscular fat content of
289 pork. *J. Anim. Sci.* 2000; 78(5):1272-1276.
- 290 21. Beech SA, Elliott R, Batterha ES. Sucrose as an energy source for growing pigs: a
291 comparison of the effects of sucrose, starch and glucose on energy and protein retention.
292 *Anim. Sci.* 1991; 53(03):383-393.

- 293 22. Fernandez X, Tornberg E, Mågård M, Göransson L. Effect of feeding a high level of
294 sugar in the diet for the last 12 days before slaughter on muscle glycolytic potential and
295 meat quality traits in pigs. *J. Sci. Food Agr.* 1992; 60(1):135-138.
- 296 23. McKean J. Feed additives for swine. 2004; Pages 1-8. Pork information gateway.
297
298