

Chemical and minerals composition of dried ruminal fluid with spray drying method with various hydrocolloids

ABSTRACT

This study was conducted to determine the chemical and **minerals** composition of dried ruminal fluid with spray drying method with various hydrocolloids. Rumen contents were taken from the slaughterhouse and powdered by spray drying with different hydrocolloids including sodium alginate (RA), guar gum (RG), chitosan (RC) and maltodextrin (RM) in two ratios (0.5 and 1% (w/v)). Samples were analyzed with a completely randomized design. The results of mean percentages showed the samples contained moisture (617 ± 0.60 to 7.29 ± 0.03 %), crude protein (9.96 ± 0.06 to 13.02 ± 0.04 %), ether extract (1.82 ± 0.11 to 3.43 ± 0.13 %), Ash (16.25 ± 0.21 to 22.67 ± 0.15 %), NDF (24.18 ± 0.12 to 27.90 ± 0.10 %), and carbohydrates (36.84 ± 0.05 to 44.54 ± 0.56 %). The **minerals** were sodium (1.98 ± 0.12 to 4.02 ± 0.17 %), potassium (0.46 ± 0.07 to 0.76 ± 0.10 %), magnesium (0.69 ± 0.05 to 1.02 ± 0.07 %), calcium (0.33 ± 0.02 to 0.70 ± 0.05 %) and phosphorous (1.22 ± 0.07 to 1.87 ± 0.01 %). The results indicated that dried ruminal fluid with spray drying method with various hydrocolloids has nutrition values that could provide nutrients and minerals for livestock. It is recommended in the diet of animals.

KEY WORDS: Proximate composition, Ruminal fluid, Spray drying, Hydrocolloids, Mineral

1. INTRODUCTION

The global shortage of animal feed increases the share of livestock feed costs and the income generated by the production of livestock products has been affected. In order to compensate for this shortage, the utilization and optimal use of agricultural waste and agricultural products as feed in ruminants is inevitable to improve livestock production [1]. Large amounts of rumen content are produced daily as a byproduct in slaughterhouses [2]. These pollutants pollute the environment by releasing toxic gases, including methane, nitrogen dioxide and other gases in the atmosphere [3]. Also, the rumen fluid contains high levels of ammonia and phosphorus [4]. Therefore, rumen fluid is a serious environmental contaminant in most slaughterhouses in developing countries [5]. The contents of the rumen are rich in raw microbial protein and contains forage materials digested in different stages of fermentation, saliva, amino acids, group B vitamins and end products of ruminal fermentation [6], rich in minerals [7]. As well as antiphenological factor [5]. Sun-dried rumen digesta has been reported to contain 11.18 % crude protein, 22.99 % crude fiber, 1.22 % ether extract, 21.54 % ash, 0.20 % calcium and 0.45 % phosphorous [8]. Sakaba *et al.* [9] also reported the composition of rumen digesta from sheep slaughtered in Zuru Abattoir was 5.83 % moisture, 15.52 % crude protein, 5.17 % lipids, 48.17 % fiber, 11.00 % ash, 19.98 % soluble carbohydrate, 19.98 % Sodium, 4.37 % Potassium, 0.42 % magnesium, 0.45 % calcium and 4.37 % phosphorous from sun-dried rumen digesta. **The benefits of recycling ruminal liquid as slaughter waste reduces environmental pollution, also, it is a source of feed for ruminants** [3]. With proper processing, the rumen content can provide a large supply of nutrients when it enters the ration [10]. And the protein needs of high-producing animals [11]. Drying can be used as a solvent reduction, which is usually water, to produce a stable and saveable product [12]. Spray drying is one of the encapsulation methods used to dry the heat-sensitive materials and a simple, quick, and economical technique for

obtaining a powder from a solution or a liquid suspension (such as an enzyme suspension) [13]. Some Binding agents, such as disaccharides, are added to the protein and improve sustainability during the build and save process. Many studies have shown the protective effects of connective materials for denaturation [14]. Therefore, some carbohydrates (such as starch, maltodextrin and dextrose), gums (such as Arabic gum, acacia gum, sodium alginate and carrageenans), proteins (such as milk or whey protein, gelatin) and chitosan are used in spray drying methods [15]. So far, no research has been done on the approximate compounds of dried ruminal fluid by spray drying method with different hydrocolloids. This study was conducted to determine the composition of this substances.

2. MATERIALS AND METHODS

2.1 Sample preparation and spray drying

The rumen contents were taken from the slaughterhouse and was transferred to a lab inside a pre-warmed containers. Then, the ruminal fluid containing digestive material was mixed with a blender for 2 minutes while injected with carbon dioxide gas. In order to separate rumen fluid from rumen solid materials filtration through four-layers cheesecloth was used. Then, The collected rumen fluid were spray dried by using different hydrocolloids including sodium alginate (Sigma-Aldrich, CAS Number, 9005-38-3, chemical Book), guar gum (Sigma-Aldrich, CAS Number, 9003-30-0), chitosan (Sigma-Aldrich, CAS Number, 9012-76-4, Tokyo chemical Industry Co. Ltd), and maltodextrin (Sigma-Aldrich, CAS Number 9050-36-6, SCBT-Santa Cruz Biotechnology) in two ratios 0.5 and 1% (w/v).The experimental groups were: **1) spray dried rumen fluid with 0.5% maltodextrin (RM0.5), 2) spray dried rumen fluid with 1% maltodextrin (RM1), 3) spray dried rumen fluid with 0.5% chitosan (RC0.5), 4) spray dried rumen fluid with 1% chitosan (RC1), 5) spray dried rumen fluid with 0.5% guar gum (RG0.5), 6) spray dried rumen fluid with 1% guar gum (RG1), 7) spray dried rumen fluid with 0.5% alginate (RA0.5), 8) spray dried rumen fluid with 1% alginate (RA1), 9) spray dried rumen fluid with no hydrocolloid (RN).** A laboratory scale spray dryer (Armfield Mini Spray Dryer, England) with two-fluid nozzles (inner diameter: 0.5 mm) was used for spray drying. The system was operated in a co-current manner with an inlet and outlet air temperature of 120 °C and 50 °C, respectively. Feed rate change in 240e640 mL/h was necessary to achieve a constant outlet temperature. A spray flow rate of 500 mL/h was used and the aspiration rate was 70%. The powdered samples were stored in two layers polyester bags and kept in the refrigerator (5 °C) until analysis.

2.2 Chemical analysis

The samples of dried ruminal fluid with spray drying method with various hydrocolloids analysed for dry matte, ash, ether extract and crud protein according to AOAC (16). The nitrogen-free extract was calculated by reducing samples % ash, % NDF, % Ether Extract, % crude protein, crude fat from 100 to the following formula.

$$\text{NFE} = 100 - (\% \text{ash} + \% \text{NDF} + \% \text{Ether Extract} + \% \text{crude protein}).$$

2.3 Determination of minerals

The minerals of samples were determined according to AOAC [16]. Sodium and potassium were determined by photometric method using FP 640, Jeumeay and phosphorous was measured through Vonado molybdate Yellow method using spectrometer Jenway 1315 UK (UV-visible). The amount of calcium and magnesium were determined using Atomic Absorption Spectrometer (Buck 210, AAS)

2.4 Data collection and statistical analysis

Data were analyzed by the GLM procedure of SAS (17) according to the randomized complete design as following model: $Y_{ij} = \mu + T_i + e_{ij}$, where y was the dependent variable, μ was overall mean, T_i was the effect of of spray drying method with different hydrocolloids. Comparison between different hydrocolloids added was done Tukey test and the significant differences were declared at $P < 0.05$.

Table 1. Approximate analysis of ruminal liquid dried with the spray drying method to various hydrocolloids

Item	Moistre (%)	CP (%)	EE (%)	Ash (%)	NDF (%)	NFE (%)
RM0.5	6.6±0.10 ^{bc}	12.54±0.03 ^d	2.94±0.18 ^b	20.20±0.75 ^c	24.18±0.12 ^f	39.26±0.23 ^d
RM1	6.23±0.17 ^{cd}	12.70±0.06 ^c	2.59±0.07 ^d	21.89±0.28 ^b	24.45±0.18 ^f	38.48±0.26 ^d
RC0.5	6.29±0.03 ^{cd}	13.02±0.04 ^a	2.80±0.07 ^{bc}	17.33±0.32 ^d	27.29±0.26 ^b	39.43±0.09 ^{cd}
RC1	6.17±0.60 ^d	12.57±0.05 ^d	3.43±0.13 ^a	16.85±0.33 ^e	27.90±0.10 ^a	36.84±0.05 ^e
RG0.5	6.80±0.05 ^b	10.51±0.07 ^e	2.28±0.12 ^e	22.67±0.15 ^a	26.58±0.33 ^c	38.42±0.27 ^d
RG1	6.24±0.05 ^{cd}	9.96±0.09 ^f	2.61±0.15 ^{cd}	20.15±0.40 ^c	27.51±0.17 ^b	40.30±0.63 ^c
RA0.5	6.80±0.05 ^b	12.92±0.20 ^b	2.77±0.07 ^{bcd}	16.66±0.25 ^e	24.46±0.39 ^f	42.37±1.02 ^b
RA1	7.29±0.03 ^a	12.30±0.12 ^d	1.82±0.11 ^f	16.25±0.21 ^e	25.45±0.16 ^d	44.54±0.56 ^a
RN	6.25±0.06 ^{cd}	10.6±0.17 ^e	2.83±0.09 ^{bc}	20.14±0.09 ^c	25.02±0.11 ^e	44.14±0.92 ^a
P-value	<.0001	<.0001	<.0001	<.0001	<.0001	<.0001
SEM	0.08	0.22	0.01	0.11	0.07	0.19

Spray dried rumen fluid with 0.5% (RM0.5) and 1% (RM1) maltodextrin, with 0.5% (RC0.5) and 1% (RC1) chitosan, with 0.5% (RG0.5) and 1% (RG1) guar gum, with 0.5% (RA0.5) and 1% (RA1) sodium alginate, and with no hydrocolloids added (RN).

The Chemical analysis % are given as mean ± slandered error.

Different superscripts indicate significant differences in a column.

Significance level was considered at $P < 0.05$.

Table 2. Mineral composition of ruminal liquid dried with the spray drying method with various hydrocolloids

Item	Sodium (%)	Potassium(%)	Calcium(%)	Magnesium(%)	Phosphorous (%)
RM0.5	2.97±0.20 ^b	0.75±0.10 ^a	0.70±0.05 ^a	0.99±0.10 ^{ab}	1.50±0.05 ^c
RM1	2.87±0.19 ^{bc}	0.63±0.10 ^{abcd}	0.58±0.06 ^b	1.02±0.07 ^a	1.65±0.04 ^b
RC0.5	2.52±0.10 ^d	0.65±0.11 ^{abcd}	0.53±0.03 ^b	0.94±0.05 ^{abc}	1.38±0.11 ^d
RC1	2.61±0.25 ^{cd}	0.68±0.11 ^{ab}	0.56±0.08 ^b	0.87±0.07 ^{de}	1.22±0.07 ^e
RG0.5	2.18±0.17 ^{fc}	0.46±0.07 ^d	0.40±0.05 ^c	0.69±0.05 ^e	1.36±0.06 ^d
RG1	1.98±0.12 ^f	0.48±0.08 ^{cd}	0.56±0.12 ^b	0.88±0.06 ^{bcd}	1.65±0.06 ^b

RA0.5	3.80±0.13 ^a	0.53±0.05 ^{bcd}	0.33±0.02 ^c	0.91±0.07 ^{abcd}	1.87±0.01 ^a
RA1	4.02±0.17 ^a	0.65±0.10 ^{abc}	0.35±0.04 ^c	0.85±0.11 ^{cd}	1.85±0.06 ^a
RN	2.41±0.24 ^{de}	0.64±0.09 ^{abcd}	0.55±0.05 ^b	0.96±0.02 ^{abc}	1.33±0.04 ^d
P-value	<.0001	<.0001	<.0001	<.0001	<.0001
SEM	0.04	0.01	0.00	0.00	0.00

Spray dried rumen fluid with 0.5% (RM0.5) and 1% (RM1) maltodextrin, with 0.5% (RC0.5) and 1% (RC1) chitosan, with 0.5% (RG0.5) and 1% (RG1) guar gum, with 0.5% (RA0.5) and 1% (RA1) sodium alginate, and with no hydrocolloids added (RN).

The mineral % are given as mean ± standard error.

Different superscripts indicate significant differences in a column.

Significance level was considered at $P < 0.05$.

3. RESULTS AND DISCUSSION

The amount of moisture has a range of $6.17 \pm 0.60 \%$ to $7.29 \pm 0.03 \%$ and is the highest in RA1 and the lowest in RC1 (Table 1). The values for moisture are lower than 14.64 % and 17.48 % reported by Gebrehawariat *et al.* [8] and Agbabiaka, Madubuike and Amadi [18]. The variation in the moisture content of the samples in this study could be attributed to the processing method. Because in this method, rumen fluid samples are dried with spray drying method samples can be exposed to extreme temperatures for a few seconds [19], which can lead to a decrease in moisture. The spray drying method produces a stable and storeable product that can be stored for a long time [19]. Among the tested hydrocolloids sodium alginate was more successful in reducing the ruminal fluid moisture content. The crude protein content of this material ranged from $9.98 \pm 0.09 \%$ to $13.02 \pm 0.04 \%$ and the highest was RC0.5, and the lowest was RG1 (Table 1). These values were lower than the 18.58 percent reported by Agbabiaka, Anukam and Nwachukwu [20] and in the study of Gebrehawariat *et al.* [8]. was 11.80 and in Sakaba *et al.* [21] study, it was reported at 15.35 %. spray drying technique may alter the 3D protein structure similar to denaturation [22]. The cutting forces that occur in the nozzle of the device also cause the protein to be denatured [23]. Protective effects of hydrocolloids have been reported in relation to the denaturation of heat [24] and the change in the 3D structure of the protein decreases in the presence of compounds such as polysaccharides, proteins or salts [22]. 0.5% chitosan during the spray drying method maintained the percentage of rumen fluid protein. Also the variation in the amount of crude protein is related to the variety and quality of forage consumed by the animal and the soil from which the forage is grown. It can be due to the population and activity of microorganisms during the pre-killing period [25,26]. And the ruminal liquid dried with spray drying with this percentage of the protein recommended in the ration of the animal.

The amount of Ether Extract has a range of $1.82 \pm 0.11 \%$ to $3.43 \pm 0.13 \%$ and the highest was RC1, and the lowest was RA1 (Table 1). These values were higher than 1.22 % percent and 1.69 % percent reported by Gebrehawariat *et al.* [8] and Agbabiaka, Madubuike and Amadi [18]. The variation in lipid is also attributed to the variety, stage of maturity of the plants used in the ration animal, drying method and hydrocolloid used. The amount of $16.25 \pm 0.21 \%$ for RA1 and $22.67 \pm 0.15 \%$ for RG0.5 for ash (Table 1) in this study was less than the 48.73 % reported by Sakaba *et al.* [9]. And the amount of ash in the study by Gebrehawariat *et al.* [8]. was 21.47%. There is no statistically significant difference in the least amount between RA1, RA0.05 and RC1. The NDF content has a range of $27.90 \pm 0.10 \%$ to $24.18 \pm 0.12 \%$. And its highest value in RC1 and its lowest value in RM0.5. There is no

statistically significant difference between RM0.05, RA1 and RA 0.05 (Table 1). These values were greater than 11.00 %, 18.44 % and 22.99 % reported by Sakaba *et al.* [9]. Sakaba *et al.* [21]. and Gebrehawariat *et al.* [8]. And less than of the 34.91 percent reported by Agbabiaka, Madubuike and Amadi [18]. In this study, the chitosan itself has been used as a fiber and has been effective in increasing NDF content. Also, the variation in the amount of NDF depends on the stage of maturity and the degree of fibrillation of the forage used in the livestock. The amount of carbohydrates ($36.84 \pm 0.05 \%$ to $44.54 \pm 0.56 \%$) in this study (Table 1) was more than 19.98 reported by Sakaba *et al.* [9]. And with amount 40.8 of and 38.13 reported by Dairo *et al.* [24], Agbabiaka Madubuike and Amadi [18]. almost equal. This is attributed to the sparing effect of carbohydrate. The chemical analysis of ruminal liquid dried with spray drying method and various hydrocolloids have shown that these materials are active after spray drying and can be used as an additive in the diet of animals. Compared to the drying method in the sun, this method does not reduce the chemical composition of these contents.

As shown in Table 2 the amount of sodium has a range of $1.98 \pm 0.12 \%$ to $4.20 \pm 0.17 \%$ percent in this study, with the highest value of RA1 being the lowest in RG1. There is no statistically significant difference between RA1 and RA0.05. That this amount of sodium was less than 19.98 percent reported by Sankaba *et al.* [9]. In the Sakaba *et al.* [9]. study was 2.46 percent. In this study, hydrocolloid sodium alginate has a sodium element and has been effective in increasing sodium content. The amount of potassium in this study ranged from $0.46 \pm 0.07 \%$ to $0.75 \pm 0.10 \%$. The amount of potassium in the Sakaba *et al.* [9]. study was 0.64 %. The amount of calcium had a range of $0.33 \pm 0.02 \%$ to $0.70 \pm 0.05 \%$ in this study. The amount of magnesium was $0.69 \pm 0.05 \%$ to $1.02 \pm 0.07 \%$ and the phosphorus content was $1.22 \pm 0.07 \%$ to $1.87 \pm 0.01 \%$ percent. This study showed that the ruminal fluid dried with spray drying method with different hydrocolloids had minerals.

4. CONCLUSION

The ruminal fluid, in addition to microorganisms, vitamins containing microbial protein, enzymes and Minerals. By using a drying method, it can be converted to a dry, storageable, portable material. The values for chemical and minerals composition obtained from the dried ruminal fluid with spray drying method with various hydrocolloids indicated the ability of the material to provide the nutrients particularly protein required by animals for normal physiological activities. The material is however deficient mineral for livestock nutrition; Therefore, supplementation is required.

RECOMMENDATIONS

From the above conclusion, farmers can use the dried ruminal fluid with spray drying method with various hydrocolloids for livestock feeding trials. More research is needed on the use of these materials in domestic animals.

Ethical Approval:

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

Consent: NA

REFERENCES

1. Negesse T, Patra AK, Dawson LJ, Tolera A, Merkel RC, Sahlu T Goetsch AL. Performance of spanish and boerxspanish doelings consuming diets with different levels of broiler litter. *Small Ruminant Research*. 2007; 69(1-3):187-197.
2. Said IF, Elkhair RMA, Shawky SM, Abdelrahman HA, Elfeki MA. Impact of feeding dried rumen content and olive pulp with or without enzymes on growth performance, carcass characteristics and some blood parameters of Molar ducks. *International Journal Agricultural Reserch*. 2015; 4:2319-2473.
3. Mondal S, Haldar S, Samanta I, Samanta G, Ghosh TK. Exploring nutritive potential of undigested rumen contents as an ingredient in feeding of goats. *Animal Nutrition and Feed Technology (India)*. 2013;13:79-88.
4. Tritt WP, Schuchardt F. Materials flow and possibilities of treating liquid and solid wastes from slaughterhouses in germany. *A review Bioresource Technology*.1992; 41(3): 235-245.
5. Cherdthong A, Wanapat M. Manipulation of in vitro ruminal fermentatationand digestibility by dried rumen digesta. *Livestock Science*. 2013; 153(1-3): 49-100.
6. Cherdthong A, Wanapat M, Seankamorsn A, Sopapong C, Anantasook N, Gunun p. Improving rumen ecology and microbial population by dried rumen digesta in beef cattle. *Tropical animal health and production*. 2015; 47(5): 921–926.
7. Elfaki MOA, Abdelatti KA. Nutritive evaluation of rumen content from cattle, camel, sheep and goats. *Global Journal of Animal Scientific Research*. 2015; 3(3): 617-621.
8. Gebrehawariat E, Animut G, Urge M, Mekasha Y. Sun Dried Bovine Rumen Content (SBRC) as an ingredient of a ration for white leghorn layers. *East African Journal of Science*. 2016;10(1):29-40.
9. Sakaba AM, Hassan AU, Harande IS, Isgogo MS, Maiyama FA, Danbare BM. Proximate composition of rumen digesta from sheep slaughtered in Zuru Abattoir, Kebbi State, Nigeria. *Journal of Agriultural Science and Practice*. 2017;2: 86-89.
10. Okpanachi U, Aribido SO, Daikwo IS. Growth and haematolo- gical response of growing rabbits to diets containing graded levels of sun dried bovine rumencontent. *African Journal of food, agriculture, nutrition and development*. 2010;10(11), 4444-4457.
11. Boorman KN, Ellis GM. Maximum nutritional response to poor quality and amino acid utilization. *British poultry science*, 1996; 37(1): 145-156.
12. Namaldi A, Calik P, Uludag Y. Effects of spray drying temperature and additives on the stability of serine alkaline protease powders. *Drying Technology*.2006; 24(11): 1495–1500.
13. Bajsic I, Kranjcevic E. Automation of industrial spray dryer. *Instrumentation Science and Technology*. 2001; 29(1):41-52.
14. Jalalipour M, Gilani K, Tajerzadeh H, Najafabadi AR, Barghi M. Characterization and aerodynamic evaluation of spray dried recombinant human growth hormone using protein stabilizing agents. *International journal of pharmaceutics* .2008; 352(1-2), 209-216.
15. Gharsallaoui A, Roudaut G, Chmbin O, Voilley A, Saurel R. Applications of spray-drying in microencapsulation of food ingredients: an overview. *Food research international*. 2007; 40(9):1107-1121.
16. AOAC. Official methods of analysis 18th ed. Association of Official Analytical Chemist, Washington, DC; 2005.
17. SAS Institute, SAS User's Guide: Statistics, Release 9.1. SAS Institute. Inc, Cary, NC: 2003.
18. Agbabiaka LA, Madubuike FN, Amadi SA. Studies on nutrients and anti-nurients of rumen digesta from three most domesticated ruminants in Nigeria. *Pakistan journal of Nutrition*. 2012; 11(7): 580-582.
19. Namaldi A, Calik P, Uludag Y.. Effects of spray drying temperature and additives on the stability of serine alkaline protease powders. *Drying Technology*. 2006; 24(11): 1495–1500.
20. Agbabiaka LA, Anukam KU, Nwachukw VN. Nutritive value of dried rumen digesta as replacement for soybean in diets of Nile Tilapia (*Oreochromis niloticus*) fingerlings. *Pakistan journal of Nutrition*. 2011; 10(6): 568-571.
21. Sakaba AM, Isgogo SM, Hassan AU, Sanchi UZ. Proximate composition of sun-dried rumen digesta from goat slaughtered in zuru central slaughterhouse,kebbi state, Nigeria. *Asian Journal Research in Animal Veterinary Sciences*. 2018;1(3):1-5.
22. Alloue WA, Destain MJ, Amighi K, Thonart P. Storage of yarrowia lipolytica lipase after spray-drying in the presence of additives. *Process Biochemistry*. 2007; 42(9): 1357-1361.

23. Gharsallaoui A, Roudaut G, Chambin O, Voilley A, Saurel R. Applications of spray-drying in microencapsulation of food ingredients: an overview. *Food Research International*, 2007;40(9):1107-1121.
24. Schüle S, Schulz-Fademrecht T, Garidel P, Bechtold-Peters K, Frieß W. Stabilization of IgG1 in spray-dried powders for inhalation. *European Journal of Pharmaceutics and Biopharmaceutics*. 2008;69(3): 793-807.
25. Dairo FA, Aina SOO, Asafa AR. Performance evaluation of growing rabbits fed varying levels of rumen content and blood rumen content mixture. *Nigerian Journal of Animal Production*. 2005; 32: 67-72.
26. Abouheif MA, Kraidees MS, Al-Sebood BA. The utilization of rumen content-barley meal in diets of growing lambs. *Asian-Australian Journal of Animal Science*. 1999; 12(8): 1234-1240.