

Adoption **Level of Agroforestry Practices in Katsina State, Nigeria**

ABSTRACT

Unsustainable forest land use practices have resulted in land degradation in Nigeria leading to low crop yield. Agroforestry is a viable option for reversing dwindling crop yields through proper soil management practices. There is notably no sufficient published information on agroforestry practices (AP) in Katsina State. The aim of this study was to assess AP for sustainable land use in the study area. Multistage stratified sampling design was used to select respondents from the study. Three Local Government Areas (LGAs) were randomly selected from each of the agro-ecological zones (Sahel, Sudan and Guinea) of Katsina State. Within each of the selected LGAs, one community was randomly selected and forty respondents were randomly sampled from each community. Using structured questionnaire, information was sought on socio-economic and AP. Data were analysed using descriptive statistics. Based on the results, multipurpose trees on farmland (79%), windbreaks (50%), woodlots (49.7%), improved fallow in shifting cultivation (32%) and home gardens (24.7%) were the common AP in the study area. Benefits of AP in the area included preservation of the environment (98.5%), provision of fruits and leaves (98%), and improvement of soil fertility (98%), erosion control (98%) and improvement of farmers' income (96%). Agroforestry practices enrich the soil with important nutrients and prevent soil erosion. The adoption of multipurpose trees on farmland in the study area will help in preventing environmental degradation, desertification and enhance food crop production.

Keywords: Degradation, Forest, Nutrients, Savanna, Sustainable, Soil.

INTRODUCTION

Cropping system and anthropogenic activities are the major factors affecting sustainable land use in most developing countries. The result is continuous stress on the natural resources base with the conversion of forested areas into croplands, the cropping of marginal lands and the use of adverse agricultural methods including inappropriate and excessive application of agro-chemicals. The need for more food has led to increased deforestation, shortened fallow periods in shifting cultivation cycles, and set in motion a degradation spiral, leading to reduced productive capacity of the land and decreased crop yields. In addition, indiscriminate fire wood gathering, timber harvesting and grazing have aggravated land degradation in many parts of Nigeria.

A best and easy method for replenishing soil nutrients would be use of inorganic fertilizers; these are beyond most of the rural farmers' budgets. Thus agroforestry practices offer an alternative solution to resource-constrained smallholder farmers, who in the absence of inorganic fertilizers would otherwise grow crops without addressing nutrient requirements and harvest little or nothing for storage (Jamala, et

32 *al.*, 2013). The presence of woody perennials in agroforestry systems may affect several bio physical and
33 bio-chemical processes that determine the health of the soil substrate (Nair, 1993).

34 In Katsina state, in the Sudan savanna ecological zone of Nigeria, adoption level of agroforestry practices
35 is low despite the recognized potential of sustainable agroforestry to contribute to more resilient farming
36 systems, food security and poverty reduction. Although factors affecting adoption of agroforestry have
37 been carried out in different part of Nigeria (Akpabio *et al.*, 2008), there is dearth information on the
38 adoption level of Agroforestry in the study area. Socio-economic study of farmers and their relationship to
39 the agroforestry would help to ascertain the opportunities for the development of agroforestry systems in
40 Katsina, Nigeria (Maluki *et al.* 2016). Studies have revealed that growing of trees is a function of socio-
41 economic characteristics of the farming community (Irshad *et al.*, 2011).

42 Population pressure today precludes practices that would enable the environment to recover from
43 extensive cropping systems or anthropogenic activities. It is expected that this study will assist
44 agricultural planners and policy makers to properly address the problem of environmental and soil
45 degradation. This paper is aimed at identifying the various agroforestry practices adopted by farmers; and
46 evaluate farmers' use of agroforestry practices in the study area with a view to ascertaining the benefits
47 derived from planting trees along with food crops in the study area.

48 **MATERIALS AND METHODS**

49 **The Study Area**

50 Katsina State lies between latitude 12° N and longitude 8° E. The state is located in the upper Sudan and
51 lower Sahel regions of Nigeria. The entire landmass of Katsina State is composed of basement complex of
52 pre- Cambrian era, which consists of meta-sediments that have been transformed into anatectic
53 migmatites and granites. The hot, dry and dust laden North East Trade wind predominates in this area for
54 as long as 7 months of the year. Rainfall is experienced in the state from June to September; with mean
55 annual rainfall from 1016 mm to 1143mm in the south and less than 635 mm in the Northern part of the
56 state. The state on the whole has a mean annual rainfall of about 840 mm. Mean relative humidity is lower
57 than 50 % in January and February and could be as high as 80 % in June and July. Temperature range is
58 often from 38° to 41°C.

59 **Data collection and analysis**

60 Multi-stage and stratified random sampling techniques were used for the study. Nine Local Government
61 Areas (LGAs) (three per each agro-ecological zone) were randomly selected out of the 34 LGAs in

62 Katsina state. Accurate data on the actual population of the rural farmers in Katsina state were not readily
63 available; equal number of farmers (120 farmers) was randomly sampled from each of the agro-ecological
64 zones through the use of random numbers. This gave a sample size of 360. Questionnaire was used to
65 collect information on the study objectives. The questionnaire was sub-divided into: source of farm land,
66 farm location and size; other agricultural activities, types of crops and trees planted and estimated farm
67 yield, estimated agricultural income, source of information and the perceived benefits of AP. Information
68 was also obtained through personal observation.

69 The data obtained from the study were collated and analysed. Descriptive statistic (such as frequencies
70 and percentages) was used to describe the variables; and inferential statistic (such as ANOVA, chi-square
71 and correlation analysis) was also used to test for significant variation within the study area.

72 **RESULTS**

73 **Demographic Attributes of the Respondents in the Study Area**

74 The demographic characteristics of the respondents in the study area showed the males were more
75 involved in agroforestry practices than the females. Most of the respondents in the Sahel and Sudan
76 Savanna were between 30 and 49 years; while in the Guinea savanna, majority of the respondents age
77 ranged from 60 years and above. The age distribution is an important factor in farming activities because
78 it affects the work force and decision-making in farming activities. The dominant age bracket among rural
79 dwellers (30 – 49 years) in Sahel and Sudan and ≥ 60 years in Guinea savanna was indication that this
80 was the age bracket that was actively involved in agroforestry practices.

81 Majority of the respondents in the three agro-ecological zones were married men and women. Most
82 marriages were polygamous and had an average of more than five children that provided labour force for
83 farming. The result showed that 93.3 % of the respondents were married, 4.2 % were single, 0.8 %
84 divorced and 1.7 % widowed. About 92 % of the respondents were Muslims, 4.6% were mixture of Islam
85 and Traditional religion, 2.6 % and 0.9 % Christians and traditional worshippers, respectively.

86 Based on these findings, majority of the respondents acquired Islamic education as their highest
87 educational attainment. On average, it represented 41.0 % across the three zones. This was followed by
88 primary education with 17.9 %.

89 Farming was the major occupation in the three agro-ecological zones of the study area. The study
 90 revealed that an average of 66.1 % of the respondents as farmers. Other occupation in the study area
 91 included trading, civil service, fishing and cattle rearing among others.

92 **Involvement of Respondents in the Use of Agro-forestry practices (AP)**

93 Based on the list of agro-forestry practices identified by ICRAF, Nair (1990), some agro-forestry practices
 94 that were adopted in the study area were selected and the respondents were made to react to the practices,
 95 that is, their involvement in the use of any of the practices. The responses were classified into five: do not
 96 practise, practised but stopped, practise occasionally, practised but do not intend to continue, and practise
 97 regularly (Table 1 to 2).

98 Multipurpose trees Agroforestry system on farmland was mostly adopted in the study area (Table 1). The
 99 highest was in Sahel savanna with 91.7 %, followed by Guinea savanna (83.3%) while Sudan had 62.5%
 100 adoption. Farmers who had never practised at all, practised but stopped, practised occasionally or
 101 practised but did not intend to continue were negligible.

102 Also, Table 1 shows majority of the respondents in Sahel and Guinea savanna had never practised home
 103 gardens, with 54.2% each. Those farmers had adopted the system were relatively few, (Sahel 35.0%,
 104 Sudan savanna 20.8%, Guinea savanna 18.3%), while those who had practised but stopped, practiced
 105 occasionally or practised but did not intend to continue recorded very low scores.

106 Table 1 indicates that majority of the respondents across the zones had not adopted *Taungya* system of
 107 agro-forestry (Sahel savanna 91.7%, Sudan savanna 87.5% and Guinea savanna 81.7%). Only very few of
 108 the respondents had adopted the *Taungya* system of agro-forestry (Sahel 5.0, Sudan savanna 8.3 and
 109 Guinea Savanna 15.0%). No farmer had adopted the system but stopped practising it across the zones.

110 The Table also shows 55.0 % of the respondents in Sahel savanna had adopted woodlot system which was
 111 the highest across the zones, thus was followed by Sudan savanna with 50.0%, and Guinea savanna with
 112 44.2%. Those that had never adopted the system were 37.5 % in Guinea savanna, 35.0 % in Sahel
 113 savanna, and 26.7% in Sudan savanna.

114 **Table 1: Adoption of Multipurpose Trees, home-gardens, *Taungya* System and Woodlots on**
 115 **Farmland in Katsina State**

| Variables | Sahel | Sudan | Guinea |
|--------------------|---------|-----------|---------|
| Multipurpose Trees | Freq. % | Freq. % | Freq. % |
| Do not Practise | 4 (3.3) | 15 (12.5) | 7 (5.8) |

| | | | |
|---|------------------|------------------|------------------|
| Practised but Stopped | 0 (0) | 10 (8.3) | 3 (2.5) |
| Practise Occasionally | 6 (5.0) | 8 (6.7) | 5 (4.2) |
| Practised but do not Intend to Continue | 0 (0) | 12 (10) | 5 (4.2) |
| Practise Regularly | 110 (91.7) | 75 (62.5) | 100 (83.3) |
| Total | 120 (100) | 120 (100) | 120 (100) |
| Home-gardens | | | |
| Do not Practise | 65 (54.2) | 48 (40) | 65 (54.2) |
| Practised but Stopped | 0 (0) | 32 (26.7) | 0 (0) |
| Practise Occasionally | 13 (10.8) | 10 (8.3) | 33 (27.5) |
| Practised but do not Intend to Continue | 0 (0) | 5 (4.2) | 0 (0) |
| Practise Regularly | 42 (35.0) | 25 (20.8) | 22 (18.3) |
| Total | 120 (100) | 120 (100) | 120 (100) |
| Taungya system | | | |
| Do not Practise | 110 (91.7) | 105 (87.5) | 98 (81.7) |
| Practised but Stopped | 0 (0) | 0 (0) | 0 (0) |
| Practise Occasionally | 4 (3.3) | 0 (0) | 0 (0) |
| Practised but do not Intend to Continue | 0 (0) | 5 (4.2) | 4 (3.3) |
| Practise Regularly | 6 (5.0) | 10 (8.3) | 18 (15.0) |
| Total | 120 (100) | 120 (100) | 120 (100) |
| Woodlots | | | |
| Do not Practise | 42 (35) | 32 (26.7) | 45 (37.5) |
| Practised but Stopped | 0 (0) | 18 (15.0) | 7 (5.8) |
| Practise Occasionally | 0 (0) | 10 (8.3) | 15 (12.5) |
| Practised but do not Intend to Continue | 12 (10.0) | 0 (0) | 0 (0) |
| Practise Regularly | 66 (55.0) | 60 (50) | 53 (44.2) |
| Total | 120 (100) | 120 (100) | 120 (100) |

116 **Adoption of border planting, windbreaks/shelterbelt, Alley Cropping System and Woody Perennial**
117 **for Sustainable Soil Management**

118 A distribution of the respondents according to adoption of border planting as shown in Table 2, the Table
119 indicates majority of the respondents had never adopted the system across the zones (Sahel savanna
120 81.7%, Sudan savanna 74.2% and Guinea savanna 91.7%). No respondent adopted the system but stopped
121 practising it. Sudan savanna recorded the highest number of adopters with 18.3 %, followed by Sahel
122 savanna (10.0%), while Guinea savanna recorded only one adopter (0.8%). Majority of the respondents
123 across the Sahel and Sudan savanna zones had adopted the windbreaks system of agro-forestry. The
124 respondents from the Sahel savanna, however, recorded the highest number of adopters with 65.0 %,
125 followed by Sudan savanna with 55.0 %, while Guinea savanna recorded only 30.0 %. Conversely, for the
126 respondents that had never adopted the system, Guinea savanna recorded the highest number with 52.5%
127 (Table 2).

128 Table 2 indicates that majority of the respondents across the entire zones had never adopted the alley
129 cropping system (Sahel savanna 95%, Sudan savanna 80% and Guinea savanna 97.5%). Very few of the
130 respondents had adopted the system (Sahel savanna 3.3%, Sudan savanna 7.5% and Guinea savanna
131 1.7%). No farmer from Sahel and Guinea savanna had adopted the system and stopped, but in the Sudan

132 savanna 6.7% had. Similarly, for those that practised the system occasionally, the Sahel and Guinea
 133 savanna also recorded zero %, while Sudan had 2.5%. The Table also shows that respondents that had not
 134 adopted the system were highest in Guinea savanna with 87.5 %. This was followed by Sahel savanna
 135 with 81.7, while Sudan savanna recorded only 31.7%. On the other hand, the Sudan savanna recorded the
 136 highest number of adopters for this system, with 53.3 %; while Sahel and Guinea savanna had very low
 137 scores. No respondents had adopted the system but stopped practising it or did not intend to continue to
 138 do so across the zones.

139 **Table 2: Adoption of border planting, windbreaks/shelterbelt, Alley Cropping System and Woody**
 140 **Perennial in the Study Area**

| border planting | Sahel | | Sudan | | Guinea | |
|---|------------|--------------|------------|--------------|------------|--------------|
| | Freq. | % | Freq. | % | Freq. | % |
| Do not Practise | 98 | (81.7) | 89 | (74.2) | 110 | (91.7) |
| Practised but Stopped | 0 | (0) | 0 | (0) | 0 | (0) |
| Practise Occasionally | 0 | (0) | 5.0 | (4.2) | 9 | (7.5) |
| Practised but do not Intend to Continue | 10 | (8.3) | 4 | (3.3) | 0 | (0.0) |
| Practise Regularly | 12 | (10.0) | 22 | (18.3) | 1 | (0.8) |
| Total | 120 | (100) | 120 | (100) | 120 | (100) |
| windbreaks/shelterbelt | | | | | | |
| Do not Practise | 39 | (32.5) | 24 | (20) | 63 | (52.5) |
| Practised but Stopped | 0 | (0) | 10 | (8.3) | 8 | (6.7) |
| Practise Occasionally | 3 | (2.5) | 6 | (5.0) | 3 | (2.5) |
| Practised but do not Intend to Continue | 0 | (0) | 14 | (11.7) | 10 | (8.3) |
| Practise Regularly | 78 | (65.0) | 66 | (55.0) | 36 | (30.0) |
| Total | 120 | (100) | 120 | (100) | 120 | (100) |
| Alley Cropping System | | | | | | |
| Do not Practise | 114 | (95) | 96 | (80.0) | 117 | (97.5) |
| Practised but Stopped | 0 | (0) | 8 | (6.7) | 0 | (0) |
| Practise Occasionally | 0 | (0) | 3 | (2.5) | 0 | (0) |
| Practised but do not Intend to Continue | 2 | (1.7) | 4 | (3.3) | 1 | (0.8) |
| Practise Regularly | 4 | (3.3) | 9 | (7.5) | 2 | (1.7) |
| Total | 120 | (100) | 120 | (100) | 120 | (100) |
| Woody Perennial | | | | | | |
| Do not Practised | 98 | (81.7) | 38 | (31.7) | 105 | (87.5) |
| Practised but Stopped | 0 | (0) | 0 | (0) | 0 | (0) |
| Practised Occasionally | 9 | (7.5) | 18 | (15.0) | 5 | (4.2) |
| Practised but do not Intend to Continue | 0 | (0) | 0 | (0) | 0 | (0) |
| Practised Regularly | 13 | (10.8) | 64 | (53.3) | 10 | (8.3) |
| Total | 120 | (100) | 120 | (100) | 120 | (100) |

141 **Tree species combined with agricultural crops among the respondents in the Study Area**

142 Table 3 shows some of the commonest trees used in combination with agricultural crops among the
 143 respondents in the study area. In the Sahel savanna zone, those tree species with high scores include:
 144 *Azadirachta indica* with 25.8% *Parkia biglobosa* with 14.2% and *Adansonia digitata* having 13.3%. In
 145 Guinea savanna, those species recorded in decreasing number include: *Adansonia digitata*, with 9.2%,
 146 *Parkia biglobosa*, with 6.7% and *Azadirachta indica* having 5.8%. Across the three zones, the highest
 147 score is recorded by *Azadirachta indica*, having 13.3%, while *Adansonia digitata* and *Parkia biglobosa*

148 recording 9.7% respectively. Other tree species recorded include: *Borassus aethiopum*, *Anacardium*
 149 *occidentale* and *Tamarindus indica*.

150 **Table 3: Identified Tree Species Combined with Agricultural Crops Among the Respondents.**

| Tree Species | Sahel | | Sudan | | Guinea | Mean |
|-------------------------------|------------|--------------|------------|--------------|------------------|--------------|
| | Freq. | % | Freq. | % | Freq. % | % |
| <i>Adansonia digitata</i> | 16 | (13.3) | 8 | (6.7) | 11 (9.2) | 9.7 |
| <i>Azadirachta indica</i> | 31 | (25.8) | 10 | (8.3) | 7 (5.8) | 13.3 |
| <i>Anacardium occidentale</i> | 2 | (1.7) | - | - | 3 (2.5) | 1.4 |
| <i>Borassus aethiopum</i> | 2 | (1.7) | - | - | 01 (0.8) | 0.8 |
| <i>Parkia biglobosa</i> | 17 | (14.2) | 6 | (5.0) | 8 (6.7) | 8.6 |
| <i>Tamarindus indica</i> | - | - | - | - | 6 (5.0) | 1.7 |
| Combination/others | 52 | (43.3) | 96 | (80) | 84 (70) | 64.4 |
| Total | 120 | (100) | 120 | (100) | 120 (100) | (100) |

151 **Constraints to adoption of agro-forestry practices in the study area**

152 Table 4 reveals some of the problems militating against the adoption of agroforestry practices in the study
 153 area. The most serious problem preventing adoption of AP in Sahel savanna zone was scanty rainfall,
 154 which accounted for (30.8 %) of the problems. This was followed by lack of land (6.7 %) and inadequate
 155 labour (5.0 %). In the Guinea savanna zone, the most serious problem was high labour demand (8.3 %),
 156 followed by lack of land (6.7 %). Lack of required seedlings and scanty rainfall had 5.0 % respectively.
 157 The problems, in decreasing order of severity across the three zones were: scanty rainfall, lack of land,
 158 high labour demand, inadequate extension personnel and lack of transportation recording the same
 159 percentage, lack of required seedlings and lack of incentives recorded the same percentage.

160 **Table 4: Identified constraints to adoption of agro-forestry practices in the study area**

| Identified Problems | Sahel | | Sudan | | Guinea | Mean |
|------------------------------------|------------|--------------|------------|--------------|------------------|--------------|
| | Freq. | % | Freq. | % | Freq. % | % |
| High Labour Demand | 6 | (5.0) | - | - | 10 (8.3) | 4.4 |
| Lack of Required Tree Seedlings | 3 | (2.5) | - | - | 1 (0.3) | 1.1 |
| Inadequate Extension Personnel | 2 | (1.7) | - | - | 4 (3.3) | 2.5 |
| Lack of Land | 8 | (6.7) | 1 | (0.8) | 8 (6.7) | 4.73 |
| Scanty Rainfall | 37 | (30.8) | 3 | (2.5) | 6 (5.0) | 12.8 |
| Lack of Transportation | 4 | (3.3) | 1 | (0.8) | 4 (3.3) | 2.5 |
| Lack of Incentive | 2 | (1.7) | - | - | 2 (1.7) | 1.1 |
| Combination | 58 | (48.3) | 115 | (95.8) | 80 (66.7) | 70.3 |
| Total | 120 | (100) | 120 | (100) | 120 (100) | (100) |

161 **DISCUSSION**

162 **Socio-economic Attributes of the Respondents in the Study Area**

163 Respondents' gender showed that there were more male than female farmers across the zones. This
 164 implies that the male gender is more involved in agroforestry practices as compared to their female
 165 counterparts. Farming generally, is almost an exclusive business for the male in the study area. This may
 166 be as a result of the strenuous nature of most farming activities in general and agroforestry practices in
 167 particular. These activities are not attractive to women who often engage in household activities. The

168 paucity of women in agroforestry practices might also be attributed to culture and religion which made
169 access to women by male extension agents difficult since there were very few women extension agents.
170 In Sahel and Sudan savanna zones, majority of the respondents fell between 30 and 49 years. Except in
171 Guinea savanna, where majority of the respondents fell from ≤ 60 years. This dominant age bracket
172 among the rural dwellers implied that this is the age bracket that is actively involved in agroforestry
173 practices. This also meant that, the farming population is mainly made up of both the old and the middle-
174 aged people. The young ones might have migrated to urban areas in search of white collar jobs while
175 others might have been in school or too young to have farms. This has an adverse effect on the economy
176 of the rural people because at old age, farmers **cannot** have optimum productivity.

177 Also, majority of the respondents across the three zones were married men and women. Most marriages
178 were polygamous and have an average of more than five children that provided labour force for farming.
179 This is because agroforestry practices are not only capital intensive but also labour intensive. **This means**
180 **that there is a good source of labour. An explanation to this is that more adult members in a household**
181 **means that more quality labour would be available for carrying out farming activities and the practice of**
182 **agroforestry would not pose any problem. This agreed with the findings of Villano and Fleming (2004).**

183 As revealed in the study, educational level of the rural dwellers was low. However, on average, Islamic
184 education recorded the highest percent, which was followed by primary education. The study also
185 revealed that inspite of the low level of western education; they had indigenous/traditional knowledge and
186 high level of awareness about farming systems, tree species, shrubs, herbs and other agroforestry
187 practices. This indigenous knowledge affects their perception and willingness to participate in
188 agroforestry practices. However, they still needed more enlightenment and training on modern
189 agroforestry techniques as a means of sustainable land management. **Farmers who acquire some level of**
190 **education are more likely to perceive new technologies than the ones who have no any form of education.**
191 **According to Amaza and Tashikalma (2003), the literacy level of farmers is important as it determines the**
192 **rate of adoption of improved technology for increased productivity. Also, Adekunle (2009) pointed out**
193 **that the level of education of farmers will directly affects their ability to adapt to change and to accept**
194 **new ideas.**

195 Frequency analysis of the respondents' occupation revealed that farming is the major occupation in the
196 three agro-ecological zones of the study area. The study reveals on average, a high % of the respondents
197 as farmers. Other occupation in the study area includes: trading, civil servants, fishing and cattle rearing
198 among others.

199 **Adoption of Agro-forestry by the Respondents in the study Area**

200 The difference on agroforestry practices among the three zones (Sahel, Sudan and Guinea savanna) in the
201 study area was examined using analysis of variance (ANOVA). The test shows that there was a significant
202 difference in the use of agroforestry practices among the three zones ($F = 63.29$, $P < 0.05$). The
203 differences in adoption might be due to differences in soil types, rainfall patterns, socio-economic reasons
204 and readiness or the people to accept innovations.

205 Adoption occurs when one has decided to make full use of the new technology as a best course of action
206 for addressing a need (Rogers, 2003). The results of this finding showed majority of the respondents were
207 involved in food crop-production, while others were involve in rearing of animals, and planting of tree
208 crops such as *Azadirachta indica*, *Parkia biglobosa*, *Adansonia digitata* among others. Tree species such
209 as *Sesbania sesban* (L.) Merr, *Tephrosia vogelii* Hook .f., *Cajanus cajan*, *Gliricidia sepium* (Jacq.) Walp.,
210 *Leucaena leucocephala* (Lam.) De Wit., *Acacia angustissima* and *Tephrosia candida* (Madagascar) are
211 suitable for agroforestry (Kwesiga *et al.*, 2003; Mafongoya *et al.*, 2003). Puri and Bangarawa, (1992)
212 pointed out that the choice of tree species is the most important factor to be considered in agroforestry
213 practices. On the other hand Foroughbackhch (1992) stated that the choice of tree species be made after
214 careful consideration of their adaptability for growth and benefit for rural populace. Leguminous species
215 such as *Faidherbia albida* and *Leucaena leucocephala* cause a considerable improvement on crop yields.
216 Okali and Nwoboshi (1997) recorded poor performance of crops on apparently rich *Tectona grandis* and
217 *Eucalyptus camadulensis* soil. The farmer's preference of forest trees would definitely be due to their
218 potentials and adaptability to the land area.

219 Respondents' involvement in AP varied from zone to zone. The differences in adoption could be that an
220 innovation which was appropriate for a given zone might not necessarily be accepted in another zone. It
221 might also be due to socio-economic reasons, complexity and incompatibility of the innovation with the
222 existing practices. Thus, majority of respondents across the zones could not adopt *Taungya*, border
223 planting and alley cropping. Very few respondents across the zones adopted these systems of agro-
224 forestry. Conversely, multipurpose trees on farmland, improved fallow in shifting cultivation, home-
225 garden, woodlots and windbreaks or shelterbelt were much more adopted by farmers. The reason for low
226 adoption of *Taungya* system of agroforestry might be that food crop might compete with tree crop. Sahel
227 and Sudan zones adopted woodlots practices more than Guinea savanna.

228 Majority of the respondents adopted multipurpose trees on farmland across the zones. The findings
229 therefore reveal that this agroforestry system was popular among the farmers across the zones hence the

230 massive adoption. This might be due to the blend of the system with indigenous or traditional farming
231 practice across the zones.

232 The study also revealed that majority of the respondents in the study area had not practiced home-gardens.
233 Those farmers that had adopted the system were relatively few. While those that had practiced but
234 stopped, practice occasionally or practiced but do not intend to continue recorded very low scores.

235 The reason for their adoption might be to stem the environmental degradation in the Sahel and Sudan
236 savanna zones. Farmers could only take fuel-wood from these plantations and no other place.
237 Indiscriminate felling of trees for timber, fuel-wood and other domestic uses and clearing of land for
238 agricultural purposes and industrial development help to remove the forest cover; thereby exposing the
239 soil to wind erosion. Adeola (2001) observed that the system is used for various purposes such as
240 provision of wood, fodder, electric-poles, fencing poles, roofing poles, soil protection, soil reclamation
241 etc.

242 Sudan Savanna had the highest adoption of woody perennials for soil conservation across the zones. The
243 farmers' interest and adoption of the system could be to check the menace of annual flooding of this zone
244 which leads to soil and gully erosion. Plants help to stabilize the soil and other conservation works
245 thereby fulfilling one of the environmental functions of agroforestry (Baumer, 1990).

246 Majority of the respondents adopted multipurpose trees on farmland across the zones; this could be due to
247 good yield obtained if tree species are combined with agricultural crops in the study area. This implies
248 that native tree species enhance high yield of agricultural crops when combined; agroforestry system was
249 popular among the farmers across the zones hence the massive adoption. This might be due to the blend
250 of the system with indigenous or traditional farming practice across the zones.

251 On sources of information/awareness on some sustainable land use practices, the study revealed that
252 extension agents recorded high scores for all the land use practices in the study area. This may be due to
253 the availability of the agricultural development programme in the area. This study therefore agreed with
254 the findings of Onumadu (2002) who observed that agricultural agents were the most important source of
255 agricultural information to farmers. This view was also supported by Azeez (2002).

256 Majority of the respondent were of the view that they obtained information on sustainable land use
257 practices/agro-forestry practices through radio. This implies that farmers in these zones had several
258 options of other sources of information that could enhance or stimulate their use of AP. Although,

259 radio/mass media and extension agents were the principal means of the awareness, these two sources of
260 information could as well be responsible for the significance of the AP. Mass media plays an important
261 role in the dissemination of information on agricultural activities as it enables even the cattle rearers that
262 roam about in the bush to have access to the information on agricultural activities through their radio. This
263 was also supported by Onumadu (2002), who observed that mass media was one of the most important
264 sources of seeking information on agro-forestry practices.

265 Other sources such as traditional and a combination of one or more of these sources also recorded
266 relatively high scores, whereas sources such as relatives and neighbour recorded very low scores; while
267 some of the respondents reported that they had no information at all. This may be due to lack of adequate
268 publicity or enlightenment. This calls for an increase in agricultural extension agents who should take up
269 the responsibility of educating, training and monitoring of farmers in the areas of food crop production.

270 The relationship between sources of information and use of agroforestry practices. It is observed that,
271 among the three agro-ecological zones, it is only in Guinea savanna that sources of information was not
272 significantly related to use of agroforestry practices by the farmers ($\chi^2 = 0.32, P > 0.05$). Conversely, in
273 Sahel and Sudan savanna zones, source of information was significantly related. In Sahel savanna zone
274 $\chi^2 = 0.01, P < 0.05$ and in Sudan savanna, $\chi^2 = 0.03, P < 0.05$.

275 Farmers in the study area may have adopted AP because of the various benefits they derive from it. These
276 benefits range from social, economic and environmental benefits. The social benefits in the study area
277 include provision of fruits and leaves, provision of shade, provision of fuel-wood, provision of fodder and
278 medicinal herbs. Correlation result on the relationship between use and benefits of agroforestry practices
279 across the three agro ecological zones of Katsina state showed a significant positive relationship between
280 the use and benefits of agroforestry practices, with a Pearson correlation (r) of 0.160, ($P < 0.05$). This
281 implies that farmers derived immense benefits from the use of agroforestry practices; and that farmers
282 who use agroforestry practices more, also benefit more.

283 CONCLUSION

284 Based on the findings of this study, windbreaks, scattered trees on farmland, woodlots, improved fallow
285 and home-gardens were the various AP that were common in the study area. There was difference in
286 adoption of AP among the three agro-ecological zones. The differences in adoption could be that an
287 innovation in which was appropriate for a given zone might not necessarily be accepted in another zone.
288 This could also be due to soil and climate type and socio-economic reasons. In order to sustain and even
289 increase our agricultural productivity and to reduce, to the barest minimum, the effects of desertification

290 and environmental degradation, the following recommendations are made: Government should encourage
291 the adoption of agroforestry as a system of multiple land use to increase wood and food production
292 thereby ensuring the optimum use of land. Provide incentives such as seedlings, transportation, inorganic
293 fertilizers and tractors to farmers so as to encourage them to participate actively in agroforestry activities.
294 The use of more indigenous tree species that can improve soil fertility and at the same time more
295 adaptable to the environment should be promoted (e.g. *Parkia biglobosa*). Application of organic
296 fertilizers and planting of leguminous trees will help to resuscitate the soil for high productivity. A study
297 to re-examine the factors limiting the adoption of some AP that have low adoption in the study area such
298 as alley cropping and *Taungya* is recommended.

299 REFERENCES

- 300 Adekunle , V. A. (2009): Contributions of agroforestry practice in Ondo State, Nigeria, to environmental
301 sustainability and sustainable agricultural production. *Afrika Focus -Volume 22, Nr. 2, 2009 -pp.*
302 *27-40.*
- 303 Adeola, A.O. (2001): Farmer's perception of social forestry impact in the semi-arid zone of Nigeria.
304 *Unpublished Ph.D. Thesis*, Department of Forest Resources Management, University of Ibadan,
305 275 Pp.
- 306 Akpabio IA, Abagale FK, Addo J, Adisenu-Doe R, Anthony KM, Apana S, Boateng EA, Esu BB,
307 Adedire MO (2008): Gender perception on constraints affecting agroforestry practices in Akwa,
308 Ibom State, Nigeria. *Agric. J. 3(5):375-381.*
- 309 Amaza, P.S and Tashikalma, A.K (2003): Technical Efficiency of Groundnuts in Adamawa State,
310 Nigeria, *Journal of Arid Agriculture, 13: 127 – 1311.*
- 311 Azeez I.O. (2002): Evaluation of media mix for disseminating conservation information in South Western
312 Nigeria. Ph.D. Thesis (unpublished), Department of Forest Resources Management, University of
313 Ibadan, Nigeria, 230 Pp.
- 314 Baumer, M. (1990): Agroforestry and Desertification. Technical Centre for Agricultural and Rural
315 Cooperation. *The Netherlands 250 Pp.*
- 316 Forough backhch, R. (1992): Establishment and Growth Potential of Firewood Species in Northeastern
317 Mexico. *Agroforestry Systems 19 (2):95-108.*
- 318 Irshad M, Khan A, Inoue M, Ashraf M, Sher H (2011): Identifying factors affecting agroforestry system
319 in Swat, Pakistan. *Afr. J. Agric. Res. 6(11):2586-2593.*
- 320 Jamala, G. Y1, H. E. Shehu, J. J. Yidau and L. Joel (2013): Factors Influencing Adoption of Agro-
321 Forestry among Smallholder Farmers in Toungo, Southeastern, Adamawa State, Nigeria; *IOSR*
322 *Journal of Environmental Science, Toxicology and Food Technology; 6 (6): 66-72.*
- 323 Mafongoya, P. L., Chintu, R., Chirwa, T. S., Matibini, J., & Chikale, S. (2003): Tephrosia species and
324 provenances for improved fallows in southern Africa. *Agroforestry Systems, 59(3), 279-288.*

- 325 Maluki J. M., Kimiti J. M., Nguloo S. and Musyoki J. K. (2016): Adoption levels of agroforestry tree
326 types and practices by smallholders in the semi-arid areas of Kenya: A case of Makueni County;
327 Journal of Agricultural Extension and Rural Development; 8 (9): 187-196.
- 328 Nair, P. K. R. (1993): An Introduction to Agroforestry. Kluwer, Boston.
- 329 Okali, D.U.U and Nwoboshi, L.C. (1997): A Preliminary Study of Maize Planted among Coppiced Teak.
330 Mimeograph. Department of Forest Resources Management University of Ibadan, Nigeria. 8pp.
- 331 Onumadu, F.N. (2002): Determinants of Adoption of Agroforestry Practices by Small Scale Farmers in
332 Katsina State, Nigeria. Unpublished P.h.D. Thesis, University of Ibadan 259 Pp.
- 333 Puri, S and Bangarawa, K.S. (1992): Effects of Trees on the Yield of Irrigated Wheat Crop in Semi-arid
334 Regions. Agroforestry System 20 (3): 229-241.
- 335 Rogers, E. M. (2003): Diffusion of Innovations (Fifth ed.). New York: The Free Press.
- 336 Villano, R. and Fleming, E. (2004): Analysis of Technical Efficiency in a Rain-fed Lowland Rice
337 Environment in Central Luzon Philippines using a stochastic frontier paper series in Agricultural
338 and Resource Economics No. 2004 – 15, University of New England, Armidale