

FUNGITOXIC POTENTIALS OF EXTRACTS OF PLANT ORIGIN AGAINST FUGAL ROOT ROT OF CASSAVA (*Manihot esculenta* Crantz) IN STORAGE

ABSTRACT

Investigations were carried out on the use of the water and ethanolic extracts of *Piper guineense*, *Ocimum gratissimum*, *Casia alata*, and *Tagetes erecta* in the management of postharvest deterioration of cassava root caused by *Aspergillus flavus* and *Rhizopus stolonifer*. Water and ethanolic extracts of the plant materials had significant differences ($p \leq 0.5$) in their rates of fungitoxicity on the pathogenic organisms. Water and ethanol extracts of *C. alata* and *T. erecta* respectively at 50% concentration gave the same highest radial growth inhibition of 80.20% on *A. flavus* *in vitro* followed by ethanol extracts of *C. alata*, *O. gratissimum*, and *P. guineense*. The ethanolic extract of *T. erecta* at 50% concentration gave the highest inhibitory effect of 53.50% on *R. stolonifer* followed by ethanol extracts of *C. alata*, *O. gratissimum*, and *P. guineense* whereas the least growth inhibition of 0.17% was recorded by aqueous extract of *P. guineense* on *R. stolonifer*. *In vivo* test of the plant extracts applied before and after inoculation with spore suspension (1×10^5 spores/ml of distilled water) of test fungi showed significant reduction in root rot incidence and severity. The lowest incidence and severity of cassava root rot of 16.5% and 1.45 respectively were recorded with *T. erecta* ethanol extracts applied before inoculation of *A. flavus* indicating that the extracts of the plant materials could be better used as protectant than eradicant in the control of post harvest fungal deterioration of cassava root. *R. stolonifer* showed stronger resistance to the extracts of the plant materials than *A. flavus* during pathogenesis *in vivo*.

Key words: Cassava, root rot, postharvest, *Piper guineense*, *Ocimum gratissimum*, *Casia alata*, and *Tagetes erecta*

INTRODUCTION

Cassava (*Manihot esculenta* Crantz), is an important crop contributing to the survival of human beings and livestock by providing a ready and cheap source of carbohydrate for food and feed as well as raw material for industries (Nweke, 2015; Markson *et al.*, 2012). It is the third largest source of carbohydrate after rice, sugar and maize in the world and a basic staple food and main

source of energy for majority of the people in Africa and many other parts of the world (Amadioha, 2012; Echebiri and Edaba, 2008; Bua and Okello, 2011; Bukanga, 1999).

Cassava is a basic staple food to more than 70% of Nigerian population (Eke-okoro and Njoku, 2012) and a reliable and convenient source of food for tens of millions of rural and urban dwellers in Nigeria in its processed form (IITA, 2010; Nweke *et al.*, 2002; Taiwo, 2006; Philip *et al.*, 2006). In addition to human consumption, cassava is used for the production of bioethanol, animal feed, and starch for industrial products (Plucknett *et al.*, 2003).

Cassava root rot diseases are some of the major constraints to achieving the full potentials in cassava production in Nigeria and many areas in Africa (Chalwe *et al.*, 1999; Onyeka *et al.*, 2008; Onyeka, 2002; Bua and Okello, 2011). Apart from reducing cassava yield, root rot diseases caused by different organisms including fungi, bacteria and other pest organisms can also reduce the quality of cassava roots harvested and their products. Some of the fungi found to be pathogenic on cassava roots include *Sclerotium rolfsii*, *Fusarium oxysporum* Schlecht, *Botryodiplodia theobromae* Pat, *Aspergillus niger* Van Tieghem, *Aspergillus flavus* Link, *Rhizopus spp*; *Fusarium solani* (Mart) Sacc., and *Macrophomina phaseolina* (Tassi) Goidanich (Okigbo *et al.*, 2009a; IITA, 1990). Different control measures have been suggested and used for the control of post-harvest cassava root rot diseases. In view of the problems associated with curing and use of synthetic chemicals in the control of storage rot of cassava root, natural plant extracts have been evaluated as pesticide alternatives in the management of postharvest root rot diseases of cassava incited by *A. flavus* and *R. stolonifer*.

MATERIALS AND METHODS

Source of plant Materials

The cassava root (TME 419 Variety) both infected and healthy (uninfected) were obtained from the National Root Crops Research Institute, Umudike, Abia State, Nigeria. TME 419 Variety chosen for high yielding but highly susceptible to root rot of cassava in storage. The fresh leaves of *Ocimum gratissimum* and *Piper guineense* were obtained from open market stalls in Umuahia, Abia State while *Cassia alata* and *Tagetes erecta* were collected from Umudike, Abia State.

Culture Medium

Potato Dextrose Agar (PDA) (39g) was poured into one liter conical flask and made up to a liter with sterile distilled water, mixed thoroughly and melted in an electric water bath and then sterilized by autoclaving at 120°C for 15 minutes. The sterile medium was allowed to cool (46°C) and 15ml dispensed into sterile Petri-dishes and allowed to solidify. The sterile solidified medium was used for all the microbial cultures and other investigations

Isolation and Identification of Fungal Pathogens

The **rotten** cassava roots were washed with tap water, surface sterilized with 70% ethanol solution and rinsed in sterile distilled water. Pieces of the rotted tissue (3 mm diameter) were collected from the boundary of the infected and healthy root and placed on the culture medium in Petri dishes. The inoculated plates were transferred into the microhumidity chamber and incubated at 26°C. The emerging fungal colonies were sub-cultured on fresh sterile culture medium of PDA to obtain pure cultures of the isolates. **Pathogenicity** test was carried out on the isolates using fresh healthy, washed and sterilized cassava roots (Amadioha, 2001). On establishment of rot condition, re-isolation was carried out to obtain pure cultures of the inoculated isolates which were then compared with the original isolates. The isolates that caused

root rot of the cassava were regarded as pathogens and were **identified by reference to the illustrated fungal genera of Sangoyomi (2004; 2010).**

Leaf Extracts.

Fresh leaves of *O. gratissimum*, *P. guineense*, *C. alata* and *T. erecta* were washed under running tap water, rinsed with sterile distilled water and **air dried at room temperature (27°C)**. The dried leaves of the test plant materials were milled into powder and separately weighed (10g, 20g, 30g, 40g and 50g) into a beaker before adding 100ml each of the extracting solvent (ethanol or sterile distilled water). Each solution was thoroughly mixed and left to stand for 24 hours and then filtered separately using a four –fold cheese cloth into a beaker. These filtrates constituted 10%, 20%, 30%, 40% and 50% concentrations of cold water or ethanol leaf extracts of the test plant materials. The purity of the extracts was confirmed using the method of Cheesbrough (2000).

Effect of Extracts on the radial growth of fungal pathogens *in vitro*

The method of Amadioha and Obi (1998) was used to evaluate the antifungal effect of extracts of the test plants against fungal growth *in vitro*. 2 ml each of the extract concentrations (10%, 20%, 30%, 40% and 50%) was separately transferred into a sterile Petri dish with the aid of a sterile pipette. Freshly prepared molten PDA (15ml) was aseptically poured into the plates. The plates were rotated gently for easy mixing of the PDA-extract media which were allowed to solidify. A 5mm diameter disc of each pathogen was then dropped separately at the centre of the solidified extract-PDA plates. The treatments were replicated three times. **The control plates consisted of only PDA (15ml) + 2ml of distilled water or 70% ethanol (no extracts) inoculated with the test fungi.** The inoculated Petri dishes were incubated at 27°C and observed daily for fungal

growth. The mycelial radial growth of each fungus was measured with a ruler along the two directions on the perpendicular lines drawn on the reverse side of the plates after the growth in the control experiment had reached the edge of the plate. The mean colony diameter of the three replicates was taken as the mean growth of each treatment. Fungitoxicity was calculated as percentage colony inhibited by the extracts (Amadioha, 2004).

$$\% \text{ Fungal Growth inhibition} = \frac{DC - DT}{DC} \times \frac{100}{1}$$

Where DC = Average diameter of colony in control experiment.

DT = Average diameter of fungal colony with extract treatment.

***In vivo* Screening of Plant Extracts against fungal pathogens**

The 50% extract concentration of each of the water and ethanol plant extracts which gave the highest radial growth inhibition of the pathogens *in vitro*, was used in this experiment. The method of Amadioha and Markson (2007a) was adopted as follows:

Group A - ten uninfected cassava roots each dipped into the extract concentration of test plants and allowed to air dried for 2 hrs before spray-inoculating with the spore suspension (1×10^5 spores/ml of distilled water) of the test fungal pathogens.

Group B - ten uninfected cassava roots each spray-inoculated with the spore suspension (1×10^5 spores/ml of distilled water), air dried for 2 hours and then dipped into the extract concentration.

The control experiment constituted the spray-inoculated cassava roots that were treated with the respective extracting solvents (water or ethanol) only. Each of the treated cassava roots including the control was enclosed separately in polyethylene bags with cotton wool soaked with distilled water (micro humidity chamber) and incubated at $28 \pm 2^\circ\text{C}$. The experiment

was replicated two times. The samples were observed daily for rot development for 14 days. The disease incidence and severity were assessed.

$$\text{Disease incidence (\%)} = \frac{\text{No. of rotted cassava roots}}{\text{Total No. of cassava roots}} \times \frac{100}{1}$$

Disease Severity was assessed (Murugan and Luaina, 2013) on a 0-5 scale as follows:

- 0. - No infection
- 1 - Slight infection ($\leq 10 - 20\%$ of cassava root infected)
- 2 - Moderate infection (21 - 40% of root infected)
- 3 - High infection (41 - 60% of root infected)
- 4 - Extensive infection (61 - 80% of root infected)
- 5 - Complete rot (81 - 100% of root infected)

$$\text{Disease severity index} = \frac{\text{Sum of all scores}}{\text{Number of plants scored (N) x Highest score (5)}} \times \frac{100}{1}$$

Where; N is the total number of cassava root assessed; 5 - the maximum score of the scale used.

RESULTS

Effect of plant extracts on the radial growth of pathogens

The *in vitro* screening of the plant extracts against the radial growth of the test fungal pathogens, *A. flavus* and *R. stolonifer* showed that the plant extracts had significant ($P \leq 0.05$) inhibitory effects on the organisms tested. The inhibitory effect of the test plants increased with concentration and differed with extracting solvents across the test organisms. *Tagetes erecta* ethanol extract recorded the highest mean inhibitory effect on *A. flavus* (66.9%) while the least was *P. guineense* aqueous extract with mean inhibition of 10.5% on *R. stolonifer*. Both ethanol

and water extracts of the test plant materials at 50% concentrations had the highest inhibitory effects on the radial growth of all the pathogenic organisms whereas the lowest mean values were recorded with 10% concentration of test plant materials. The water and ethanol extracts of *C. alata* and *T. erecta* at 50% concentration gave the highest radial growth inhibition of 80.20% and 82.0% respectively of *A. flavus* in culture. All the concentrations tested recorded more inhibitory effects with ethanol as extracting solvent than water and this was significant ($p \leq 0.05$) (Table 1). *R. stolonifer* showed a stronger resistance across all the plant extracts tested.

Table 1: Percentage growth inhibition of aqueous and ethanol plant extracts on *R. stolonifer* and *A. flavus* in culture.

Plant Extract Concentration	Fungal Radial Growth Inhibition (%)			
	<i>A. flavus</i>		<i>R. stolonifer</i>	
	WE	EE	WE	EE
<i>Piper guineense</i>				
10	6.17*±1.4**	23.0±1.6	0.17±0.8	7.33±1.0
20	23.0±0.4	30.0±1.2	3.33±1.5	16.8±0.7
30	37.0±0.2	40.0±0.9	12.7±0.8	25.5±0.9
40	43.7±0.8	49.0±0.7	17.5±1.3	30.5±1.0
50	50.2±1.6	54.5±1.3	21.5±1.6	40.5±0.6
Control	0.00	0.00	0.00	0.00
Mean	30.8±0.9	38.5±1.1	10.5±1.2	23.6±0.8
<i>Occimum gratissimum</i>				
10	13.3±1.7	25.5±0.9	0.50±0.8	10.5±1.5
20	32.8±0.8	39.3±0.6	11.8±0.6	25.2±0.9
30	40.5±1.4	53.2±1.4	19.7±0.9	30.2±0.7
40	56.5±0.7	63.2±1.2	24.5±1.3	41.8±0.5
50	62.8±0.5	67.5±0.9	26.2±1.2	48.2±0.8
Control	0.00	0.00	0.00	0.00
Mean	40.0±1.0	48.5±1.0	16.0±1.0	30.6±0.9
<i>Cassia alata</i>				
10	24.3 ±0.5	34.5±1.3	1.67±1.4	7.33±0.9
20	37.3 ±1.3	46.7±1.5	6.50±0.8	15.5±1.2
30	63.3±0.8	64.3±0.8	10.9±1.0	20.5±0.8

40	71.5±0.4	71.8±0.6	24.8±0.7	30.8±1.3
50	80.2±0.3	74.0±0.7	32.3±0.9	34.8±0.4
Control	0.00	0.00	0.00	0.00
Mean	53.9±0.7	57.1±1.0	14.7±1.0	21.3±0.9

Tagetes erecta

10	28.5 ±0.8	45.0±1.3	6.17±0.8	16.0±0.7
20	46.2±0.5	64.7±0.6	21.8±0.5	24.5±0.4
30	61.5±1.0	72.3±0.3	29.8±0.7	33.5±0.7
40	73.2±0.9	76.8±0.5	35.0±0.4	40.2±0.3
50	79.3±0.6	82.0±0.2	49.2±0.9	53.5±0.6
Control	0.00	0.00	0.00	0.00
Mean	56.5±0.8	66.9±0.6	27.9±0.7	33.0±0.5

LSD (5%) Conc.	3.78	3.03
LSD (5%) Extract	1.96	1.58

WE = water extract, EE = ethanol extract

*Values are means of three replicates in two separate experiments.

** Standard Error

Effect of plant extracts on disease incidence and severity *in vivo*

The *in vivo* screening of plant extracts applied before and after spray-inoculating with the spore suspension (1×10^5 spores/ml of distilled water) of the test fungi indicated that incidence of cassava root rot was significantly reduced by the treatment both before and after inoculation when compared with the control experiment (Table 2). The same trend was observed in the severity of cassava root rot (Table 3). *P. guineense* water extract had the highest percentage disease incidence of 48.6% when it was applied after the inoculation with spore suspension of *R. stolonifer* followed by *O. graticimum*, *C. alata* and *T. erecta*. The lowest incidence and severity of cassava tuber rot of 16.5% and 1.45 respectively were recorded with *T. erecta* ethanol extracts applied before inoculation of *A. flavus*. Generally, the extracts of *T. erecta* had a stronger inhibitory effect on the pathogens than the other three plant extracts whereas *R. stolonifer*

showed stronger resistance to the plant extracts than *A. flavus* both before and after spray-inoculating with the pathogenic organisms.

Table 2: Effect of aqueous and ethanol plant extracts applied before and after inoculation on the disease incidence by *R. stolonifer* and *A. flavus* in vivo

Treatment	Pathogens and Disease Incidence (%)			
Plant Extracts	<i>A. flavus</i>		<i>R. stolonifer</i>	
	A	B	A	B
<i>Piper guineense</i>				
Water extract	43.3*±1.4**	34.2±0.9	48.6±1.5	38.9±0.8
Ethanol extract	32.3±2.1	28.2±1.6	40.4±0.9	32.0±1.4
<i>Ocimum gratissimum</i>				
Water extract	40.5±1.6	27.0±0.8	47.5±1.3	36.8±1.4
Ethanol extract	30.5±0.7	21.9±1.4	38.2±0.8	27.2±1.0
<i>Cassia alata</i>				
Water extract	40.3±0.9	25.4±1.5	43.3±0.6	33.8±1.7
Ethanol extract	30.3±0.8	20.3±0.6	31.9±1.0	22.3±2.0
<i>Tagetes erecta</i>				
Water extract	34.2±1.5	24.5±1.2	40.5±0.9	30.5±1.5
Ethanol extract	26.8±1.3	16.8±1.7	30.3±0.8	20.5±0.6
Control	50.50±1.4	45.68±1.2	62.83±1.6	58.60±1.0
LSD (5%)	3.71		1.48	

*Values are means of three replicates in two separate experiments.

** Standard Error

187 **Table 3: Effect of aqueous and ethanol plant extracts applied before and after inoculation**
 188 **on the disease severity of cassava root incited by *R. stolonifer* and *A. flavus*.**

Treatment	Pathogens and Disease Severity Index			
Plant Extracts	<i>A. flavus</i>		<i>R. stolonifer</i>	
	A	B	A	B
<i>Piper guineense</i>				
Water extract	4.30*±1.4**	3.47±0.8	3.67±1.0	4.73±0.9
Ethanol extract	3.14±0.8	2.20±0.7	2.20±1.6	2.47±1.8
<i>Ocimum gratissimum</i>				
Water extract	4.17±1.0	3.20±1.5	3.13±0.9	4.10±0.7
Ethanol extract	3.03±1.2	2.20±1.8	3.20±0.7	1.97±1.3
<i>Cassia alata</i>				
Water extract	3.43±1.5	3.50±1.2	3.33±0.8	3.30±0.9
Ethanol extract	3.17±1.6	1.97±0.8	2.17±1.3	2.10±0.7
<i>Tagetes erecta</i>				
Water extract	3.30±0.6	2.50±1.0	3.10±0.8	3.17±0.5
Ethanol extract	3.23±0.7	1.45±0.4	2.37±0.9	1.77±1.2
Control	6.7±0.4	4.37±1.5	5.47±1.4	4.96±1.8
LSD (5%)	0.42		0.38	

189 *Values are means of three replicates in two separate experiments.

190 ** Standard Error

DISCUSSION

The pathogens, *Aspergillus flavus* and *Rhizopus stolonifer* inciting cassava root rot in this study have been previously linked with postharvest rot of cassava, yam and cocoyam (Okigbo *et al.*, 2015; 2014; 2009a). Amadioha and Markson (2007b) and Okigbo *et al.*, (2009b) implicated *Botryodiplodia acerina* and *A. niger* as the leading cause of postharvest fungal rot of cassava which is at variance with *R. stolonifer* and *A. flavus* in this study. The difference could be due to the variety and age of the cassava root used and location especially the prevailing environmental factors where the studies were conducted.

The result of the inhibitory potentials of the water and ethanolic extracts of test plant materials against the radial growth of the two fungal pathogens showed significant differences ($p \leq 0.5$) in their rates of fungitoxicity on *A. flavus* and *R. stolonifer*. The ethanolic extract of *T. erecta* at 50% concentration gave the highest inhibitory effect on *A. flavus* whereas the least radial growth inhibition was recorded by 10% water extract of *P. guineense* on *R. stolonifer*. This implies that the two fungal pathogens showed differences in their rates of resistance or susceptibility to the extracts of the test plants with *R. stolonifer* being less susceptible than *A. flavus* indicating that the fungus may have devised means of resisting the effect of the plant extracts (Umana *et al.*, 2016). The different concentrations of the plant extracts also showed significant differences in their mean growth inhibitory effects with the higher concentrations (50%) having more inhibitory effects on the pathogens than the lower concentrations. This corroborates the work of Suleiman (2010) and Amadioha (2006) who recorded a significant difference in mycelial growth inhibition by various plant extract concentrations with higher concentrations giving remarkable fungitoxic

effect. This shows that the antimicrobial potentials of the test plant materials can effectively be realized at higher concentrations of extracts (Nwinyi *et al.*, 2009). Also, the higher concentrations of the test plant materials may have contained more active ingredients due to higher dilution in the extracting solvent than the lower concentration with low dilution due to some inhibitory factors (Umana *et al.*, 2016). This finding also agrees with the reports of Amadioha, (2000) and Okigbo *et al.*, (2009a) that the difference in the fungitoxicity of extracts may be due to the differences in the solubility of the active ingredients or compounds in extracting medium. The study on the fungitoxic potentials of the various extracts of the plant materials showed that *T. erecta* and *C. alata* were more fungitoxic and exhibited the highest percentage growth inhibition on the radial growth of the rot pathogens suggesting that the plant materials contain some active compounds/phytochemicals that affected the radial growth of the rot pathogens in culture. The ethanolic extracts recorded the highest radial growth inhibition in culture across all the concentrations and plant materials. This observation suggests that ethanol as extracting medium dissolved more active compounds present in the plant materials than water which dissolved less active principles or compounds (Anukworji *et al.*, 2012). *P. guineense* gave the lowest growth inhibitory effect whereas *T. erecta* recorded the highest inhibitory effects across the test fungi indicating that the pathogenic organisms reacted differently to the phytochemicals of test plant materials extracted by different extracting solvents.

The results of *in vivo* test of the plant extracts applied before and after inoculation with spore suspension of test fungi and their effect on the incidence and severity of cassava root rot indicated significant differences ($p \leq 0.05$) with the application of the extracts before inoculation recording a lower disease incidence and severity by the rot causing organisms than application of the extracts after inoculation. However, the water and ethanolic extracts of the test plant

materials showed significant differences in their reduction of disease incidence and severity caused by the test organisms when compared with the control experiment. The least percentage tuber rot was recorded in tubers treated with *T. erecta* ethanol extract applied before inoculation of *A. flavus* while the highest incidence of cassava root rot was recorded with *P. guineense* water extract applied after inoculation of *R. stolonifer*. It was observed that *T. erecta* and *C. alata* reduced the growth and spread of the test fungi during pathogenesis than *P. guineense* and *O. graticimum* *in vivo* while *R. stolonifer* had more resistance to the plant extracts than *A. flavus*. Extracts of *T. erecta* and *C. alata* could therefore be exploited as biopesticide and alternative to synthetic chemicals by resource poor farmers in the control of storage rot of cassava caused *A. flavus* and *R. stolonifer*.

CONCLUSION

The water and ethanolic extracts of *P. guineense*, *O. graticimum*, *C. alata*, and *T. erecta* were effective in the radial growth inhibition *in vitro* and the development and spread of postharvest root rot of cassava caused by *A. flavus* and *R. stolonifer* *in vivo*. The extracts had significant differences ($p \leq 0.5$) in their rates of fungitoxicity on the pathogenic organisms with ethanolic extracts giving higher inhibitory effect than the aqueous extracts at 50% concentration. Application of the plant extracts before inoculation with spore suspension (1×10^5 spores/ml of distilled water) of test fungi was more effective in reducing the incidence and severity of cassava root rot disease than when the extracts were applied after inoculation with the pathogens indicating that the extracts of the plant materials could be better used as protectant than eradicant in the control of post harvest fungal deterioration of cassava root incited by *A. flavus* and *R. stolonifer*.

REFERENCES

- Amadioha, A. C. (2000). Fungitoxic effects of some leaf extracts against *Rhizopus oryzae* causing tuber rots of potato. *Arch Phytopathol.* 34: 1-9.
- Amadioha, A.C. (2001). Fungicidal Activity of Some Plant Extracts against *Rhizoctonia Bataticola*. *Arch Phytopathol.*, 33 (6): 509-517.
- Amadioha, A.C. (2004). Control of black rot of potato caused by *Rhizoctonia bataticola* using some plant leaf extracts. *Arch Phytopathol*, 37: 111-117.
- Amadioha, A.C. (2006) Prospect in developing of eco-friendly biopesticides from neem (*Azadirachta indica*) in Nigeria. Proceedings of the 2006 International Neem Conference Kunming, China, November 11-15, 2006, p 51-58.
- Amadioha, A.C. (2012) Reducing food losses through sustainable methods of plant disease management: An imperative for actualization of food security in Nigeria. 13th Inaugural Lecture, Michael Okpara University of Agriculture, Umudike. June 12, 2012.
- Amadioha, A. C. and Obi V. I. (1998) Fungitoxic activity of extracts from *Azadirachta indica* and *Xylopi aethiopica* on *Colletotrichum lindemuthianum* in cowpea. *Journal of Herbs, Spices and Medicinal Plants*, 6 (2): 33-40.
- Amadioha, A. C. and Markson, A. A. (2007a) Postharvest control of cassava tuber rot caused by *Botrydiplo dia acerina* using extracts of plant origin. *Arch Phytopathol*, 40 (5): 359-366.
- Amadioha, A.C. and Markson, A. A. (2007b). Control of storage rot of cassava tuber caused by *Rhizopus oryzae* using some plant extracts. *Arch Phytopathol*, 40 (6): 381-388.
- Anukworji, C.A., Putheti, R.R., and Okibgo, R.N. (2012) Isolation of fungi causing rot of cocoyam (*Colocasia esculenta* (L.) Schott) and control with plant extracts: (*Allium sativum* L., *Garcinia kola*, *Azadiachta indica* and *Carica papaya* L.). *Global Adv. Res. J. Agric Sci.*, 1(2): 33-47.
- Barnett, H. L. and Hunter, B. B. (1987) Illustrated Genera of Imperfect Fungi 4th Edition. Macmillan Inc. New York, pp. 218.
- Bua, B. and Okello, C. (2011) Isolation and Identification of Cassava Root Rot Disease Causal Pathogens from Lira District, Uganda. African Crop Science Conference Proceedings, 10: 183-186.
- Bukanga, M. (1999) Cassava: Postharvest Operations. International Institute of Tropical Agriculture, Ibadan, Nigeria. (<http://www.cgiar.org/iita/>)

- Chalwe A., Malambo, C., Zimba, K., Nawa, I and Muimba-Kankolongo, A. (1999) Cassava root rots and associated micro-organisms in Zambia. In "Food security and crop diversification in SADC countries: the role of cassava and sweet potato, Akoroda, M.O. and Teri, J.M. (eds). Pp 317 – 324. Proceedings of the scientific workshop of SARRNET, 17-19 August 1998, Zambia.
- Cheesbrough, M. (2000) *District Laboratory Practice in Tropical Countries*, Pt- 2. Cambridge University Press, pp. 64-65.
- Echebiri, R. N. and Edaba, M.E. I. (2008) Production and utilization of cassava in Nigeria: Prospects for food security and infant nutrition. *Patnsukjournal*, 4 (1): 38-52.
- Eke-okoro, O. N. and Njoku D. N. (2012) A Review of Cassava Development in Nigeria from 1940-2010. *ARPN Journal of Agricultural and Biological Science*, 7 (1): 59-64.
- IITA (1990) International Institute of Tropical Agriculture, Ibadan, Nigeria. Cassava in Tropical Africa: A Reference Manual. p.176.
- IITA (2010) International Institute of Tropical Agriculture, Ibadan, Nigeria. Postharvest Technology". Annual report. p. 62-80.
- Markson, A. A., Umana, E. J., Udo, S. E., Omosun, G., Madunagu, B. E., and Osim, M. O. (2012) Tissue masceration and starch depletion caused by some rot pathogens of cassava (*Manihot esculenta* Crantz). *International Journal of Current Research*, 4 (04): 99-104.
- Murugan, K. and Lubaina, A.S. (2013) Physiological and biochemical characterization of *Senna alata* (L.) Roxb. Leaf Extract - A plant based fungicide against *Alternaria* Leaf Spot in Sesame. *World Journal of Pharmacy and Pharmaceutical Sciences*, 2 (6): 5790-5801.
- Nweke, F. U. (2015). Effect of some plant leaf extracts on mycelia growth and spore germination of *Botryodiplodia theobromae* causal organism of yam tuber rot. *Journal of Biology, Agriculture and Healthcare*, 5 (8): 67-71.
- Nweke, F.I., Spender, D.S.C. and Lynam, J. (2002). The Cassava Transformation: Africa's Best Kept Secret. Michigan State University Press, East Lansing, Michigan, USA.
- Okigbo, R. N., Ramesh Putheti and Achus, i C. T. (2009a) Postharvest deterioration of cassava and its control using extracts of *Azadirachta Indica* and *Aframomum Melegueta*. *E-Journal of Chemistry*, 6 (4): 1274-1280.
- Okigbo, R.N., Eme, U.E., Aseidu, R., Ramesh, P. (2009b) Effect of crude extracts of *Allium sativum* Linn, *Cymbopogon citratus* C.D. Stapf and *Terminalia catappa* on rot causing fungi of *Dioscorea* species. *Nigerian Journal of Botany*, 22 (2): 359-369.
- Okigbo, R.N., Anukwuorji, C.A. and Igweonu, M.Z. (2014) Fungitoxic effect of *Azadirachta indica* and *Vernonia amygdalina* extracts on *Aspergillus niger* and *Fusarium*

- oxysporium; the Causal organisms of yam (*Dioscorea rotundata*) rot. *The Bioscientist*, 2 (1): 70-86.
- Okigbo, R.N., Opara, P.U. and Anuagasi, C.L. (2015) Efficacy of extracts of water yam (*Dioscorea alata*) and aerial yam (*Dioscorea bulbifera*) peels in the control of white yam (*Dioscorea rotundata*) rot. *Journal of Agricultural Technology*, 11 (8): 1823-1842.
- Onyeka, T.J. (2002). Cassava root rot fungi in Nigeria; variability in *Botryodiplodia theobromae* isolates and evaluation of cassava germplasm for root rot resistance. PhD thesis, University of Ibadan.
- Onyeka, T. J., Owolade, O. F., Ogunjobi, A. A., Dixon, A. G. O., Okechukwu, R., Bandyopadhyay, R., and Bamkefa, B. (2008) Prevalence and severity of bacterial blight and anthracnose diseases of cassava in different agroecological zones of Nigeria. *African Journal of Agricultural Research*, 3 (4): 297-304.
- Philips, T.P, Taylor, D.S., Sanni, L. and Akoroda, M. (2006) A cassava industrial revolution in Nigeria; the potential for a new crop. IITA, Ibadan, Nigeria. IFAD, FAO of the United Nations, Rome, Italy.
- Plucknett, D.L., Philips, T. P. and Kagbo, R. B. (2003) The global cassava strategy for cassava: Transforming a traditional tropical root crop. The global cassava development strategy and implementation plan, Report of the FAO/WHO Expert Consultation Rome, 1: pp. 17-21.
- Sangoyomi, T.E., Asiedu, R., and Ekpo, E.J.A. (2010) Effects of ten plant extracts on mycelial growth and conidial production of four fungi associated with yam tuber rot, *AJRTC*, 8 (1): 24-30.
- Sangoyomi, T.E. (2004) Postharvest fungal deterioration of yam (*Dioscorea rotundata* poir) and its control. Ph.D thesis IITA, Ibadan Nigeria, pp. 179.
- Suleiman, M.N. (2010) Fungitoxic activity of neem and pawpaw leaves extracts of *Alternaria solani* causal organism of yam rots. *Advanced Environmental Biology*, 4 (2): 159-161.
- Taiwo, K. A. (2006). Utilization potentials of cassava in Nigeria: The domestic and industrial products. *Food Reviews International*, 22 (1): 29-42.
- Umana, E. J., Markson, A. A., Udo, S.E., Okon, E. I. and Akwaji, P. I. (2016) Control of postharvest rot of sweet orange (*Citrus sinensis*) with extracts of plant origin. *Nig. J. of Biop.* 1 (1): 88-93.