

Original Research Article

PREVALENCE AND DETERMINANTS OF MALNUTRITION IN UNDER FIVE CHILDREN IN **SELECTED HEALTH FACILITIES IN EGUME**, KOGI STATE, NIGERIA.

ABSTRACT

Aims: This study was conducted to determine the prevalence and determinants of malnutrition in under five children in selected health facilities in Egume, Dekina LGA, Kogi State.

Study Design: This was a **cross-sectional descriptive study with historic data** that assessed the Prevalence and determinants of malnutrition of under-five children **in selected health facilities in** Egume Dekina Local Government Area, Kogi State.

Study Area: The study was conducted at General Hospital and St. Luke clinic in Egume, Dekina LGA, Kogi State.

Methodology: The study focused on Mothers and their under five children who were treated as in- or out-patients at the General Hospital and St. luke clinic Egume within the past five years, Interview was conducted with the aid of a validated, questionnaire, after obtaining informed consent from mothers and ethical clearance from the Health Research Ethics Committee, Kogi State Ministry of Health. Data was obtained through reviews of available health records. Anthropometric measurement was carried out following standard procedures.

Results: The study showed that 21.3 %, 25.6 %, 47.4 % and 5.7 % of the sampled children were mildly undernourished, moderately undernourished, severely undernourished and healthy respectively.

Conclusion: The present study further revealed that a child's age and gender, mother's education and nutrition and other health variables such as clean water and toilet are determinants of child malnutrition.

Keywords: Malnutrition, under nutrition, Children, Women, under five, Prevalence.

1. INTRODUCTION

Malnutrition is a condition that results from eating a diet in which one or more nutrients are either not enough or are too much such that the diet causes health problems (1). **Insufficient intake of energy and nutrients results in under nutrition while excessive intake of nutrients results in over nutrition.** Adequate nutrition is essential in early childhood to ensure healthy growth, proper organ formation and function, a strong immune system, neurological and cognitive development. Economic growth and human development require well nourished populations **that could** learn new skills, think critically and contribute to their communities. Child malnutrition impacts cognitive function and contributes to poverty through impeding individuals' ability to lead productive lives. In addition, it is estimated that more than one-third of under-five deaths are attributable to under nutrition (2, 3). Children are prone to diseases, infection and malnutrition because their immune system is not fully developed. This phase of life is critical, hence the need for sufficient nutrients that will aid in their growth and development. (4)

According to the Nigeria Demographic and Health Survey (5), the percentage of stunting among Nigerian children under-five years declined from 41 % in 2008 to 37 % in 2013. However, the percentage of children who are wasted (Low weight for height) increased from 14 % in 2008 to 18 % in 2013. Under nutrition is more common in developing countries aside under-five children, certain groups have higher rates of under nutrition including women who are pregnant or breast feeding and the elderly under nutrition becomes more common due to physical, psychological and social factors (6).

For a child to be properly fed, to avoid malnutrition he/she must absorb or consume the right quality and quantity of food in the right number of times a day. Children 6 -to- 25 months of age should receive foods from the four main food groups which would provide the appropriate nutrients including fruits, vegetables, eggs, milk, fish and meat. Children should receive solid, semi-solid, or soft foods two –to – four times daily (7).

2. MATERIALS AND METHOD

2.1 Study design and duration

The study design was a cross-sectional descriptive study with historic data that assessed the Prevalence and determinant of malnutrition of under-five children in selected health facilities in Egume Dekina LGA, Kogi State, from January, 2012 to December, 2017.

2.2 Study Area

This study was conducted at General Hospital and St. Luke clinic in Egume, Dekina LGA, Kogi State. Egume, the headquarters of Okura district is an important town in Dekina Local Government area Kogi state, Nigeria. Its geographical coordinates are 7° 29' 0" North, 7° 12' 0" East and its original name (with diacritics) is Egume. The town is almost in the centre of Igala land. Egume has only one ward.

2.3 Study Population

Records of under five children who were treated as in-patient and out-patient and discharged in the past five years at the general Hospital Egume, Dekina LGA, Kogi State. Also, mothers with under five children who attend the St. Luke hospital clinic in Egume within the sample size participated in the study.

2.4 Inclusion and Exclusion criteria

Eligible mothers of under-five children who are willing to participate and give their consent were included in the study. Unwilling mothers with under-five children that are sick and requiring hospitalization were excluded from the study. Under-five children and their mothers on visit were also excluded.

2.5 Sample Size Determination

A minimum sample size was determined according to methods by Charan and Biswas (8).

$$n = \frac{Z^2 p q}{d^2}$$

Where:

n = Minimum sample size

Z = Standard deviation at 95% confidence interval = 1.96

p = Anticipated proportion/Prevalence Rate of Malnutrition = %

q = Complementary probability = 1 – p

d = Error margin/tolerance = 5% = 0.05

In Kogi state, a prevalence of 10 % was reported (9) . Therefore, at 10 % prevalence of malnutrition, using 5 % precision at 95 % confidence interval, the minimum sample size n for this study was calculated as 138. Therefore, 138 patients were selected for the study.

Sampling

The sampling method used was a systematic sampling; of the equal – probability modality. This sampling interval was elucidated using the formula:

$$K = \frac{N}{n}$$

Where:

K = sampling interval by which every Kth element/subject was selected from the sampling frame.

N = population size of patients = 556

n = sample size = 138 patients

K = $556/138 = 4$

Therefore, within 3 weeks of the research, every 4th patient was recruited from the entire sampling frame of 556 patients to comprise the 138 patients as sample size, given the logical homogeneous composition of the population.

2.6 Ethical Clearance and Consent Form

Ethical clearance was obtained from the Health Research Ethics Committee, Kogi State Ministry of Health in accordance with the code of ethics on human experimentation drafted by the World Medical Association in 1964.

Informed consent for inclusion into this study was obtained from the mothers with under five children who attend the selected health facilities to participate in the study using standard protocol.

2.7 Data collection.

Tools used for the data collection include; infantometer, height board, questionnaire, weighing scale, etc. Questionnaire was administered to respondent. Interview was conducted with validated questionnaire after obtaining informed consent from mothers with children under the age of five years. The questionnaire was pre-tested by administering the questions to a small number of representatives, who were mothers with under five children at both health facilities before the survey.

Relevant data on the factors related to malnutrition was obtained with the help of the schedule and review of any available relevant health records. For assessing the nutritional status of the subjects, anthropometric measurements were carried out following standard operating procedures. The data collected include weight, recumbent length (if the child is not able to stand without support), and height for children above 2 years.

2.8 Anthropometry

2.8.1 Age determination

Age was determined using available records or using local calendars.

2.8.2 Weight determination

Weight was measured to the nearest 0.1 kg using UNICEF Seca weighing scale with subjects in minimal clothing. The scale was occasionally calibrated using validated standard weights to avoid faulty reading.

2.8.3 Length/height determination

Height was measured using a length/height board, with the participant standing on a firm/level surface and it was measured to the nearest 0.1 cm. recumbent length was measured using an infantometer. Each measurement was done twice, and the mean of the two readings were recorded.

2.9 Data Analysis

Statistical analysis was performed using statistical package for social sciences (SPSS) 20.0. Results obtained were expressed as frequencies and percentages. The data obtained from questionnaire were subjected to descriptive statistics.

3. RESULTS

3.1 Socio-demographic status of mothers attending the selected health facilities in Egume, Kogi state.

The socio-demographic status of mothers attending St. Luke clinic and General hospital, Egume is presented in table 1. The results revealed that the age range between 31-40 (64 %) participated more in the study. 80 % of the mothers are married. 60.1 % are Igalas. A high percentage of the mothers (62.3 %) had senior secondary school education. Also more of the mothers are traders (56 %). 44.2 % of the mothers studied had their monthly income between N5, 000 - N14, 000.

Table 1: Socio-demographic status of mothers.

Characteristics	Group	Frequency (F)	Percentage (%)
Age	≤ 20	0	0
	21 – 30	33	24.0
	31 – 40	88	64.0
	>40	17	12.3
Marital Status	Single	22	16.0
	Married	110	80.0
	Divorced	3	2.2
	Widow	3	2.2
Ethnicity	Igala	83	60.1
	Hausa	5	3.6
	Yoruba	11	8.0
	Igbo	25	18.1
	Others	14	10.1
Education	Primary	14	10.1
	Junior Secondary	17	12.3
	Senior Secondary	86	62.3
	Tertiary Education	19	13.7
	No Formal Education	2	1.4
Occupation	Civil Servant	11	8.0
	Trader	77	56.0

	Farmer	17	12.3
	Artisan	8	5.7
	Full time house wife	16	12.0
	Others	9	6.50
Monthly Income	< N5,000	22	16.0
	N5,000 - N14,000	61	44.2
	N15,000 - N24,000	39	28.3
	N25,000 - N34,000	5	3.6
	N35,000 - N44,000	0	0
	N45,000 - N54,000	8	5.7
	≥ N55,000	3	2.2

N: Naira

3.2. Household care resources of mothers attending the selected health facilities in Egume, Kogi state.

The household care resources of mothers attending St. Luke clinic and General hospital Egume as shown in table 2 revealed that 52.2 % of the population use wood as source of heat for cooking. Main source of drinking water for 50 % is borehole. More than half (52.2 %) of the mothers use bush as their primary method of refuse disposal. 41.3 % make use of pit latrine. 50 % have no means of household food storage.

Table 2: Household care resources of mothers

Characteristics	Group	Frequency (F)	Percentage (%)
Main Source of Cooking	Wood	72	52.2
	Kerosene	47	34.1
	Gas	19	14.0
Main Source of Drinking Water	Public Tap	39	28.3
	Public well	5	3.6
	Private well	5	3.6
	River/Stream	17	12.3
	Borehole water	69	50.0
	Rain water	3	2.2
Primary Method of Refuse Disposal	Bush	72	52.2
	Refuse dump	61	44.2
	Others	5	3.6
Main Type of Toilet	Bush	28	20.3
	Pit latrine	57	41.3
	VIP latrine	14	10.1
	Water System	39	28.3

Household Food Storage Method	Refrigerator	22	16.0
	No means of storage	69	50.0
	Others	47	34.1

3.3: Child information and Hygiene Practices of mothers attending the selected health facilities in Egume, Kogi State.

Child information and Hygiene Practices of mothers attending General Hospital and St. Luke's clinic , Egume as presented in table 3 showed that they were more male (54.3 %) participants in the study than females (45.6 %). 68.1 % of the mothers wash their hands regularly with soap and water. Also 38 % of the mothers wash their fruits before eating it with the skin, also 16 % of mothers pre-chew food before giving to their child.

Table 3: Child information and Hygiene Practices of mothers.

Characteristics	Group	Frequency (F)	Percentage (%)
Sex of Child	Male	75	54.3
	Female	63	45.6
Age of Child (months)	≤ 1	14	10.1
	1 – 12	97	70.3
	13 – 24	5	3.6
	25 – 36	8	5.7
	37 – 48	6	4.3
	49 – 60	8	5.7
Number of siblings	<2	68	49.3
	>2	70	50.7
Hygiene Practices in Food Preparation			
Do you wash your hand with soap and water regularly?	Yes	94	68.1
	No	44	32.0
Do you wash fruits before eating with the skin	Always	52	38.0
	Sometimes	75	54.3
	Never	11	8.0
Do you masticate/ pre-chew foods before giving to your child?	Always	22	16.0
	Sometimes	58	42.0
	Never	58	42.0

3.4: Breastfeeding Knowledge and Practices of mothers attending the selected health facilities in Egume, Kogi State.

The Breastfeeding Knowledge and Practices of mothers attending General hospital and St. Luke's clinic, Egume, Kogi State as presented in table 4 showed that majority of the mothers (100 %) breastfed their children. 13.0 % of the study population breastfeed their children for the duration of 19 - 24 months. 88% introduced colostrums at birth. 60.1 % of the mothers initiated breast milk immediately after delivery. 70.3 % of the mothers breastfeed their child exclusively. 66 % of mothers used Guinea corn gruel as complimentary food. 56 % of the mothers introduced their children to complementary food at the age of 4 - 6 months, while 68.1 % of mothers feed children complementary food with aid of spoon and cup. 70.1 % of mothers prepared and stored their complementary food in warmers.

Table 4: Breastfeeding Knowledge and Practices of mothers.

Characteristics	Group	Frequency (F)	Percentage (%)
Do you breastfeed your child?	Yes	138	100.0
	No	0	0
Duration of breast feeding (months)	≤6	5	3.6
	7 – 12	0	0
	13 – 18	115	83.3
	19 – 24	18	13.0
Breastfeeding Initiation	Immediately after delivery	83	60.1
	Within the first 12 hours	25	18.1
	Within the first 24 day	22	16.0
	Within the first 2 days	5	3.6
	Others	3	2.2
Colostrum	Fed the Child	121	88.0
	Discarded	17	12.3
Breastfeeding frequency (per day)	3 – 5 times	3	2.2
	6 - 9 times	5	3.6
	>10 times	3	2.2
	Breastfed on demand	55	40.0
	Others	72	52.2
Exclusive breastfeeding	Yes	97	70.3
	No	41	30.0
Complementary foods introduced	Commercial	47	34.0
	Guinea corn gruel	91	66.0

Time of introduction of complementary foods	1 – 3 months	31	22.4
	4 – 6 months	77	56.0
	Above 6 months	30	22.0
Mode of feeding complementary foods	Spoon and cup	94	68.1
	Feeding bottles	44	32.0
	Others		
Preparation of complementary foods	On demand	41	30.0
	Prepared and stored in warmers	97	70.1

3.5: Nutritional Status of under-five children based on Weight for Age (WFA) between January 2012 to December 2017 at the selected health facilities in Egume, Kogi State.

Nutritional status of under-five children based on weight for age (WFA) between January 2012 to December 2017 at General hospital and St.Luke's clinic, Egume, Kogi State as presented in figure 1 shows that 11.6 % of the male children were mildly underweight, 15.4 % where moderately underweight, 25.8 % where severely underweight, and 1.9 % were healthy. It also shows that 9.7 % of females were mildly underweight, 10.2 % where moderately underweight, 21.6 % where severely underweight, and 3.8 % were healthy with the standard Z-scores <-1 to>-2, <-2 to>-3, <-3, 0 to +1 respectively.




Figure 1: Nutritional Status of under-five children based on Weight for Age (WFA) between January 2012 to December 2017 at General Hospital and St. Lukes clinic, Egume, Dekina, Kogi State.

Key: Mild = (< -1 to > -2 Z score)
 Moderate = (< -2 to > -3 Z score)
 Severe = (< -3 Z score)
 Healthy = (0 to $+1$ Z score)

4. DISCUSSION

Results from the analyses confirmed that child's variables (age and gender), mother's variables (education and nutrition), feeding practices and health variables (clean water and toilet) were the determinants of child malnutrition.

In this study, 64.0 % of the *mothers* were between the ages of 31 and 40 years, 80 % of the mothers were married and 60.1 % of them were from the Igala tribe.

The women see business as self-employment with quick results that could afford them minimum assets for marriage or for sending funds back home to their families. This present study revealed that few mothers (1.4 %) had no formal education, 62.3 % had secondary education. Several studies have shown that a mother's education is associated with good nutritional practices, most especially under-five child nutrition (10, 11, 12). Formal education of mothers directly transfers health knowledge to future mothers (13). Children born to educated women suffer less from under nutrition which usually manifests as underweight, wasting and stunting in children. Maternal education has been associated with nutrition outcomes among children in various settings (14). The literacy and numeracy skills that women acquire in school enhance their ability to recognize illness and seek treatment for their children. Mother's occupation is one of the indicators for access to adequate food supplies, use of health services, availability of improved water sources, and sanitation facilities which are prime determinants of child nutritional status (15). This study showed that few of the mothers are civil servant (8 %) and majority of them are traders (56 %). Weight-for-age is a composite index of weight-for-height and height-for-age, and does not distinguish between acute malnutrition (wasting) and chronic malnutrition (stunting). A child can be underweight for his age because he is stunted, because he is wasted, or both. Weight-for-age is a good

overall indicator of a population's nutritional health (16). The observed prevalence of under nutrition in this study could be due to the occupation of the care givers as they did not apparently appear constrained to cater for their children. Indeed, maternal occupation and educational characteristics determines child care practices as a correlation exists between height for age and the number of hours mothers worked outside the home (17).

A high prevalence sometimes reflects the low socio-demographic status of the mothers which affects the dietary intake. In this study, the socio-demographic status of the mothers revealed that few of the mothers (8%) were civil servants, although only 11.5 % earn a monthly income above N25 000, a greater percentage (60.2 %) earn N14000 or less. Such a relatively low income is likely to affect the nutritional status of subjects in such homes considering the cost of living in Egume. Poor breastfeeding and complementary feeding practices are widespread. Globally, it is estimated that only 34.8 % of infants are exclusively breastfed for the first 6 months of life, the majority receiving some other food or fluid in the early months (18). Complementary foods are often introduced too early or too late and are often nutritionally inadequate and unsafe (18). This contradicts the results from this study which indicate that majority of the mothers (70.3 %) breast fed exclusively. also all the mothers studied (100 %) breastfed their children. while 56 % of them introduced complementary food between 4 - 6 months. As a global public health recommendation, infants should be exclusively breastfed for the first 6 months of life to achieve optimal growth, development and health. The high percentage of exclusive breastfeeding observed was as a result of increased awareness and campaign. The present study further revealed that child's variables (age and gender), mother's variables (education and nutrition) and health variables (clean water and toilet) were the significant determinants of child malnutrition. Mother's education and body mass index are significant due to the fact that child nutrition will improve with increased mother's education and nutrition. Availability of clean water and toilet also reduces the possibility of infection. The cross sectional study of Nutritional status of under-five children studied showed that the case of severe underweight was high with 47.4 % of which male was 25.8 % and female 21.6 %. The significant determinants of underweight include, gender of child, education of mother, body mass index of mother and access to clean water. The gender variable revealed that male children are more likely to be underweight, compared to their female counterparts. The differences observed across studies in regard to the relationship between gender and the child's nutritional status may indicate that gender is not a biological factor in malnutrition, but that there may be other gender-related social or cultural factors that affect a child's growth. The results from this study are consistent with those from the study by Rayhan and Khan (19) on factors causing malnutrition among under-five children in Bangladesh that showed that low birth weight was positively associated with child underweight. This study revealed a significant association between child illness and child underweight, consistent with the findings of a study by Turyashemererwa et al (20), of child underweight. Also the finding that males were more underweight and wasted than females agrees with the study by Ozor *et al.*,(21). But disagrees with the findings of Banerjee *et al.*, (22), Harishankar *et al.*, (23) and Bhalani *et al.*,(24).

CONCLUSION

Results from the analysis confirm that child's variables (age and gender), mother's variables (education and nutrition), feeding practices and health variables (clean water and toilet) were the significant determinants of child malnutrition. To a large extent, women are responsible for feeding and caring for young children. The quality of feeding and care given to the children would reflect the level of education of the mother when other factors are fixed. This then gives a focus to policymakers in the designing of strategies aimed at combating malnutrition among children below five years.

REFERENCES

1. United Nations Children's fund. Facts for life. Newyork. pp.61 and 75. 2010; ISBN 978-92-806-4466-1.
2. Liu, L, Johnson, H.L, Cousens, S, Perin, J, Scott, S, Lawn, J.E, Rudan, I, Campbell, H, Cibulskis, R, Li, M, Mathers, C, Black, R.E. Global, regional and national causes of child mortality: an updated systematic analysis for 2010 with time trends since 2000. Lancet. 2012; **379**: 2151-61.
3. Black, R.E, Allen, H, Bhutta, Z.A, Caulfield, L.E, de Onis, M, Ezzati, M, Mathers, C, Rivera, J (2008). Maternal and child undernutrition: global and regional exposures and health consequences. Lancet. 2008; **371**: 243-60.
4. United Nations Children's fund. The state of the world's children. 2009.
5. National population commission. National demographic and health Survey, 2013. ICF Micro. 2014.
6. Powell, J, Nadia, I, Ronnie, J, Shirley, L. The characteristics and motivations of online health information seekers: cross-sectional survey and qualitative interview study. Journal of medical internet research. 2011; **13**(1).
7. WHO. The child growth standards. Available from <http://www.who.int/childgrowth/standard/en/>. 2004.
8. Charan, J and Biswas, T. How to calculate sample size for different study designs in medical research. Indian journal of psychological medicine. 2013; **35**(2): 121.
9. World bank. Multiple Indicator cluster survey (MICS). 2011
10. Babatunde, R.O. and Qaim, M. Impact of Off-farm Income on Food Security and Nutrition in Nigeria. Food Policy. 2010; **35**: 303-311.
11. Webb P. and Block, S. Nutritional Information and Formal Schooling as Inputs to Child Nutrition. Economic Development and Cultural Change, 2004; **55**: 801-820.
12. Shrimpton, R., Victoria, C.G., Onis, M.de., Lima, R.C., Blossner, M., and Clugston, G. Worldwide Timing of growth faltering: Implications for nutritional interventions. Paediatrics. 2001; **107**: 75-81.
13. Desai S and Alva S. Maternal education and child health: is there a strong causal relationship? Demography. 1998; **35**(1):71-81.
14. Abuya, B.A., Onsomu, E.O., Kimani, J.K., Moore, D. Influence of maternal education on child immunization and stunting in Kenya. Maternal and child health journal. 2011; **15**(8): 1389-1399.
15. Marie, T.R., Carol, E, L., Margaret, A.K., Klemesu, D, Daniel, M., Saul, S.M. Good care practices can mitigate the negative effects of poverty and low maternal schooling on children's nutritional status: evidence from Accra. World development. 1999; **27** (11): 1993-2009.
16. Ogden, C.L., Carroll, M.D., Kit, B.K., Flegal, K.M. Prevalence of childhood and adult obesity in the United states, 2011-2012. Jama. 2014; **311**(8): 806-814.
17. Roshita, A., Schubert, E, Whittaker, M (2012). Child care and feeding practices of urban middle class working and non working Indonesian mothers: a qualitative study of the socio- economic and cultural environment. Maternal and child nutrition. 2012; **8**(3): 299-314.

18. Caetano, M.C., Ortiz Ortiz, T.T., Lopes da Siva, S.G., Souza, F.I, Sani, R.O. Complementary feeding: inappropriate practices in infants. *Jornal de pediatria*. 2010; **86** (3): 196-201.
19. Rayhan, I and Khan, SH. Factors causing malnutrition among under five children in Bangladesh. *Pakistan Journal of Nutrition*. 2006; **5**(6): 558 - 562.
20. Turyashemerwa, F.M., Kikafunda, J.K., Agaba, E. Prevalence of early childhood malnutrition and influencing factors in peri urban areas of Kabarole district, Western Ugandan. *African journal of Food, Agriculture, Nutrition and Development*. 2009; **9** (4).
21. Ozor, M.O., Iyamu, O.A., Osifo, U.C. Prevalence of under nutrition among under five year children in Ekpoma, Edo-Nigeria. *International journal of community Research*. 2014; **3** (1): 34-38.
22. Banerjee, B and Bandyopadhyay, L. Gender difference in nutritional status. *Indian paediatrics*. 2005; **42**: 400.
23. Harishankar, Dwivedi, S., Dabral, S.B., Walia, D.K. Nutritional status of children under 6 years of age. *Indian Journal of Preventive and social medicine*. 2004; **3**(4): 156-62.
24. Bhalani, K.D, Kotecha, P.V. Nutritional status and gender differences in the children of less than 5 years of age attending ICDS Anganawad's in vadodara city. *Indian journal of community medicine*. 2002; **27**: 124-9.

UNDER PEER REVIEW