2 3

4

- 5
- 6

7 8

# ABSTRACT

**Introduction:** The ABO blood group system is unique in immunology and haematology because it is the only system in which antibodies are consistently and predictably present in the serum of normal individuals whose red cells lack the antigens. Several epidemiological studies have reported that the distribution of different ABO blood groups vary markedly among the populations of different geographical areas reflecting racial differences

Anthropometric Indices among Young Adults of Yoruba

**ABO Blood Group system: Its Association with** 

**Ethnicity** 

Aims: The aim of this study was to determine the association between ABO blood group and anthropometric indices among apparently healthy young adults of Yoruba ethnicity.

**Study design:** This is a cross-sectional descriptive study.

**Place and Duration of Study:** The study was carried out at Department of Physiological Sciences and University Health Centre, Obafemi Awolowo University, Ile-Ife between May 2016 to June 2017.

**Methodology:** Eighty apparently healthy young adults who were purposely selected participated in the study after screening for the ABO blood groups following the standard protocol. They were divided into four equal groups; 20 in each of blood groups A, B, AB and O who were age- and sex - matched. The weight (kg) and height (cm) of the participants were measured following standard protocol. Body mass index (BMI) and body surface area (BSA) were estimated from weight and height using Quetelex and Mosteller formulae respectively. Chi-square was used to compare grouped data while the comparison of means of parameters among the four blood groups was done using Analysis of Variance (ANOVA). A p-value of < 0.05 was taken statistically significant.

**Results:** Out of the total 80 participants, 15% were males. The mean  $\pm$  SD of height(m) of participants with blood groups A, B, O and AB were 1.65  $\pm$  0.06, 1.63  $\pm$  0.08, 1.62  $\pm$  0.08 and 1.63  $\pm$  0.08 (F= 0.349, p= 0.790) respectively. The mean  $\pm$  SD of weight(kg) of the participants with blood group A,B, O and AB were 56.15  $\pm$  8.71, 56.00  $\pm$  11.21, 57.10  $\pm$  12.73 and 58.05  $\pm$  10.35 (F= 0.154, p= 0.927) respectively. The mean  $\pm$  SD of BMI in kg/m<sup>2</sup> for blood groups A, B, O and AB were 20.74  $\pm$  3.22, 20.86  $\pm$  2.91, 21.79  $\pm$  5.10 and 21.91  $\pm$  4.21 (F= 0.472, p= 0.703) respectively while the mean  $\pm$  SD of BSA(m<sup>2</sup>) for blood group A, B, O and AB were 1.60  $\pm$  0.12, 1.59  $\pm$  0.19, 1.60  $\pm$  0.18 and 1.62  $\pm$  0.15 (F= 0.098, p= 0.961) respectively.

**Conclusion:** ABO Blood group system may be predisposed to metabolic syndrome due to the higher mean of weight, BSA and BMI found in this group.

9 10

Keywords: ABO blood groups, Anthropometry, Apparently healthy, Young adults, Ethnicity

11

# 12 1. INTRODUCTION

13 The ABO blood group system was the first and most important human blood group system [1]. It was 14 credited to be discovered by Karl Landsteiner in 1900 [2]. The ABO blood group system is unique in 15 immunology and haematology because it is the only system in which antibodies are consistently and 16 predictably present in the serum of normal individuals whose red cells lack the antigens [3]. The 17 antibody production becomes detectable at three to six months of age [4]. Several epidemiological 18 studies have reported that the distribution of different ABO blood groups varies markedly among the 19 populations of different geographical areas reflecting racial differences [5-7]. Human blood groups are 20 genetically determined antigens expressed on the surface of red blood cells [8]. The antigens may 21 also be found in body fluids (urine, saliva and amniotic fluid) of secretors [9]. Blood groups may also 22 be expressed on organ cellular surfaces. The association between ABO blood groups and disease 23 susceptibility has generated a lot of interest [10-17]. Its association with anthropometric indices 24 among young adults is rarely studied in Nigeria. Anthropometry is the scientific study of the measurements and proportions of the human body. It can also be explained as a branch of 25 26 anthropology dealing with measurement of the human body to determine differences in individuals 27 and groups; especially for the purpose of classification and comparison [18]. Body mass index (BMI) 28 is regarded as most popular among the anthropometric indices. It is an internationally accepted index 29 for assessing obesity [19,20]. BMI is a measure of weight adjusted for height, calculated as weight in kilograms divided by the square of height in meters [20]. Age and ethnicity had been reported to play 30 a significant role in the anthropometric dimensions [21,22]. BMI as a trait is determined not only by 31 32 environmental factors but also largely by interaction with inherited genetic factors [23]. Several studies 33 have indicated that genetic factors account for a substantial portion of variation in human adiposity 34 [24-26]. The ABO blood group system is known to be one of such genetic make-up of an individual 35 that may provide much valuable information for early detection of vulnerable groups for abnormal BMI such as obesity [27]. Data is sparsed with regards to the association between ABO blood group and 36 37 anthropometric indices among apparently healthy young adults of Yoruba ethnicity in Nigeria, hence 38 this study.

## 40 2. MATERIAL AND METHODS

This study was carried out at Obafemi Awolowo University, Ile-Ife. It was a cross-sectional descriptive study involving apparently healthy young adults between the ages of 18 and 40 years (inclusive) of Yoruba ethnicity. The target population were the students of Obafemi Awolowo University community, Ile-Ife, Nigeria. Ethical approval was obtained from the Institute of Public Health of the Obafemi Awolowo University, Ile-Ife. Eighty participants were purposively selected using Estimated sample size was be determined using STATA statistical software (version 10) formula for estimation of sample size studies;

- $N = \frac{4 \times \sigma^2 (Z_{crit} + Z_{pwr})^2}{D^2}$
- 49 **D**<sup>2</sup> 50 **N** = Total number of sample size,

51  $\sigma = 0.05$ ,

39

48

57 58

- 52 Z<sub>crit</sub> = Standard normal deviate corresponding to selected significance criteria. It is a constant 53 factor and it equals to 1.960
- 54  $Z_{pwr}$  = Standard normal deviate correspondence to selected statistical power. It is a constant 55 factor and equal to 1.282,
- 56 D =the effect size of 4% = 0.04.
  - $N = \frac{4 \times (0.05)^2 (1.960 + 1.282)^2}{(1.960 + 1.282)^2} = 65$

 $(0.04)^2$ 

59 N was rounded up to 80 which was equally divided among the four groups

Blood samples were obtained from a superficial cubital vein (2ml) from the participants at resting 60 position after cleaning with 75% ethyl alcohol using 5ml syringe and the sample was dispensed into 61 Sodium EDTA anticoagulant bottle. All sample collections followed standard procedure. A drop of each 62 antisera A, B and AB were placed on a clean white tile in three different places marked X, Y and Z. 63 Drops of blood were added to drops of antisera at points X, Y and Z respectively and mixed with the 64 65 aid of glass rods. Then, the mixture was rocked gently for 60 seconds to observe for agglutination. The 66 results of agglutination were recorded immediately after mixing. The tests were done in duplicates to ensure the validity of results. The agglutinations in X & Z were considered as blood group A and 67 68 agglutinations in Y & Z were considered as blood group B. The agglutinations at points X, Y and Z 69 were considered as group AB. If no agglutination at points X, Y and Z, the sample was considered as 70 blood group O. Agglutination at point X or Y or Z were considered invalid and the test was repeated. Heights and weight of all participants were measured to the nearest 1 cm and 0.5kg respectively using 71 72 ZT120 Health Scale. The body mass index was calculated from height in metres and weight in kg [28] and body surface area (BSA) was calculated from the weight (kg) and height (cm) using Mosteller 73 formula, BSA= [(weight in kg × height in cm) / 3600]<sup>1/2</sup> [29]. The data were analyzed by the aids of IBM 74 75 version 20.0 software using descriptive statistics and cross-tabulations. Comparison of means of parameters among the four blood groups (A, B, AB and O) was done using Analysis of Variance 76 (ANOVA). Comparison of grouped data was done using chi-square. A p value of < 0.05 was taken as 77 78 statistically significant. 79

## 80 3. RESULTS

The mean  $\pm$  SD and range of age in years, weight in kg, height in metres, body surface area in metre square, body mass index in kg/m<sup>2</sup> were 21.50  $\pm$  3.52 (18-28), 56.83  $\pm$  10.67 (39-95), 1.63  $\pm$  0.07 (1.48-1.81), 1.60  $\pm$  0.14 (1.30  $\pm$ 2.08), 21.33  $\pm$  3.92 (15.42-38.54) respectively. This study showed that 12% were underweight, 66% had had normal weight, 10% had overweight and 3% had obesity (Figure 1). The highest and lowest mean height was found in blood groups A and O respectively. The highest mean weight, body surface area and body mass index were found in blood group AB. The
relationship between ABO blood group and anthropometric parameters were not statistically
significant (p-value > 0.05) as shown in table 1. The highest frequency of obesity was found in blood
group AB. The distribution of ABO blood group according to the BMI class (underweight, normal
weight, overweight and obesity) is as shown in figure 2.



Figure 2: Distribution of ABO blood group according to body mass index



Table 1: Relationship between ABO blood group and Anthropometric Parameters

Anthropometry	Blood Group	Parameters (Mean ± SD)	F	p-value
Weight (kg)	A	56.15 ± 8.71	0.154	0.927
	В	56.00 ± 11.21		
	0	57.10 ± 12.73		

	AB	58.05 ± 10.35		
Height (m)	A	1.65 ± 0.06		
	В	1.63 ± 0.08	0.349	0.790
	0	1.62 ± 0.08		
	AB	1.63 ± 0.08		
BSA (m <sup>2</sup> )	A	1.60 ± 0.12		
	В	1.59 ± 0.19	0.098	0.961
	0	1.60 ±0.18		
	AB	1.62 ± 0.15		
BMI (kg/m <sup>2</sup> )	A	20.74 ± 3.22		
	В	20.86 ± 2.91	0.472	0.703
	0	21.79 ± 5.10		
	AB	21.91 ± 4.21		

102 N=80. Statistically significant p-value- \*

103 SD- standard deviation, BSA- body surface area, BMI- body mass index

104

### 105 106 **4. DISCUSSION**

107 The mean weight (56.83 ± 10.67kg) obtained from this study was similar to 56.37 ± 6.53kg reported in 108 a study among young adults within the age range of 18-40years in India [30]. The mean height, 109 weight and BMI of the study population were similar to the earlier report of a study done within the 110 same environment [31-33]. It is also inconsistency with the report of the study done among the undergraduates of the University of Benin, Nigeria within the same age range [34]. The percentage of 111 112 the participants that were either overweight (10%) or obese (3%) was far lower than 33.9% and 113 25.4% respectively that was reported by Jafari et al in 2012 in adult Iran population between the ages 114 of 40-57 years [35]. The difference may be as a result of the difference in age and genetic make-up. 115 The mean body surface area gotten from the study is similar to the report of El Missiri et al 2016 in a 116 study conducted among young healthy Egyptian adults [36].

117 From this study, the highest mean height was found in blood group A which is in line with the report of 118 Chaitra et al in a study done among 120 young adults of Kasturba Medical College, Mangalore in 119 India within the age range of 18-22 years [37]. This is in contrast with blood group B reported by Kuar 120 in 2014 in a study done among post-menopausal females between 45 to 80 years of age living in 121 India [17]. This contrast may be due to the age difference among the participants. Highest mean BMI 122 was found in blood group AB which is supported by the studies of Ainee et al, Amela et al and Chaitra 123 et al in studies done among young adults [37-39]. The high BMI in this blood group may predispose 124 them to metabolic syndrome. This was in contrast with a study by Jafari et al which reported the 125 highest BMI in blood group A [35]. ABO blood group system did not have a significant effect on body 126 anthropometric parameters (height, weight, BSA and BMI). This was in support of the report of Jafari 127 et al in 2012 in a population-based study of 50,045 men and women who reside in rural and urban 128 areas of Golestan Province of Iran [35]. This was also in line with the report of Chaitra et al in 2014 129 and Amela et al in 2017 among other studies [37,39,40]. The lowest mean BMI was in blood group A, 130 this could be due to the preference of this blood group for fruits and vegetables as earlier reported by 131 Lam [41]. It was also found that blood group AB has the highest percentage of participants with

132 obesity (10%) when compared with other blood groups. This corroborates the findings shown in table 1.

133

134 135

#### **5. CONCLUSION** 136

137 The mean ± SD and range of age in years, weight in kg, height in metres, body surface area in metre 138 square, body mass index in kg/m<sup>2</sup> were 21.50  $\pm$  3.52 (18-28), 56.83  $\pm$  10.67 (39-95), 1.63  $\pm$  0.07 139 (1.48-1.81), 1.60 ± 0.14 (1.30 ±2.08), 21.33 ± 3.92 (15.42-38.54) respectively. This study showed that 140 12% were underweight, 66% had had normal weight, 10% had overweight and 3% had obesity. The 141 highest mean weight, body surface area and body mass index were found in blood group AB. 142 Though ABO blood group has no significant effect on body anthropometric parameters blood group 143 AB may be predisposed to metabolic syndrome due to the higher mean of weight, BSA and BMI 144 found in this group.

145

152

154 155

158

159 160

172

#### CONSENT (WHERE EVER APPLICABLE) 146

147 Written informed consent was obtained from each participant. 148

#### **ETHICAL APPROVAL** 149

150 Ethical approval for this study was obtained from the Institute of Public Health of the Obafemi 151 Awolowo University, Ile-Ife.

#### 153 REFERENCES

- 1. Seeley RR, Stephens TD, Tate P. Anatomy and Physiology. 4th Edition. The McGraw Hill Companies, Inc., USA, 1998; 1098,
- Watkins WM. The ABO blood group system: historical background. Transfus Med 2001; 11: 156 2 157 243-265
  - 3. Daniels GL (2002). Human Blood Groups. 2nd ed. Oxford, United Kingdom: Blackwell Science.
    - 4. Puneet KK. Blood transfusion in clinical practice, first edition, 2012, Copyright Intech.
- 161 5. Sana Ullah, Ahmad T. Distribution of ABO and Rh (D) Blood Groups in the Population of 162 District Dir Lower, Khyber Pakhtunkhwa Pakistan. World Applied Sci J. 2015; 33: 123-35.
- 163 6. Anees M, Mirza MS. Distribution of ABO Group alleles in Guirat region of Punjab, Pakistan. 164 Proc. Pak. Acad. Sci. 2005; 4: 233-238.
- 165 Bakare AA, Azeez MA, Agbolade JO. Gene frequencies of ABO and rhesus blood groups and 7. 166 haemoglobin variants in Ogbomoso, south-west Nigeria. African Journal of Biotechnology. 167 2006; 5(3): 224-229.
- 168 8. Mondal B, Maiti S, Biswas BK, Ghosh D, Paul S. Prevalence of hemoglobinopathy, ABO and 169 rhesus blood groups in rural areas of West Bengal, India. Journal of Research in Medical 170 Sciences. 2012; 17(8): 772-776. 171
  - 9. Sheryle AW. Immunohematology for medical laboratory Technician, first edition Delmar, Cengage Learning. 2010.
- 173 10. Kucevic I, Vasic J, Svorcan JZ, Radisic B. Evaluation of ABO blood groups distribution in 174 patients with diabetes mellitus Type 2 from Serbia. Serbian Journal of Experimental and 175 Clinical Research. 2013; 14(3):
- 176 11. Bener A, Yousafzai MT. The distribution of the ABO blood groups among diabetes mellitus 177 patients. Nigerian Journal of Clinical Practice. 2014; 17 (5): 565
- 178 12. Moinzadeh F, Najafabady GM and Toghiani. Type 2 diabetes mellitus and ABO/Rh blood 179 groups. Journal Research Medical Sciences. 2014; 19(4): 382
- 180 13. Wolpin BM, Kraft P, Gross M, Helzlsouer K, Bueno-de-Mesquita HB, Steplowski E, et al. Pancreatic cancer risk and ABO blood group alleles: results from the pancreatic cancer cohort 181 182 consortium. Cancer Research. 2010; 70: 1015-1023.
- 183 14. Xie J, Qureshi AA, Li Y, Han J. ABO blood group and incidence of skin cancer. PLoS One. 184 2010; 5(8): e11972.
- 185 15. Bhattacharyva S, Ganaraja B, Ramesh BM. Correlation between the blood groups, BMI and 186 prehypertension among medical students. J. Chinese Clinical Medicine. 2010: 51(2): 78-82 16. Nasreen AR. Blood groups and hypertension. J. Bagh Coll Dentistry. 2006; 18(2): 68-70 187
- 188 17. Kaur M. Association between ABO blood group and hypertension among post-menopausal 189 females of North India. 2014

- 18. Must A, Anderson SE. Body mass index in children and adolescents: considerations for
   population-based applications, International Journal of Obesity. 2006; 30: 590–594.
- Ajayi OI, Akinbo DB, Okafor AM. Correlation between Body Mass Index and Hematological Indices in Young Adult Nigerians with Different Hemoglobin Genotypes. Am. J. Biomed. Sci. 2017; 9(1): 38-46;
- Billewicz, WZ, Kemsley WFF, Thomson AM. Indices of adiposity. British Journal of Preventive
   and Social Medicine. 1962; *16: 183*
- 197 21. Lean MEJ, Han TS, Bush H, Anderson AS, Bradby H, Williams R. Ethnic differences in anthropometric and lifestyle measures related to coronary heart disease risk between South Asian, Italian and general population British women living in the west of Scotland. Int J Obesity. 2001; 25: 1800-1805.
  - 22. Yu-Cheng L, Wang JM, Wang EM. The comparison of anthropometric characteristics among four peoples in East Asia. Applied Ergonomics. 2004; 35: 173-178.

201 202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223 224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

- 23. Loos RJ, Bouchard C. Obesity--is it a genetic disorder? Journal of Internal Medicine. 2003; 254(5): 401-425.
- 24. Allison DB, Kaprio J, Korkeila M, Koskenvuo M, Neale M, Hayakawa K. The heritability of body mass index among an international sample of monozygotic twins reared apart. International Journal of Obesity and Related Metabolic Disorder. 1996; 20: 501–506.
- Bouchard C, Tremblay A, Després, JP, Nadeau A, Lupien PJ, Thériault G. The response to long-term overfeeding in identical twins. New England Journal of Medicine. 1990; 322: 1477– 1482.
  - 26. Price R A, Gottesman II. Body fat in identical twins reared apart: roles for genes and environment. Behavioural Genetics. 1991; 21: 1–7.
  - Speliotes EK, Willer CJ, Berndt SI, Monda KL, Thorleifsson G, Jackson AU, et al., Association analyses of 249,796 individuals reveal 18 new loci associated with body mass index. Nature Genetics. 2010. 42(11). 937-948.
    - 28. Keys A, Fidanza F, Karvonen MJ, Kimura N, Taylor HL. Indices of relative weight and obesity. Journal of Chronic Diseases. 1972; 25(6-7): 329-349.
    - 29. Mosteller RD. Simplified calculation of body surface area. New England Journal of Medicine. 1987; 317(17): 1098.
    - 30. Mahendrappa SK, Sathyanarayana TB. The normal variations in heart rate and QRS complex of healthy individuals in different age groups.\_International Journal of Research in Medical Sciences. 2014; 2(3): 861-865.
    - Johnson OE, Adedoyin RA, Awotidebe TO, Mbada CE, Otolorin OA, Owoseni I. Cardiovascular risk among Undergraduates in a Nigerian University. International Scholar Journal. 2013; 2(5): 85-89.
    - Ogunlade O, Asafa MA. Pattern and Prevalence of Underweight, Overweight and Obesity Among Young Adult Nigerians. American Journal of Biomedical and Life Sciences. 2015; 3: 12-15
    - 33. Ogunlade O, Asafa MA, Ayoka AO, Akintomide AO. Association between body anthropometric parameters and indices of cardiac autonomic function among apparently healthy young adults. Achieve of Medicine and Health Sciences. 2016; 4(1): 26-29.
    - Odili VU, Egor FO, Osarenmwinda IM, Obieche AO. Prevalence of Overweight and Obesity among Pharmacy Students in a University in Benin City, Nigeria. Tropical Journal of Pharmaceutical Research. 2014; 13(12): 2109-2114.
    - 35. Jafari E, Sebghatollahi V, Kolahdoozan S, Elahi E, Pourshams A. Body Mass Index and ABO Blood Groups among Different Ethnicities of the Golestan Cohort Study Subjects. Govaresh. 2012; 17(1): 50-54.
  - 36. El Missiri, AM, El-Meniawy KA, Sakr SA, Mohamed AS. Normal reference values of echocardiographic measurements in young Egyptian adults. The Egyptian Heart Journal. 2016; 68: 209-215.
- 37. Chaitra U, Bhagyalakshmi K, Vinodini N, Anupama N, Rekha DK, Nayanatara A, Shreedhara
  A, Sheila RP, Prashanth KC. Comparison of heart rate variability in different abo blood groups
  of young adults. International Journal of Medical Science and Public Health. 2014; 3(12):
  1466-1469.
- 38. Ainee A, Hussain S, Kauser T, Qureshi TM., Nadeem M, Rashid F. Studies on Comparison of
  Body Mass Index (BMI) of School Going Children Having Different Blood Groups (A, B, Ab
  and O) of Sargodha District. Pak J Nutri. 2014; 13(3): 164-167.

- Amela H, Edina H, Snježana H, Emir H, Aldijana A, Mirela H. Blood Group, Hypertension and
  Obesity in the student population of northeast Bosnia And Herzegovina. IFMBE Proceedings.
  2017; 62: 774-777.
- 40. Mascie-Taylor CG, Lasker GW. Lack of an association between ABO and Rh blood group polymorphisms and stature, body weight, and BMI in a cohort of British women. Hum Biol. 1990; 62: 573-576.
- 41. Lam M. Blood Type Diet. Dr Lam website.https://www.drlam.com/blood\_type\_diet/.

### 255 ABBREVIATIONS

ANOVA: Analysis of Variance; BMI: Body Mass Index; BSA: Body Surface Area; EDTA: Ethylene Diamine Tetraacetic Acid; WHO: World Health Organisation.