

Protective Effect of *Dorema glabrum* on induced Oxidative Stress by diazinon in

Hippocampus of Rat

Parisa Sistani, Homeira Hatami Nemati, Mina Adampourezare

Abstract

Introduction: Diazinon (DZN) administration produces lipid peroxidation as an indicator of oxidative stress in the brain. Some medicinal plants such as *Dorema glabrum* has antioxidant properties, so can be used as an antioxidant that may protect neurons from oxidative stress. The aim of present study was to investigate the effect of *D.glabrum* against DZN-induced oxidative stress in hippocampus.

Methods: Twenty-four adult male Wistar rats were used in this study. The rats randomly were divided into four groups including a control group, and two groups received different doses of *D.glabrum* (40 and 80 mg/kg) as pre-treatment for 21 days with DZN (100 mg/Kg) that was injected intraperitoneally (ip) in last day of *D. glabrum* usage, and one group received only DZN. Thiobarbituric acid reactive substances (TBARS), which are the indicators of lipid peroxidation, and the activities of antioxidant enzymes (glutathione peroxidase, superoxide dismutase and catalase) were determined in the rats' hippocampus.

Results: Administration of DZN significantly increased TBARS levels and superoxide dismutase activity and decreased glutathione peroxidase activity but there were no significant changes in catalase activity in the hippocampus. Combined *D.glabrum* and DZN treatment, caused a significant increase in glutathione peroxidase, a significant decrease of TBARS and a significant decrease in superoxide dismutase and again no significant changes in catalase activity in the rats' hippocampus when compared to the rats treated with DZN.

Conclusion: Our study demonstrated that *D.glabrum* had an amelioratory effect on oxidative stress induced by DZN.

Key words: Diazinon, Oxidative stress, *Dorema glabrum*, Hippocampus, male rat

Introduction:

Diazinon is a thiophosphorus organophosphate pesticide (OPs) which is used commonly as insecticide, nematicide and acaricide in soil, fruit, plants, vegetable crops in agriculture, urban pest control, and veterinary medicine [1]. DZN inhibit acetylcholinesterase activity, which is an important brain neurotransmitter, and cause accumulation of acetylcholine in the synapses [2]. Symptoms of acute DZN poisoning include headache, dizziness, nausea, tearing, and sweating. Some symptoms including headaches, blurred vision, and memory problems can last for months or years [3]. Low level repeated exposure to OPs causes inflammatory responses in cultured astrocytes [4] and upregulates

31 inflammatory cytokines in vivo [5] and inflammatory cytokines have induce significant impairment in spatial memory
32 [6]. DZN at high doses induces oxidative stress and the production of free radicals in rats by alteration of antioxidant
33 enzyme activity, depletion of glutathione S-transferase (GST), and lipid peroxidation [7]. Endogenous enzymatic and non-
34 enzymatic antioxidants are so important for the conversion of reactive oxygen species (ROS) to harmless metabolites as
35 well as to protect and restore normal cellular metabolism and functions [8].The brain is particularly vulnerable to
36 oxidative stress [2] because it metabolizes about 20% of total body oxygen and has a limited amount of antioxidant
37 capacity [9]. The hippocampus, which is a part of the limbic system, is a critical center for memory and learning
38 processes and plays an important role in forming and saving spatial memory. Reactive oxygen species (ROS) are
39 involved in several diseases including ischemic injury, aluminum toxicity, Parkinson's disease, Alzheimer's disease
40 and Down's syndrome all of which affect cognitive processes [11,12,13].

41 ROS can be detoxified by an enzyme defense system, containing superoxide dismutase (SOD), catalase (CAT), and
42 selenium-dependent glutathione peroxidase (GPx), or non-enzymatic systems by the scavenging action of reduced
43 glutathione, while organic peroxides can be detoxified by glutathione S-transferase (GST) [14].

44 Many insecticides have hydrophobic molecules that can bind extensively to biological membranes, especially to the
45 phospholipids bilayers [15].

46 *Dorema glabrum* belongs to the family of Umbelliferae distributing throughout the Mediterranean to Central
47 Asiaaltitudes [16]. The most of the active constituents of this plant are polar phenolic components, which shows
48 antioxidant activity and has many beneficial pharmacological effects [17]. In a previously work, the crude extract of the
49 plant demonstrated antioxidant activity and anti-lipidemic effects [18].

50 Members of the genus *Dorema* (Apiaceae) possess antispasmodic, carminative, expectorant, diaphoretic, mild diuretic,
51 emmenagogue, stimulant, vasodilator [19], antimicrobial and antifungal [20, 21,22] and hepatoprotector [23] properties
52 and are extensively used as a green vegetable or as a folk medicine for treatment of many diseases [24]. According to
53 the common folk believes of Azeri and Armenian people, *D.glabrum* can suppress different kinds of cancer.

54 The present study was designed to investigate the effect of DZN on enzymes of superoxide dismutase (SOD), catalase
55 (CAT), glutathione peroxidase (GPx) activities, total thiol (TSH) and thiobarbituric acid reactive substances (TBARS)
56 as an indicator of lipid peroxidation in the hippocampal areas of the rats brain as well as on antioxidant enzymes
57 activities and TBARS levels.

58

59 **Materials and Methods:**

60 **Plant Material:**

61 Seeds of *D. glabrum Fisch* were collected during the fruiting stage from slopes of Aras river; Jolfa, Eastern Azerbaijan,
62 Iran. Air dried and then powdered seeds were subjected to extraction by refluxing ethanol using a soxhlet (DGE). Then
63 the extract was dried by a rotary evaporator (Heidolph, Germany).

64

65 **Animals:**

66 Twenty-four adult male Wistar rats (weighting approximately 250-350 g) used in present study. Animals were obtained
67 from the animal house of the veterinary department University of Tabriz. The rats were kept under constant
68 temperature 20-22°C and 12/12h cycle of light and darkness and access to enough food and water throughout the
69 **experiment**. Ethics of working with laboratory animals have been followed during all procedures.

70

71 **Animal treatment schedule:**

72 The rats randomly **were divided** into four groups including group I (C): normal control rats, group II (DZN): received
73 DZN in single dose (100 mg/kg), group III (DGE 40): received (40 mg/kg) of DGE and (100 mg/kg) of DZN, group IV
74 (DGE 80): received DGE (80 mg/kg) and DZN (100 mg/kg). This two experimental groups **were received** different
75 doses of DGE (40 and 80 mg/kg) as pre-treatment for 21 days with DZN (100 mg/kg) that **was injected** intraperitoneally
76 in last day of DGE usage.

77

78 **Chemicals:**

79 DZN was applied from Aria chemistry Co. (Iran) containing 96% active ingredies. It was diluted with corn oil as DZN
80 solvent. Thiobarbituric acid (TBA), trichloroacetic acid (TCA), H₂O₂ (30%), ethylene diamino tetraacetic acid
81 (EDTA), Tris-HCl, 2,2'-dinitro-5,5'-dithio-dibenzoic acid (DTNB), ethanol of technical grade and the other chemicals
82 used in this study were procured from Merck Co. (Germany).

83

84 **Tissue preparation:**

85 Brains (hippocampus) were removed from the animals under ether anesthesia after 21 days of treatment and washed
86 with cold saline buffer. Then washed tissues were immediately stored at -80°C. To obtain the enzymatic extract, tissues
87 were homogenized in ice-cold KCl 1.15% to yield 10% (W/V) homogenate. Then the homogenates were centrifuged at
88 1000 rpm for 10 min at 4°C. The supernatants were separated and used for enzyme activity of SOD, CAT and GPx
89 which was expressed in international units per mg protein (IU/mg protein). Biomarkers for tissue damage were
90 measured using UV kinetics methodology and total protein was determined using bovine serum albumin (BSA) as
91 standard and the values were expressed as mg/dl.

92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116

Analytical procedures

Thiobarbituric acid reactive substances (TBARS) were measured as an index of lipid peroxidation by the method of Satho [25]. SOD was determined according to the method described by Ukeda [26]. CAT was measured by monitoring the decomposition of hydrogen peroxide, as described by Aebi [27]. GPx was evaluated by the method of Paglia and Valentine (21). Protein was measured by the method of Lowry [28] using bovine serum albumin as standard. Total thiol content (TSH) was measured in homogenate by the method of Hu [22].

Statistical analysis:

Statistical analysis was performed using one-way analysis of variance (ANOVA) followed by Tukey’s post hoc test (level of significance $P < 0.05$) using SPSS version 16, statistical program. The results are expressed as mean \pm SEM, and were obtained from at least 6 rats in each group. Statistical analysis was based on comparing the values between the DZN and control groups, while DZN-treated groups concomitantly with DGE were compared with their corresponding group of DZN-treated rats.

Results:

Lipid peroxidation:

Lipid peroxidation (LPO) is refers to oxidation of lipids by free radicals and it is one of the main manifestations of oxidative damage in tissues and cells. In the hippocampus, increased lipid peroxidation was observed by a significant increase in MDA (malondialdehyd) levels by 30.64% (expressed as nano moles of TBARS/g of protein) in the DZN group when compared to control group ($P < 0.01$) (Fig1). The DGE 40 group, showed decreased levels of MDA by 31.11% when compared with DZN group ($P < 0.0001$) and the DGE 80 group showed decreased levels of MDA by 29.87% when compared with DZN group ($P < 0.01$) (Fig1).




Fig 1: Effect of DGE 40 and DGE 80 on MDA level of DZN treated rats and effect of DZN (100 mg/Kg) on MDA level of normal rats in the hippocampus. Values are mean±SEM (n=6). ** P<0.01, compared to control group, ### P<0.001, compared to DZN group.

Antioxidant enzymes:

SOD, CAT and GPx play crucial role in the cellular antioxidant defense mechanism. The activities of SOD (U/mg of protein) in hippocampus were increased significantly by 11.56% in DZN treated animals compared to control group (P<0.001). Whereas in the DGE 40 group, a significant decrease by 27.4% were observed in the activities of SOD compared to DZN group (P<0.05). In the DGE 80 group, a significant decrease by 21.61% was observed (p<0.01). In rats' hippocampus non-significant differences were observed between the groups in CAT activity (p>0.05) (Fig3). GPx activity was diminished by 39% by DZN (p<0.05). In rats supplemented with DGE 40, 80 we observed an increase about 64.3%, (p<0.05) and 92.8%, (p<0.05) respectively (Fig4).




Fig2: Effect of DGE 40 and DGE 80 on SOD level of DZN treated rats and effect of DZN (100 mg/Kg) on SOD level of normal rats. Values are mean±SEM (n=6). *** P<0.001, compared to control group, * p<0.05, compared to DZN group.




Fig3: Activities of CAT in control group and treated rats with DZN, DGE40 and DGE 80.




Fig4: Effect of DGE 40 and DGE 80 on GPx activities of DZN treated rats and effect of DZN (100 mg/Kg) on GPx activities of normal rats in the hippocampus. * p<0.05, compared to control group, # p<0.05 and ## p<0.01, compared to DZN group.

Total thiol content:

Rats which received DZN showed non-significant values of total thiol (TSH) when compared with control rats (p>0.05). Rats in DGE 40 group showed non-significant values of total thiol when compared with DZN group (p>0.05). The DGE 80 group showed a significant increase by 85.6% when compared with DZN group (p<0.01) (Fig5).




Fig5: Values of total thiol in control group and treated rats with DZN, DGE40 and DGE 80. ^{##} p<0.01, compared to DZN group.

Discussion:

Recently, an increasing amount of attention has been concentrated towards free radical mediated damage triggered by pesticide exposure in biological systems [29]. Some herbal drugs with antioxidant activity have gained importance as the dietary intake of antioxidants obtained from natural sources is considered to be relatively safe and without undesirable side effects [18]. The aim of present study was to investigate the effect of *D.glabrum* against DZN-induced oxidative stress in hippocampus.

The current study indicated that DZN treatment induced oxidative stress, and increased MDA levels and SOD activities in the hippocampus but decreased GPx activities and no significant changes showed in CAT activities. Salehi et al., showed that intraperitoneal injection of DZN leads to an increase in SOD activity and MDA levels in DZN doses higher than 50 mg/kg and no-significant differences in CAT activity. The increase of MDA levels observed in the hippocampus following DZN exposure may be attributable to the excessive production of ROS. In a study by Salehi et al. it was identified that DZN increased SOD activities at doses of 50 and 100 mg/kg in compared with the control group but there were no significant changes observed in brain CAT activity. Also MDA concentration was significantly increased at 100 mg/kg dose in comparison with the control group. Abbasnejad et al. indicated that intraperitoneal injection of DZN leads to an increase in CAT and SOD activities in doses higher than 30 mg/kg and increase in MDA levels in 100 mg/kg doses in compared to control group. Our results are in agreement with this studies [30, 31, 32]. Under the oxidative stress conditions, SOD acts as the first line of defense against superoxide as it converts the superoxide anion to H₂O₂ which is converted to H₂O by CAT and GPx. Therefore the increased lipid peroxidation may be interpreted here by an inhibition of SOD, CAT, GPx activities and other antioxidants in the brain tissue leading to membrane injury and neuron death [33,34].

174 In our investigation, ethanolic extract of *D. glabrum* showed an amelioratory effect on DZN-induced changes in MDA
175 levels and antioxidant enzymes (SOD, CAT and GPx). Ghollassi Mood, Yousefzadi et al., Shahidi et al. and Kumar et
176 al. showed that members of the genus *Dorema* (Apiaceae), has a lot of biological properties such as antioxidant,
177 antispasmodic, expectorant, carminative, diaphoretic, mild diuretic, emmenagogue, stimulant, vasodilator [35,36]
178 antimicrobial, antifungal [37,38] and hepatoprotector [39] properties. Most of the biological action of *D. glabrum* is
179 probably ascribed to its antioxidant potential. In the present study, DGE as pre-treatment with DZN that was injected
180 intraperitoneally in last day of DGE usage, caused a significant increase in glutathione peroxidase, a significant decrease
181 of TBARS and a significant decrease in superoxide dismutase and no significant changes in catalase activity in the rats'
182 hippocampus when compared to the rats treated with DZN. Our study demonstrated that *dorema* had an amelioratory
183 effect on oxidative stress induced by DZN.

184
185 **References:**

- 186
187
188 1. Zhou J.F., Xu G.B., Fang W.J. Relationship between acute organophosphorus pesticide poisoning and damages
189 induced by free radicals. *Sci. Biomed Environ*, **15**: 177–186, 2002.
- 190
191 2. Casrellano C., Cabis S., Puglisi P. Psychopharmacology of memory modulation: evidence for multiple interactions
192 among neurotransmitters and hormones. *Res. Behave Brain* **77**: 1-21, 1996.
- 193
194 3. Reigart J.R., and Roberts J.R. Recognition and management of pesticide poisoning. Washington, D.C.: U.S. EPA.,
195 Fifth edition, 34-38, 1999.
- 196
197 4. Mense SM., Sengupta A., Lan C., et al. The common insecticides cyfluthrin and chlorpyrifos alter the expression of
198 a subset of genes with diverse functions in primary human astrocytes. *Sci. Toxicol*, **93**(1):125–35, 2006.
- 199
200 5. Singh AK., Jiang Y. Lipopolysaccharide (LPS) induced activation of the immune system in control rats and rats
201 chronically exposed to a low level of the organothiophosphate insecticide, acephate. *Toxicol Ind Health*. **19**(2–6): 93–
202 108, 2003.
- 203
204 6. Wenk GL., McGann K., Hauss-Wegrzyniak B., et al. The toxicity of tumor necrosis factor-alpha upon cholinergic
205 neurons within the nucleus basalis and the role of norepinephrine in the regulation of inflammation: implications for
206 Alzheimer's disease. *Neuroscience*, **121**(3):719–29, 2003.

207
208 7. Jafari M., Salehi M., Ahmadi S., et al. The role of oxidative stress in diazinon-induced tissues toxicity in wistar and
209 Norway rats. *Toxicol Mech Methods*, **22(8)**: 638-47, 2012.
210
211 8. Bebe F.N., Panemangalore M. Exposure to low doses of endosulfan and chlorpyrifos modifies endogenous
212 antioxidants in tissue of rats. *Sci. Environ. Health*, **38**: 349–363, 2003.
213
214 9. Floyd R. Antioxidants, oxidative stress, and degenerative neurological disorders. *Proc Soc. Exp. Biol. Med.* **222**:
215 236–245, 1999.
216
217 11. Poeggeler B., Reiter R.J., Tan D.X., Chen L.D., Manchester L.C. Melatonin, hydroxyl radical mediated oxidative
218 damage, and aging: a hypothesis. *Res. Pineal* **14**: 151-168, 1993.
219
220
221 12. Reiter R., Poeggeler B., Tan D., et al. Antioxidant capacity of melatonin: a novel action not requiring a receptor.
222 *Lett. Neuroendocrinol*, **15**: 103-116,1993.
223
224 13. Cizova H., Lojek H., Kubala L., et al. The effect of intestinal ischemia duration on changes in plasma antioxidant
225 defense status in rats. *Res. Physiol*, **53**: 523-531, 2004.
226
227 14. Halliwell B., Gutteridge J. *Free Radicals in Biology and Medicine*, Oxford University Press, London, pp. 936,
228 1999.
229
230 15. Lee A., East J. Balgaug, Interactions of insecticides with biological membranes. *Sci. Pest.*, **32**: 317–327, 1991.
231
232 16. Dehghan G., Shafiee A., Ghahreman MH. Antioxidant Potential of Various Extracts from *Ferula szovitsiana* in
233 Relation to Their Phenolic Content. *Pharmaceutical Biology*, **45(9)**: 691–699, 2007.
234
235 17. Mandenova I. *Heracleum*. In: Rechinger KH, editor. *Flora Iranica, Umbelliferae* 1987; 162: 492-502.
236
237

238
239 18. Dehghan G., Khoshkam Z. Tin (II)-quercetin complex: Synthesis, spectral characterization and antioxidant activity.
240 Food Chemistry.2012, 131, 422-427.
241
242 19. Keyhani J., Keyhani E., Kamali J. Thermal stability of catalases active in dormant saffron (*crocus sativus L*) corms.
243 Mol. Biol. Rep.,**29**:125-128, 2002.
244
245 20. Gao R., Yuan Z., Zhao Z., et al. Mechanism of pyrogallol autoxidation and determination of superoxide dismutase
246 enzyme activity. Bioelectrochem. Bioenerg. **45**: 41-45,1998.
247 21. Paglia DE., Valentine WN. Studies on the quantitative and qualitative characterization of erythrocyte glutathione
248 peroxidase. J Clin La. Med. **70**: 158–169, 1967.
249
250 22. Hu ML., Dillard CJ. Plasma SH and GSH measurement. Methods Enzymol, **233**: 385–7, 1994.
251
252 23. Saavedra A., Carreton A., Xifro O., et al. Increased PKA signaling disrupt recognition memory and spatial
253 memory: role in Huntington’s disease. Human Molecular Genetics, **20(21)**: 4232-4247, 2011.
254
255 24. Banerjee BD., Seth V., Bhattacharya A., et al. Biochemical effects of some pesticides on lipid peroxidation and
256 free-radical scavengers. Toxicol Lett **107(1–3)**: 33–47, 1999.
257
258 25. Satho K. Serum lipid peroxidation in cerebrovascular disorders determined by a new colorimetric method. Clin
259 Chem Acta.,**90**:37–43, 1978.
260
261 26. Ukeda H., Maeda S., Ishii T., et al. Spectrophotometric assay for superoxide dismutase based on tetrazolium salt 3’-
262 1-(phenylamino)-carbonyl-3, 4-tetrazolium]-bis(4-methoxy-6-nitro)benzenesulfonic acid hydrate reduction by
263 xanthine-xanthine oxidase. Anal Biochem., **251**: 206–9, 1997.
264
265 27. Aebi H. Catalase *in vitro* assay. Meth Enzymol., **105**: 121–6, 1984.
266
267 28. Lowry OH., Roserbrough NJ., Farr AL., et al. Protein measurement with the folin phenol reagent. J Biol Chem.,
268 **193**: 265–75, 1951.

269
270 29. Banerjee BD, Seth V, Bhattacharya A, Pasha ST, Chakraborty AK. Biochemical effects of some pesticides on lipid
271 peroxidation and free-radical scavengers. *Toxicol Lett.* 1999;107(1–3):33–47.
272
273 30. Salehi M., Jafari M., Asgari A., et al. Study of diazinon effect on antioxidant enzymes and lipid peroxidation in rat's
274 brain. *Iranian Journal of medical sciences*, **70**: 25-31, 2010.
275
276 31. Abbasnejad, M. Jafari, M.Asgari, A. Haji Hossaini, R. Hajigholamali, M. Salehi, M. and Salimian, M. Acute
277 Toxicity Effect of Diazinon on Antioxidant System and Lipid Peroxidation in Kidney Tissues of Rats. *Scientific-*
278 *Research Journal of Shahed University*, **17(83)**: 35-42, 2009.
279
280 32. Yilmaz M., Yilmaz M., Altuntas I. Diazinon-induced brain toxicity and protection by vitamins E plus C. *Toxicol Ind*
281 *health*,**28(1)**:51-7,2012.
282
283
284 33. Fridovick I. Superoxide radical: an endogenous toxicant. *Annu Rev Pharmacol Toxicol* **23**:239–253, 1975.
285 34. Nehru B., Anand P. Oxidative damage following chronic aluminium exposure in adult and pup rat brains. *J Trace*
286 *Elem Med Biol.*, **19**: 203–208, 2005.
287
288
289
290 35. Ghollassi Mood S.. A contribution to some ethnobotanical aspects of Birjand Flora (Iran). *Pakistan Journal of*
291 *Botany*, **40(4)**: 1783-1791, 2008.
292
293 36. Yousefzadi M., Heidari M., Akbarpour M., et al. In vitro cytotoxic activity of the essential oil of *Dorema*
294 *ammoniacum* D. Don. *Middle-East Journal of Scientific Research*, **7(4)**, 511-514, 2011.
295
296
297 37. Shahidi GH., Moein MR., Foroumadi AR., et al. Cytotoxic activity of medicinal plants used in Iranian traditional
298 medicine on two strains of *Saccharomyces cerevisiae*. *DARU*, **10(4)**, 162-164,2002.
299

- 300 38. Kumar VP., Chauhan NS., Padh H., et al. Search for antibacterial and antifungal agents from selected Indian
301 medicinal plants. *Journal of Ethnopharmacology*, **107**: 182-188, 2006.
- 302
- 303 39. Govind P. Medicinal plants against liver diseases. *International Research Journal of Pharmacy*, **2(5)**: 115-121,
304 2011.