

Case study

Intrahepatic multicystic biliary hamartoma: presentation of a case report and magnetic resonance imaging / magnetic resonance cholangiopancreatography findings

Abstract : Biliary hamartomas, known as von Meyenburg complexes (VMCs), are benign liver malformations. They are histologically characterized by cystic dilated bile ducts surrounded by numerous fibrous stromal elements measuring up to 5 mm in diameter. Incidental detection of VMCs by autopsy is difficult. Detection of VMCs by imaging is also difficult because of their asymptomatic nature and small size and also rarity. Moreover, they are easily confused with metastatic diseases of the liver, especially in imaging .

A 39-year-old man presented to our hospital with a 6-month history of recurrent nonspecific abdominal pain. Abdominal ultrasonography (US) revealed multiple cystic lesions in the liver. The diagnosis of metastases was suggested. However, final diagnosis of VMCs was confirmed by magnetic resonance imaging and magnetic resonance cholangiopancreatography

.This case report highlights the routine differential diagnosis of biliary microhamartomas by magnetic resonance imaging and magnetic resonance cholangiopancreatography.

Key words : biliary microhamartomas, magnetic resonance imaging (MRI), magnetic resonance cholangiopancreatography(MRCP)

Introduction

Biliary hamartomas, known as von Meyenburg complexes (VMCs), are benign liver malformations. They are histologically characterized by cystic dilated bile ducts surrounded

26 by numerous fibrous stromal elements [1,2] measuring up to 5 mm in diameter. Incidental
27 detection of VMCs by autopsy is difficult. Detection of VMCs by imaging is also difficult
28 because of their asymptomatic nature and small size [3]. VMCs are also rare. Moreover, they
29 are easily confused with metastatic diseases of the liver, especially in imaging [4].

30

31 Therefore, an understanding of the imaging traits of VMCs is needed to establish a list of
32 differential diagnoses, which will decrease the need for methods such as biopsy or laparotomy
33 [5]. We herein report a case of VMCs and describe the routine diagnostic magnetic resonance
34 imaging (MRI) and magnetic resonance cholangiopancreatography (MRCP) findings of
35 biliary microhamartomas.

36

37

38 **Case report**

39

40 A 39-year-old man presented to our hospital with a 6-month history of recurrent nonspecific
41 abdominal pain. Physical examination findings were unremarkable. Laboratory examination
42 results were normal with the exception of a slight elevation of gamma-glutamyl transferase
43 (142 mg/dL; reference range, 0–55 mg/dL). Abdominal ultrasonography (US) revealed
44 multiple cystic lesions in the liver that appeared similar to metastases. Subsequent MRI
45 showed multiple small cysts that were hypointense on T1-weighted images (Fig. 1a, b) and
46 hyperintense on T2-weighted images; they were scattered in the liver parenchyma (Fig. 2a, b).
47 MRCP showed small cysts distributed uniformly within the contour of the liver, creating a
48 “starry sky” configuration (Fig. 3a, b).

49

50 The patient was diagnosed with multiple VMCs based on the typical MRI features.
51 Verification using these imaging techniques within the 6-month follow-up confirmed the
52 diagnosis of VMCs.

53

54

55 **Discussion**

56

57 A VMC is a benign congenital malformation of the biliary duct. It was first defined in 1918
58 by von Meyenburg [6]. Although jaundice and portal hypertension may be caused by a mass
59 effect, patients are usually asymptomatic [7]. VMCs may be single or multiple, with sizes
60 ranging from 1 to 15 mm [1]. Because of the small size of the lesions, an ultimate description
61 is difficult to attain.

62

63 The prevalence of VMCs by autopsy ranges from 0.6% to 2.8% [8]. Histologically, the
64 lesions include disorganized and dilated bile ducts and ductules surrounded by fibrous stroma
65 [9]. US imaging shows hypoechoic, hyperechoic, or mixed heterogenic echoic structures
66 [1,3,4]. The multiple comet-tail sign is considered to be a specific US finding of VMCs [3].
67 Additionally, lesional echogenicity might be related to the number and size of dilated bile
68 ducts and the degree of fibrosis [9]. In contrast, enhanced computed tomography shows that
69 VMCs are usually of low attenuation with irregular margins. Most reported cases have
70 suggested that VMCs do not demonstrate contrast enhancement [3,9]. On MRI, VMCs are
71 defined as hypointense on T1 and hyperintense on T2 compared to the surrounding liver
72 parenchyma [1,9].

73

Although VMCs are benign, some reports have described hepatic malignancies with a background of VMCs, including hepatocellular carcinoma and cholangiocarcinoma [10]. VMCs are rare and usually only seen as multiple small nodules. They are sometimes confused with metastatic liver disease, microabscesses, diffuse primary hepatocellular carcinoma, biliary cysts, or Caroli disease [1,6,8].

Conclusion

The use of various imaging modalities with follow-up has proven helpful for the diagnosis of VMCs. A correct diagnosis is easier to reach when typical imaging findings are present. Otherwise, histological verification may be needed.

Consent Disclaimer:

As per international standard or university standard, patient's consent has been collected and preserved by the author.

References

1. Lev-Toaff AS, Bach AM, Wechsler RJ, Hilpert PL, Gatalica Z, Rubin R. The radiologic and pathologic spectrum of biliary hamartomas. *AJR Am J Roentgenol* 1995; 165: 309-313
2. Wei SC, Huang GT, Chen CH, Sheu JC, Tsang YM, Hsu HC, Chen DS. Bile duct hamartomas. A report of two cases. *J Clin Gastroenterol* 1997; 25: 608-611

3. Luo TY, Itai Y, Eguchi N, Kurosaki Y, Onaya H, Ahmadi Y, Niitsu M, Tsunoda HS. Von Meyenburg complexes of the liver: imaging findings. *J Comput Assist Tomogr* 1998; 22: 372-378
4. Cooke JC, Cooke DA. The appearances of multiple biliary hamartomas of the liver (von Meyenberg complexes) on computed tomography. *Clin Radiol* 1987; 38: 101-102
5. Mortelé B, Mortelé K, Seynaeve P, Vandeveld D, Kunnen M, Ros PR. Hepatic bile duct hamartomas (von Meyenburg Complexes): MR and MR cholangiography findings. *J Comput Assist Tomogr* 2002; 26: 438-443
6. Zheng RQ, Zhang B, Kudo M, Oanda H, Inoue H. Imaging findings of biliary hamartomas. *World J Gastroenterol*, 2005; 13(40):6354-6359.
7. Wohlgemuth WA, Bottger J, Bohndorf JB. MRI, CT, US and ERCP in the evaluation of bile duct hamartomas (von Meyenburg complex): A case report. *Eur Radiol*, 1998; 8:1623-1626.
8. Markhard BK, Rubens DJ, Huang J, Dogra VS. Sonographic Features of Biliary Hamartomas with Histopathologic Correlation. *J Ultrasound Med*, 2006; 25:1631-1633.
9. Yong Moon Shin. Biliary hamartoma presented as a single mass. *The Korean Journal of Hepatology*, 2011; 17:331-334.
10. Maher MM, Dervan P, Keogh B, Murray JG. Bile duct hamartomas (von Meyenburg complexes): Value of MR imaging in diagnosis. *Abdominal Imaging*, 1999; 24:171-173.

Figure 1A: T2-weighted three-dimensional magnetic resonance cholangiopancreatography images (coronal plane). Multiple hyperintense cysts with scattered placement are observed in the liver parenchyma, the largest diameter reaching about 2 cm. No significant association between the cysts and biliary ducts is present.


Figure 1b: T2-weighted three-dimensional magnetic resonance cholangiopancreatography images (coronal plane). Multiple hyperintense cysts with scattered placement are observed in the liver parenchyma, the largest diameter reaching about 2 cm. No significant association between the cysts and biliary ducts is present.

Figure 2a :T1-weighted contrast-enhanced axial fat-suppressed sequences. (a, b) Multiple hypointense cysts, the largest of which is 2 cm in diameter, are observed in the liver parenchyma without contrast enhancement.

Figure 2b :T1-weighted contrast-enhanced axial fat-suppressed sequences. (a, b) Multiple hypointense cysts, the largest of which is 2 cm in diameter, are observed in the liver parenchyma without contrast enhancement.

Figure 3a :Multiple hyperintense cysts in the liver parenchyma. (a) Coronal-plane T2-weighted sequence, (b) axial fat-suppressed T2-weighted sequence

Figure 3b: Multiple hyperintense cysts in the liver parenchyma. (a) Coronal-plane T2-weighted sequence, (b) axial fat-suppressed T2-weighted sequence.




142

143

144

145

Figure1A




146

147

148

Figure 1b


149



150

151 Figure2a

152




153

154

155 Figure 2b

156




157

158

159

Figure 3a

160



161

162

163

164

Figure 3b