

**Determination of Some Biochemical Parameters in Streptozotocin-Induced Diabetic Albino Rats Pre-Treated and Post-Treated with *Vernonia Amygdalina* and *Gongronema Latifolium* Extracts**

**ABSTRACT**

The use of herbs in the management of diabetes mellitus and its complications have been reported. This study was thus aimed at determining the levels of some biochemical parameters in streptozotocin-induced diabetic albino rats pre-treated and post-treated with *Vernonia amygdalina* and *Gongronema latifolium* extracts. Fifty (50) albino rats weighing between 150 – 250g were used for this study. 25 albino rats were used for each phase of the treatment. The pre-treatment phase involved the treatment of the rats with 400mg/kg b.w (singly) and 200mg/kg b.w (combined) extracts for 14 days, after which diabetes mellitus was induced using streptozotocin before the rats were sacrificed. The post-treatment phase involved the inducement of diabetes with streptozotocin after which the rats were treated with 400mg/kg b.w (singly) and 200mg/kg b.w (combined) extracts for 28 days before the animals were sacrificed. Blood was collected via cardiac puncture and plasma/serum was collected. Total cholesterol (TC), high density lipoprotein cholesterol (HDL-C), low density lipoprotein cholesterol (LDL-C), malondialdehyde (MDA), and total antioxidant capacity (TAC) were determined using standard procedures while glucose was measured with a glucometer. Cardiovascular risk ratios such as Castelli risk ratio I and II and atherogenic index of plasma were also calculated. Results showed that extracts of *V. amygdalina* (only), *G. Latifolium* (only) and V.A+ GL (combined) significantly ( $p < 0.05$ ) reduced the TC, LDL-C, FBS, MDA and TAC levels while HDL-C level significantly ( $p < 0.05$ ) increased. The cardiac risk indices (CRI-I and CRI-II, and Atherogenic index of plasma (AIP) in both the pre-treated and post-treated rats were also reduced when compared to diabetic control. It is thus evident that these plants' extracts possess hypoglycaemic, hypolipidaemic, and antioxidant properties and thus could be used to reduce cardiovascular risks. Therefore, the traditional use of these plants in the management of cardiovascular complications is highly recommended.

**Keywords:** Diabetes mellitus, *Vernonia amygdalina*, *Gongronema latifolium*, hypolipidaemic, hypoglycaemia, antioxidant, hyperglycaemia.

**1.0 INTRODUCTION**

A chronic hyperglycaemic condition, stemming from the decreased supply of insulin, its action or both, resulting in a cluster of metabolic disorder is termed diabetes mellitus (DM) [1]. Mainly two types of diabetes mellitus occur, such as type I and II. The former is also called juvenile diabetes or insulin-dependent diabetes mellitus, as it occurs mainly in the young, characterized by destruction of beta cells which may be due to an autoimmune process or accident, usually leading to absolute deficiency of insulin [2]; patients with this type of diabetes will require insulin therapy to maintain normal blood glucose concentration,

39 while the latter is also called maturity-onset diabetes or non-insulin dependent diabetes  
40 mellitus representing ninety percent of all cases of diabetes mellitus [3], and may be due to  
41 insensitivity of target tissues to insulin [1].

42 Oxidative stress is a state whereby there is an imbalance between the generation and  
43 neutralization of reactive oxygen and nitrogen species (RONS) such that the antioxidant  
44 capacity of the cell becomes overwhelmed [4]. When free radicals are overproduced,  
45 biomolecules such as lipids, proteins and DNA may get destroyed, leading to the induction of  
46 several disease conditions such as cancer, diabetes, cardiovascular diseases, aging and other  
47 degenerative diseases. The diabetogenic potential of streptozotocin is dependent on its ability  
48 to generate reactive oxygen species [5], which exert toxic effects on the beta cells of the  
49 pancreas, decreasing its ability to produce insulin; these free radicals may be responsible for  
50 the diabetes-induced pathological conditions. The antioxidant capacity of plasma is said to be  
51 the primary measure and marker in the evaluation of the status and potential of oxidative  
52 stress in the body. To prevent cellular biomolecules from being damaged, certain compounds  
53 occur in the plasma, and function against the oxidative stressors in the body. The total sum  
54 potential of all the antioxidant molecules in the plasma is a reflection of the antioxidant  
55 capacity of the plasma. In all processes where reduced plasma antioxidant potential is  
56 reported, prevalence of oxidative stress is also reported [6]. In diabetics with poor glycaemic  
57 control, the plasma antioxidant level is significantly low, while diabetics with good  
58 glycaemic control have higher plasma antioxidant level. Oxidative stress in diabetics coexists  
59 with a decrease in the antioxidant status, which in turn, may further elevate the deleterious  
60 effects of the free radicals.

61 Coronary artery disease is one of the pathologies associated with diabetes mellitus, with  
62 dyslipidemia been identified as one of the most important risk factors. Low HDL-C, high

63 Triglyceride and high LDL-C levels have been associated with an increased incidence of  
64 coronary artery disease [7]. However, the absence of an abnormal lipid profile does not  
65 completely rule out the possibility of coronary artery disease, thus high risk persons may be  
66 identified through predictions using different combinations of the lipid profile parameters;  
67 these include the Atherogenic Index of Plasma (AIP) and Castelli Risk Index (CRI) and  
68 Atherogenic Coefficient (AC) [8]. The AIP is calculated as  $\text{Log TG/HDL-C}$ , CRI-I as  
69  $\text{TC/HDL-C}$ , CRI-II as  $\text{LDL-C/HDL-C}$ , and AC as  $\text{TC-HDL-C/HDL-C}$  [9].

70 The World Health Organization stated that eighty percent of the emerging world's population  
71 depends on the use of herbal medicine. Herbs being the oldest form of healthcare produce  
72 several plant extracts and phytochemicals with several therapeutic benefits and affordable  
73 treatment [10]. Plants major constituents include terpenoids, flavonoids, glycosides, alkaloids  
74 and carotenoids, which are often believed to possess antidiabetic effect.

75 *Vernonia amygdalina* (bitter leaf) is the most prominent species belonging to the family  
76 Asteraceae [11], and mainly found in the tropical parts of Africa, [12], where it is used as a  
77 vegetable or flavour in soups. It is commonly referred to as bitter leaf due to its bitter taste in  
78 nature; the bitter taste is due to its anti-nutritional components such as alkaloids, saponins,  
79 glycosides and tannins. The leaves have found relevance in traditional folk medicine as  
80 antihelminth, a laxative herb and an antimalarial as they are known as quinine substitute [13].  
81 It is also used in the treatment of cough and hypertension [14,15]

82 *Gongronema latifolium* (utazi) is a nutritional and medicinal plant which is edible, and  
83 commonly found in Nigeria, particularly in the rain forest zones, and the tropics in other  
84 African countries. As a result of their nutritional and ethnomedicinal values, several studies  
85 have been reported regarding methanolic and ethanolic extracts of the herbs. *Gongronema*  
86 *latifolium* (Asclepiadaceae) is also a tropical rainforest plant primarily used as spice and

vegetable in traditional folk medicine [16,17]. Reports by various authors showed that it contains essential oils, saponins and pregnans among others [18, 19]. The leaves of *Gongronema latifolium* have protective role against diabetes, hypertension, stomach upsets and pains, and typhoid fever [20].

Most studies conducted on the herbs focussed mostly on the efficacy of the extracts following induction of organ damage. The literature on the prophylactic efficacy of the aqueous extracts of these herbs before induction of organ damage to evaluate the therapeutic potentials of the herbs in protecting the body organs against xenobiotic assaults are scarce. Thus, this study was designed to determine the serum levels of some biochemical parameters in streptozotocin-induced diabetic albino rats pre-treated and post treated with *Vernonia amygdalina* and *Gongronema latifolium*.

## 2. MATERIALS AND METHODS

This study was carried out at the Animal House, Department of Physiology, University of Port Harcourt, Nigeria. Fifty (50) albino rats weighing between 150 – 250g were used for this study. They were allowed to acclimatize a week prior to experimentation. They were kept in properly ventilated cages, at a room temperature of about 27<sup>0</sup>C and 12 hours light/dark cycle, and the animals were fed with growers' marsh and water obtained from tap *ad libitum*.

Fresh mature leaves of *V. amygdalina* and *G. Latifolium* were purchased from the Mile 1 market, Port Harcourt, Nigeria. Botanical identification was confirmed at the Herbarium, Department of Plant Science and Biotechnology, University of Port Harcourt Rivers State. The leaves were washed and air-dried in the shade for 4 days, and the dried leaves were milled into powder with a blender and stored in an airtight plastic bag and kept from sunlight. 1g of the powdered herb was soaked in 100ml of distilled water for 24 hrs and filtered with a

muslin cloth to produce a concentration of 10mg/ml of the aqueous extract used for the experiment.

In other to induce diabetes mellitus, the adult albino rats were allowed to fast overnight, and were then injected with a single intraperitoneal dose of streptozotocin at 50 mg/kg b.w in 0.1 molar citrate buffer, pH 4.5, while the control animals were injected intraperitoneally with citrate buffer alone (1 ml/kg b.w). All animals were allowed free access to feed and water after streptozotocin (STZ) injection, and they were left undisturbed for a minimum of 72 hours for hyperglycaemia to develop. After that, fasting blood glucose levels of the animals were measured with One Touch Ultra Mini Glucometer. Animals with blood glucose greater than or equal to 13.8 mmol/l were considered hyperglycaemic.

During the experimental period, there was strict adherence to ethical regulations required for handling experimental animal in accordance with National and Institutional Guidelines for Protection of Animal Welfare [21].

There were two phases; phase I (the pre-treatment phase) which was for 14 days, and phase II (the post-treatment phase) which was for 28 days, and the dosage of the plant extracts was 400mg/kg, administered twice daily using the method of Atangwho *et al.* [22]. In phase I (pre-treatment phase), the animals were divided into five (5) groups with 5 animals in each group.

Group A (Normal Control): consisted of rats which were maintained on food (Grower's marsh) regime and water 0.1 M. citrate buffer

Group B (Diabetic Control): consisted of rats injected with streptozotocin at 50 mg/kg b.w in 0.1 M. citrate buffer, and maintained on food (Grower's marsh) regime and water

Group C: consisted of rats pre-treated with 400mg/kg of *Vernonia amygdalin* twice daily, and food (Grower's marsh) regime and water prior to induction with streptozotocin

133 Group D: consisted of rats pre-treated with 400mg/kg of *Gongronema latifolium* twice daily,  
134 and food (Grower's marsh) regime and water prior to induction with streptozotocin.

135 Group E: consisted of rats pre-treated with combined extracts of *Vernonia amygdalina*  
136 (200mg/kg) and *Gongronema latifolium* (200mg/kg) twice daily, and food (Grower's marsh)  
137 regime and water prior to induction with streptozotocin.

138

139 The rats in groups C, D and E after being pre-treated with the various extracts for 14 days,  
140 were then allowed to fast overnight on day 15, before injecting with streptozotocin (50  
141 mg/kg), after which they were left for 72 hours, and were then sacrificed.

142 In phase II (post-treatment phase), the animals were divided into five (5) groups with 5  
143 animals in each group.

144 Group A (Normal Control): consisted of rats which were maintained on food (Grower's  
145 marsh) regime and water 0.1 M. citrate buffer

146 Group B (Diabetic Controls): consisted of rats injected with streptozotocin at 50 mg/kg b.w  
147 in 0.1 M. citrate buffer, and maintained on food (Grower's marsh) regime and water

148 Group C: consisted of diabetic rats treated with 400mg/kg of *Vernonia amygdalina* twice  
149 daily post diabetic induction with food (Grower's marsh) regime and water.

150 Group D: consisted of diabetic rats treated with 400mg/kg of *Gongronema latifolium* twice  
151 daily post diabetic induction with food (Grower's marsh) regime and water.

152 Group E: consisted of diabetic rats treated with combined extracts of 200mg/kg each of  
153 *Vernonia amygdalina* and *Gongronema latifolium* twice daily post diabetic induction with  
154 food (Grower's marsh) regime and water.

The rats in groups C, D and E after successful diabetic induction, were treated with the various extracts for 28 days. After the last dose on day 28<sup>th</sup>, the animals were left fasting overnight and sacrificed on the morning of day 29.

About 6ml of whole blood was obtained through cardiac thoracic puncture using a sterile syringe and needle; 3ml was poured into an EDTA and 3ml into a plain tube (allowed for some minutes to clot). Samples were spun for 10 minutes at 4000 rpm to obtain plasma and serum. Serum samples were used for the analysis of total cholesterol (TC), triglycerides (TG) and high density lipoprotein cholesterol (HDL-C), while plasma samples were used for total antioxidant capacity (TAC) and malondialdehyde (MDA) levels.

## **2.1 DETERMINATION OF SERUM TOTAL CHOLESTEROL**

The enzymatic procedure for total cholesterol determination in serum based upon the Trinder [23] method as modified by the Centers for Disease Control and Prevention was used. The method is popularly known as the enzymatic endpoint method. The cholesterol is determined after enzymatic hydrolysis and oxidation. The indicator quinoneimine is formed from hydrogen peroxide and 4-aminoantipyrine in the presence of phenol and peroxidase.

## **2.2. Determination Of High-Density Cholesterol**

The method of Lopes-Virella *et al.* [24] for the determination of high-density cholesterol in serum was employed. Low density lipoproteins and very low density lipoproteins (LDL and VLDL) and chylomicron fractions are precipitated quantitatively by the addition of phosphotungstic acid in the presence of magnesium ions. After centrifugation, the cholesterol concentration in the HDL (high density lipoprotein) fraction, which remains in the supernatant, is determined by CHOD-PAP method.

178 The low density lipoprotein cholesterol (LDL-C) was calculated using the formula below:

179  $LDL-C \text{ (mg/dl)} = \text{Total cholesterol} + (\text{HDL} - \text{Cholesterol} + \text{Triglyceride}/5) \text{ [25]}.$

180 The Castelli's Risk Index I and II were calculated from the formulas below:

181 Castelli's Risk Index (CRI)-I =  $TC/HDL-C$

182 Castelli's Risk Index (CRI)-II =  $LDL-C/HDL-C$

183 The Atherogenic Index of Plasma (AIP) is calculated as  $\text{Log } TG/HDL-C \text{ [9]}$

### 184 **2.3. Determination of Serum Triglycerides (TG)**

185 The colorimetric method of Tietz [26] was employed. The triglycerides are determined after  
186 enzymatic hydrolysis with lipases. The indicator is a quinoneimine formed from hydrogen-  
187 peroxide, 4-amino phenazone and 4-chlorophenol under the catalytic influence of peroxidase.

### 188 **2.4. Estimation of MDA Levels**

189 Malondialdehyde (MDA), a marker of Lipid peroxidation was determined as thiobarbituric  
190 acid reactive substance according to Okhawa *et al.*, [27] with slight modification by Atawodi  
191 *et al.*, [28] using trichloroacetic acid (TCA) and thiobarbituric acid. The product of the  
192 reaction is a coloured complex which absorbs light at 533nm and can hence be measured.  
193 Exactly 2 ml of 15% trichloroacetic acid was measured into a test tube, 2 ml thiobarbituric  
194 acid was added and 100ul of the serum was added. The mixture was incubated at 80°C for  
195 30minutes in a water and allowed to cool for some time followed by centrifugation at  
196 3000rpm for 10minutes. A clear supernatant was collected and the absorbance of it was  
197 determined at 533nm spectrophotometrically. TBARS concentrations were expressed as  
198 nmol/L of protein.



199      Conc. nmol/L Protein = Absorbance of sample /  $1.56 \times 10^{-5}$  x protein Conc. (mg).

## 200      **2.5.      Estimation of Total Antioxidant Activity**

201      Total antioxidant activity was determined according to the method described by Buico *et al.*,  
202      [29]. Preformed radical monocation of 2,2-azinobis-(3 ethyl benzothiazoline 6 sulfonic acid)  
203      (ABTS), a blue green chromophore was generated by reacting 7M ABTS stock solution with  
204      2.45M potassium persulfate solution in acetate buffer(PH 4.5). The solution was kept in the  
205      dark at room temperature for 12 to 24 hours before use. The ABTS solution was diluted to an  
206      absorbance of 1.00 at 734 nm. 50µl of sample was added to 950µl of diluted ABTS. The  
207      sample was properly mixed and incubated in the dark in a water bath at 37<sup>0</sup>C for 20minutes.  
208      The absorbance was read at 734nm. Trolox was used for the calibration of the method.  
209      Inhibition of absorbance level versus Trolox concentration curve was used to express the  
210      serum/plasma TAC in trolox equivalent

## 211      **2.6.      Statistical analysis**

212      Values obtained were presented as mean ± standard error of mean (SEM). The statistical tool  
213      used was the one way analysis of variance (ANOVA) followed by the Tukey's multiple  
214      comparison tests using the IBM SPSS Version 23 Software. Results were considered  
215      statistically significant at 95% confidence interval (p<0.05).

## 216      **3.0 RESULTS**

217      The table 1 shows that the MDA levels in the albino rats pre-treated and post-treated with the  
218      combined extracts of *V. Amydalina* and *G. latifolium*were significantly (p<0.05) reduced  
219      when compared with the response obtained for the single administration of the extracts level.  
220      The TAC level, however, were reduced significantly (p<0.05) in the rats that were post  
221      treated for 28 days after induction of diabetes mellitus in all the groups. The pre-treatment of

222 the rats with the extracts for 14 days whether alone or in combination showed poor glycaemic  
223 response while significantly reduced level of fasting blood sugar was obtained in the 28 days  
224 post-treated rats after induction of diabetes mellitus.

225

UNDER PEER REVIEW

226 **Table 1. Comparison of mean  $\pm$  SD of MDA, TAC and FBS levels in the pre-treated and post treated albino rats.**

Groups	MDA (nmol/L) (14 days)	MDA(nmol/L) (28 days)	TAC (mmolFe/L) (14 days)	TAC (mmolFe/l) (28 days)	FBS (mmol/l) 14 (days)	FBS (mmol/l) 28 days
Group A (N.C)	92.07 $\pm$ 9.90 <sup>a</sup>	101.35 $\pm$ 7.35 <sup>b</sup>	3.65 $\pm$ 0.19 <sup>a</sup>	3.33 $\pm$ 0.39 <sup>a</sup>	4.06 $\pm$ 0.024 <sup>a</sup>	4.78 $\pm$ 0.32 <sup>a</sup>
Group B (D.C)	214.81 $\pm$ 21.19 <sup>b</sup>	220 $\pm$ 6.85 <sup>a</sup>	9.65 $\pm$ 1.19 <sup>b</sup>	12.94 $\pm$ 0.71 <sup>b</sup>	19.44 $\pm$ 1.87 <sup>b</sup>	21 $\pm$ 2.10 <sup>b</sup>
Group C (V. <i>amygdalina</i> )	191.41 $\pm$ 20.37 <sup>d</sup>	223.46 $\pm$ 13.63 <sup>c</sup>	8.85 $\pm$ 0.97 <sup>c</sup>	5.42 $\pm$ 1.23 <sup>c</sup>	16.8 $\pm$ 0.99 <sup>c</sup>	6.08 $\pm$ 0.84 <sup>c</sup>
Group D (G. <i>latifolium</i> )	217.58 $\pm$ 27.31 <sup>b</sup>	158.64 $\pm$ 35.61 <sup>d</sup>	10.01 $\pm$ 1.24 <sup>b</sup>	5.52 $\pm$ 0.49 <sup>c</sup>	15.5 $\pm$ 1.29 <sup>c</sup>	5.1 $\pm$ 0.84 <sup>e</sup>
Group E (combined extract)	184.27 $\pm$ 23.47 <sup>c</sup>	189.17 $\pm$ 5.67 <sup>e</sup>	10.85 $\pm$ 1.22 <sup>e</sup>	4.15 $\pm$ 0.10 <sup>d</sup>	23.3 $\pm$ 1.55 <sup>d</sup>	6.2 $\pm$ 0.68 <sup>c</sup>
F value	4.405	8.038	11.872	32.501	54.840	37.669
P value	0.002 <sup>*</sup>	<0.001 <sup>*</sup>	<0.001 <sup>*</sup>	<0.001 <sup>*</sup>	<0.001 <sup>*</sup>	<0.001 <sup>*</sup>

227 Notes: Groups with different superscript are significantly different at p<0.05, \*significant at p<0.05, MDA=malondialdehyde, TAC=total antioxidant capacity, FBS= fasting  
228 blood sugar

229

The effect of pre-treatment and post treatment of the albino rats with the extracts on lipid profiles is shown in table 2. The table shows that while no significant ( $p>0.05$ ) reduction in the total cholesterol and triglycerides levels in the groups were observed in the pre-treated rats before induction of diabetes mellitus, the values of total cholesterol and triglycerides were significantly ( $p<0.05$ ) reduced with all the extracts. However, while the combined extracts showed more potential in causing significant ( $p<0.05$ ) increase in the HDL-C level in rats pre-treated for 14 days, *G. latifolium* demonstrated the highest rise in HDL-C levels after 28 days of post treatment. The LDL-C level was also significantly reduced in the rats that were administered the extracts in both phases.

The CRI-I and CRI-II obtained in the albino rats pre-treated with the extracts as shown in table 3 was lowest in the animals that received the combined extracts of *G. latifolium* and *V. Amygdalina* and *G. Latifolium* respectively while the AIP was lowest in the rats that received *V. Amygdalina* and indices were far lower than that obtained from the diabetic control group B.

The comparison of the atherogenic potentials of the extracts is shown in table 3. *G. Latifolium*, *V. Amygdalina* and the combined extracts showed obvious potentials in reducing atherogenic risk based on CRI-I, CRI-II and AIP in the rats in the both phases of the study.

253 **Table 2. Comparison of mean  $\pm$ SD of lipid profile in rats pre-treated and post-treated with the extracts.**

Groups	TC (mmol/l) 14 days	TC (mmol/l) 28 days	TG (mmol/l) 14 days	TG (mmol/l) 28 days	HDL-C (mmol/l) 14 days	HDL (mmol/l) 28 days	LDL-C (mmol/l) 14 days	LDL (mmol/l) 28 days
Group A (N.C)	1.72 $\pm$ 0.2 <sup>a</sup>	1.88 $\pm$ 0.05 <sup>a</sup>	0.62 $\pm$ 0.18 <sup>a</sup>	0.5 $\pm$ 0.06 <sup>a</sup>	0.92 $\pm$ 0.24 <sup>a</sup>	1.16 $\pm$ 0.04 <sup>a</sup>	0.51 $\pm$ 0.22 <sup>a</sup>	0.47 $\pm$ 0.04 <sup>a</sup>
Group B (D.C)	1.78 $\pm$ 0.07 <sup>a</sup>	1.97 $\pm$ 0.07 <sup>a</sup>	0.78 $\pm$ 0.17 <sup>a</sup>	0.9 $\pm$ 0.20 <sup>b</sup>	0.65 $\pm$ 0.32 <sup>b</sup>	0.69 $\pm$ 0.05 <sup>b</sup>	0.77 $\pm$ 0.37 <sup>b</sup>	0.86 $\pm$ 0.05 <sup>b</sup>
Group C ( <i>V. amygdalina</i> )	1.87 $\pm$ 0.18 <sup>a</sup>	1.47 $\pm$ 0.19 <sup>b</sup>	0.84 $\pm$ 0.42 <sup>a</sup>	0.5 $\pm$ 0.07 <sup>a</sup>	0.87 $\pm$ 0.19 <sup>c</sup>	0.75 $\pm$ 0.10 <sup>b</sup>	0.69 $\pm$ 0.22 <sup>c</sup>	0.49 $\pm$ 0.10 <sup>a</sup>
Group D ( <i>G. latifolium</i> )	1.76 $\pm$ 0.18 <sup>a</sup>		0.70 $\pm$ 0.3 <sup>a</sup>		0.96 $\pm$ 0.09 <sup>a</sup>		0.44 $\pm$ 0.07 <sup>d</sup>	
		1.57 $\pm$ 0.19 <sup>b</sup>		0.64 $\pm$ 0.16 <sup>a</sup>		0.83 $\pm$ 0.60 <sup>c</sup>		0.45 $\pm$ 0.11 <sup>a</sup>
Group E (combined extract)	1.98 $\pm$ 0.34 <sup>a</sup>		0.63 $\pm$ 0.17 <sup>a</sup>		1.11 $\pm$ 0.14 <sup>d</sup>		0.59 $\pm$ 0.19 <sup>e</sup>	
		1.48 $\pm$ 0.13 <sup>b</sup>		0.37 $\pm$ 0.38 <sup>c</sup>		0.75 $\pm$ 0.29 <sup>b</sup>		0.5 $\pm$ 0.14 <sup>a</sup>
F value	2.285	2.881	1.396	5.881	5.763	7.555	2.475	3.506
P value	0.052*	0.005	0.241*	0.003*	<0.001*	0.001*	0.038*	0.025*

254 Note: Groups with the different superscript are significantly different from each at p<0.05, \*significant at <0.05, TC=total cholesterol, HDL-C=high density lipoprotein  
255 cholesterol, LDL-C=low density lipoprotein cholesterol, TG= triglycerides.

256

257

258 **Table 3. Comparison of mean  $\pm$ SD of atherogenic indices in rats pre-treated and post-treated with the extracts.**

Groups	CRI-I (14 days)	CRI-I (28 days)	CRI-II (14 days)	CRI-II (28 days)	AIP (14 days)	AIP (28 days)
Group A (N.C)	1.95 $\pm$ 0.19	1.64 $\pm$ 0.08	0.62 $\pm$ 0.17	0.41 $\pm$ 0.05	-0.17 $\pm$ 0.08	-0.37 $\pm$ 0.05

Group B (D.C)	3.60±1.08	2.88 ± 0.17	1.91±0.91	1.27 ± 0.12	0.12 ± 0.11	0.12 ±0.03
Group C ( <i>V. amygdalina</i> )	2.21 ± 0.16	1.99 ± 0.13	0.82 ± 0.11	0.68 ± 0.12	-0.05 ± 0.06	-0.17 ±0.04
Group D ( <i>G. latifolium</i> )	1.84±0.11	1.88 ± 0.13	0.47± 0.05	0.55 ± 0.13	-0.17 ± 0.09	- 0.15 ± 0.06
Group E (combined extract)	1.79 ± 0.1	1.97 ± 0.13	0.53 ±0.08	0.66 ± 0.16	-0.25 ± 0.03	-0.31 ± 0.042

---

Note: : CRI- I= Castelli Risk Index I, CRI-II= Castelli Risk Index II, AIP= Atherogenic Index of Plasma

## 4.0 DISCUSSION

The results from this study showed that in both phases of the study, injection of the rats with streptozotocin (diabetic control group) induced a significant increase in the blood sugar level when compared with the normal control group. This observation is in agreement with that of Akpaso *et al.* [30]. The diabetogenic potential of streptozotocin is dependent on its ability to generate reactive oxygen species [5], which exert toxic effects on the beta cells of the pancreas, decreasing its ability to produce insulin. The hypoglycaemic potential of the *V. amygdalina* and *G. latifolium* was observed to be more pronounced in rats that received the extracts after induction of diabetes mellitus by streptozotocin. This observation is in agreement with the reports of Uchenna *et al.* [31] and Owu *et al.* [32] who reported that the leaves of the plants possess anti-diabetic potential in that it reduces blood glucose levels in streptozotocin-induced diabetic rats.

The mean plasma malondialdehyde level for the diabetic control was observed to be significantly ( $p < 0.05$ ) higher than that of the normal control in both phases of the study. This may be a resultant effect of lipid peroxidation, which may be attributed to hyperglycaemia-induced oxidative stress. This report agrees with that of Akpan and Usuh [33] who stated that rats treated with streptozotocin induced a significant increase in the levels of malondialdehyde. The report by Szkudelski [5] that streptozotocin induces increased generation of reactive oxygen species also agrees with the observation in this study. Pre-treatment of the rats with combination of *V. amygdalina* and *G. amygdalina* extracts showed lower significant ( $p < 0.05$ ) reduction in the malondialdehyde level. However, single administration of *G. amygdalina* for 28 days after streptozotocin-induced diabetes mellitus also showed very remarkable decrease in malondialdehyde level. Nwanjo *et al.*, [34] had

earlier reported that *V. amygdalina* extract has the potential to reduce malondialdehyde levels in rats.

Total antioxidant capacity (TAC) is the measure of the amount of free radicals scavenged by a test solution [35], and it is used to evaluate the antioxidant capacity of biological samples [36]. Total antioxidant capacity (TAC) includes both enzymatic antioxidants, such as catalase, and non-enzymatic antioxidants, such as ascorbic acid[37].Studies on total antioxidant capacity in albino rats administered with *V. amygdalina* and *G. latifolium* extracts following diabetic induction is scarce. The total antioxidant capacity rats pretreated with the extracts of *V. amygdalina* and *G. latifolium* were significantly higher when compared to that seen the animals post-treated with the herbs after diabetic induction. This observation could suggest that pre-treatment with the plants could have the potential to protect the body from cellular damage than when it is used therapeutically after the damage has been done. The increase in total antioxidant capacity in the body has been reported to be due to defensive mechanism by the body's antioxidant system in response to the increased oxidative stress [38].

The result of the study further showed that the total cholesterol concentration obtained in the rats pretreated with either single extract of *V. amygdalina* and *G. latifolium* or combination of *V. amygdalina* and *G. latifolium* were not significantly different from the concentration in the diabetic control group. However, in the rats post-treated for 28 days after streptozotocin diabetic induction, the total cholesterol level was significantly ( $p<0.05$ ) reduced when compared to the diabetic control group. The reduction in total cholesterol concentration was not significantly ( $p<0.05$ ) different between *V. amygdalina* and *G. latifolium* aqueous extracts at the dose of 40mg/kg b.w. of the rats. Ugwu *et al.*, [39] had reported that diets preparations made with *V. amygdalina* and *G. latifolium* decreased the serum total cholesterol levels,



however, their observation that *Vernonia amygdalina* diet induced a significantly lower ( $p<0.05$ ) serum total cholesterol when compared to the *Gongronema latifolium* diet preparation was not made in this study. The triglycerides levels in the rats that received the combined extracts of *V. amygdalina* and *G. latifolium* in the pretreated rats were insignificantly ( $p>0.05$ ) reduced when compared to the diabetic control rats. The study also showed that the triglycerides level in rats given the combined extracts for 28 days after induction of diabetes was significantly ( $p<0.05$ ) reduced when compared to the levels in the groups that received the extracts singly. Agwu *et al.*, [39] also reported the lowering of the serum triacylglycerols by the two diet preparations *V. amygdalina* and *G. latifolium* was not significant to each other at equal concentration when compared. The result is in line with the results obtained by Nwanjo [40] and Ugochukwu *et al.*, [17]. The results suggest that the plants could reduce hepatic triacylglycerols biosynthesis and favor the redistribution of cholesterol among the lipoprotein molecules. Adaramoye *et al.*, [41] observed no significant difference in plasma triacylglycerol levels of rats fed on *Telfairia occidentalis* supplemented diets when compared to cholesterol-fed rats. The combined extract also resulted in a significantly reduced triglycerides level in the 28 days post induction rats when compared to the levels from the plants singly.

The result of this study also that treatment aqueous extract of *Gongronema latifolium* and *Vernonia amygdalina* resulted in a significant ( $p<0.05$ ) increase in the level of HDL-C when compared with the diabetic control group. The increase in HDL-C was more pronounced in the rats that were pre-treated with extracts for 14 days before diabetic induction. Furthermore, the increase was more evident in rats that were pre-treated with the combined extracts for 14 days. Since HDL-C is often regarded as the good cholesterol, this study thus shows that these plants have potential protective role against cardiovascular disease (CVD). The comparison of the effects of these plants in increasing the levels of HDL-C in serum in this study showed

that *G. latifolium* induces a significantly higher HDL-C concentration than *V. amygdalina*. This finding is in sharp contrast with the finding of Agwu *et al.*, [39] the fact that *Vernonia amygdalina* induced a significantly higher HDL-C concentration compared to *Gongronema latifolium*. There was a non-significant decrease in the mean HDL level between the diabetic and the normal control groups.

In this study, the effects of pretreatment for 14 days and post treatment for 28 days of aqueous extracts of *Gongronema latifolium* and *Vernonia amygdalina* singly and in combination on the serum LDL-C were also compared. The results show that both preparations significantly lowered the serum LDL-C values though the *Gongronema latifolium* preparation produced a significantly lower serum LDL-C concentration relative to the *Vernonia amygdalina* extracts in both phases of treatment. LDL-C is associated with CVD because they transport cholesterol to the arteries which could lead to the formation of plaque. Therefore, plasma LDL-C level may be used for monitoring the treatment of patients with elevated cholesterol levels. From the results obtained, the plants elicited beneficial effects by lowering the serum LDL in rats.

Evidence from the present study confirms the effects of aqueous extracts of *Gongronema latifolium* and *Vernonia amygdalina* preparations on lipid levels in experimental animals. *Gongronema latifolium* and *Vernonia amygdalina* in single preparations and in combinations were found to be very effective in reducing the levels of serum cholesterol, triacylglycerols and LDL-C thereby exhibiting hypocholestromic effects. They also increased the levels of serum HDL-C in the experimental animals.

Estimation of cardiovascular risk has become the cornerstone of cardiovascular prevention. Although atherogenesis is a multifactorial process, abnormalities in lipoprotein metabolism are one of the key factors, representing around 50% of the population-attributable risk of

developing cardiovascular disease [42]. The total/high-density lipoprotein (HDL) cholesterol ratio, known as the atherogenic or Castelli risk index I and or the LDL/HDL-cholesterol ratio also known as Castelli risk index II are two important components and indicators of vascular risk, the predictive value of which is greater than the isolated parameters. In this respect, an increase in total cholesterol concentration, and specifically LDL-cholesterol, is an atherogenic lipid marker, whereas reduced HDL cholesterol concentration is correlated with numerous risk factors, including the components of the metabolic syndrome, and probably involves independent risk [43]. The Castelli Risk indices (CRI-I) obtained from the animals following pre-treatment the extracts before induction of diabetes mellitus showed that *G. latifolium* reduced the atherogenic effect of diabetes better than *V. amygdalina* while the combined extract was better effect than both of them. Similar CRI-I findings were observed from 28 days post treatment with the extracts after induction of diabetes except that the CRI-I for the animals treated with the combined extracts was higher than that obtained from the 14 days pre-treatment. The CRI-II obtained from the study followed the same trend as the CRI-I for both the 14 days pre-treatment and the 28 days post treatment. Thus, *V. amygdalina* and *G. latifolium* exhibited the capacity to reduced cardiovascular risk.

Many clinical studies make effort to introduce a better marker of atherogenic dyslipidemia that can predict the risk of CVD to be useful for evaluating response to treatment instead of the classical ratio [44]. It has been shown that Atherogenic Index of Plasma (AIP) is a strong marker to predict the risk of atherosclerosis and coronary heart disease [45,46]. AIP is calculated according to the formula,  $\log (TG/HDL-C)$  [46]. The extracts of *V. amygdalina* and *G. Latifolium* either in combination or singly reasonably caused a reduction of the AIP in the experimental animals in both phases of the study implying that these plants can reasonably be employed in the treatment of cardiovascular and coronary heart disease. This

finding is the first to be reported about the medicinal value of *V. amygdalina* and *G. Latifolium* respectively.

## 5.0 CONCLUSION

The hypoglycaemic, hypolipidaemic and antioxidant properties of *V. amygdalina* and *G. latifolium* has been demonstrated this study implying that aqueous extracts of the plants can be useful as prophylactic preventive therapy in conditions of dyslipidaemia, cardiovascular disease and coronary heart disease. The novel discovery that the plants have immense potential to reduce cardiovascular risk ratios such as total/high-density lipoprotein (HDL) cholesterol ratio, known as the atherogenic or Castelli risk index I (CRI-I), LDL/HDL-cholesterol ratio also known as Castelli risk indexII (CRI-II) and Atherogenic index of Plasma (AIP), calculated according to the formula,  $\log (TG/HDL-C)$  has added immensely to the numerous ethnopharmacological usefulness of *V. amygdalina* and *G. latifolium* in the management of cardiovascular disease and associated conditions.

**CONFLICT OF INTEREST:** None

## REFERENCES

1. Ozuogwu JC, Obimba KC, Belonwu CD, Unakalamba CB. The pathogenesis and pathophysiology of type 1 and type 2 diabetes mellitus. *J Physio Pathophysio*. 2013; 4(4): 46-57.
2. Kumar PJ, Clark M. Textbook of Clinical Medicine. Pub: Saunders, London, UK. 2002; 1099-1121.

3. Gonzalez E, Johansson S, Wallander M, Rodriguez L. Trends in the prevalence and incidence of diabetes in the UK: 1996-2005. *J Epidemiol Comm Health.* 2009; 63: 332-336.
4. Sellamuthu PS, Arulselvan P, Kamalraj S, Fakurazi S, M. Kandasamy M 2013. Protective nature of mangiferin on oxidative stress and antioxidant status in tissues of streptozotocin-induced diabetic rats. *ISRN Pharmacol.*2013; 10.1155/2013/750109
5. Szkudelski T. The mechanism of alloxan and streptozotocin action in B cells of the rat pancreas. *Physio Res Acad Scient Bohemoslovaca.*2001;50: 537–46
6. Schiffrini EL. Antioxidants in hypertension and cardiovascular disease. *Mol Interv.* 2010; 10(6):354-362.
7. Fein R, Scheuer J. Heart Disease i Diabetes Mellitus: Theory and Practice. In: Rifkin. H. & Porte, D. Eds. Elsevier, New York.1990; 812-823.
8. Pinto X, Munoz A, Prat J, Masana L. Lipoprotein ratios: physiological significance and clinical usefulness in cardiovascular prevention. *Vasc Health Risk Manag.* 5:757-765.
9. Bhardwaj S, Bhattacharjee J, Bhatnagar MK, Tyagi S. Atherogenic index of plasma, castelli risk index and atherogenic coefficient- new parameters in assessing cardiovascular risk. *ResArt Biol Sci.* 2013; 3 (3): 359-364
10. Adikw E, Bokolo B. Ethanolic leaf extract of *Ocimum gratissimum* ameliorates methotrexate-induced nephrotoxicity in albino rats. *Asian J Med Sci.*2018; 9 (5): 37-43.
11. Johri RK, Singh C. Medicinal uses of *Vernonia* species. *J Med Aromatic Plant Sci.* 1997;19: 744–752
12. Erasto P, Grierson DS, Afolayan AJ. Antioxidant constituents in *Vernonia amygdalina* leaves. *Pharmaceut Bio.*2007;45(3): 195-199
13. Farombi EO. African indigenous plants with chemotherapeutic potentials and biotechnological approach to the production of bioactive prophylactic agents. *Afr. J. Biotech.*2003; 2: 602- 671
14. Kambizi L, Afolayan AJ. An ethnobotanical study of plants used for the treatment of sexually transmitted diseases (njovhera) in Guruve District, Zimbabwe. *J. Ethnopharmacol.* 2001; 77(1):5-9.
15. Amira CA, Okubadejo NU. Frequency of complementary and alternative medicine utilization in hypertensive patients attending an urban tertiary care in Nigeria. *BMC Compl. Altern. Med.* 2007;7: 30-48.
16. Ugochukwu NH, Babady NE. Antioxidant effects of *Gongronema latifolium* in hepatocytes insulin dependent diabetes mellitus. *Filoterapia.* 2002; 73 (7-8): 612-618.
17. Ugochukwu NH, Babady NE, Cobourne M, Gasset SR. (2003). The effect of *Gongronema latifolium* extracts on serum lipid profile and oxidative stress in hepatocytes of diabetic rats. *J Biol Sci.*2003; 28 (1): 1-5.
18. Morebise O, Fafuso MA, Makinde JM, Olajide OA, Awe EO. Antiinflammatory property of *Gongronema latifolium*, *Phytother Res.* 2002; 16:s75-s77
19. Schneider C, Rotscheidt K, Breitmaier E. Four new pregnane glycosides from *Gongronema latifolium* (Asclepiadaceae). *Annalen Der Chemie.* 1993;10:1057-1062.
20. Etim OE, Akpan EJ, Usuh IF. Hepatotoxicity of carbon tetrachloride: protective effect of *Gongronema latifolium*. *Pak. J. Pharm. Sci.* 2008;21:268-274.
21. Public Health Service. Public Health Service Policy on Humane Care and Use of Laboratory Animals. Publication of the Department of Health and Human Services. National Institute of Health . Office of Laboratory Animal Welfare. 2015
22. Atangwho IJ, Ebong PE, Eyong EU, Williams IO, Eteng MU. Comparative Chemical composition of leaves of some antidiabetic medicinal plants: *Azadirachta indica*, *Vernonia amygdalina* and *Gongronema latifolium*. *Afr J Biotec.*2009; 8: 4685-4689.

23. Trinder P. The principle of assay for enzymatic determination of cholesterol. *Ann Clin Biochem.* 1969; 6:24.
24. Lopez-Virella MF, Stone P, Eltis S, Colwell JA. (1977). Cholesterol determination in HDL separated by three different methods. *ClinChem.* 1977; 23: 822-884.
25. Friedewald WT, Levy RI, Fredrickson DS. (1972) Estimation of the concentration of low density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin. Chem.* 1972; 18:499-502
26. Tietz NW. *Clinical Guide to Laboratory Tests.* 3<sup>rd</sup> edition, Philadelphia, W.B. Saunders Company, PhiladelphiaUSA. 1995; 1-178.
27. Okhawa H, Ohisi N, Yagi K. Assay of lipid peroxide in animal tissues by thiobarbituric acid reaction. *Ann Biochem.* 1979; 95: 351-358.
28. Atawodi SE, Adekunle OO, Bala I. Antioxidant, organ protective and ameliorative properties of methanol extract of *Anogeissus leiocarpus* stem bark against carbon tetrachloride-induced liver injury. *Inter J Pharmaceut Sci Res.* 2011; 2(6): 1443-1448.
29. Buico A, Cassino C, Ravera M, Betta P, Osella D. Oxidative stress and total antioxidant capacity in human plasma. *Redox Report.* 2009; 14(3): 125-131.
30. Akpaso MI, Atangwho IJ, Akpantah A, Fisher VA, Igiri A, Ebong E. Effect of Combined Leaf Extracts of *Vernonia amygdalina* (Bitter leaf) and *Gongronema latifolium* (utazi) on the Pancreatic B-cells of Streptozotocin. *Br J Med Med Res.* 2011; 1 (1): 24-34.
31. Uchenna VO, Chinwe EO, John MO, Ijeoma OE. Hypoglycaemic indices of *Vernonia amygdalina* on postprandial blood glucose concentration of healthy humans. *Afri J Biotech.* 2008; 7: 4581-4585.
32. Owu DD, Nwokocha CR, Obembe AO, Essien AD, Ikpi DE, Osim EE. Effect of *Gongronema latifolium* ethanol leaf extract on gastric acid secretion and cytoprotection in streptozotocin-induced diabetic rats. *West Ind Med J.* 2012; 61(9): 852-860
33. Usuh IF, Akpan HD (2015). Antioxidative Efficacy of Combined Leaves Extracts of *Gongronema Latifolium* and *Ocimum Gratissimum* on Streptozotocin-Induced diabetic rat models. *Int. Inv. J. Med. Med. Sci.* 2015; 2(6): 88-95
34. Nwanjo HU, Okafor MC, Oze GO. Anti-lipid peroxidative activity of *Gongronema latifolium* in Streptozotocin-induced diabetic rats. *Nig J Physio Sci.* 2006; 21(1-2): 6165.
35. Ghiselli A, Serafini M, Natella F, Scaccini C. Total antioxidant capacity as a tool to assess redox status: critical view and experimental data. *Free Rad Bio Med.* 2000; 29: 1106–1114
36. Marques SS, Magalhães LM, Tóth IV, Segundo MA. Insights on antioxidant assays for biological samples based on the reduction of copper complexes: the importance of analytical conditions. *Inter J Mol Sci.* 2014; 15:11387–1402.
37. Gupta S, Choi A, YU HY, Czerniak, SM, Holick, EA, Paolella, LJ, Agarwal AA, Combelles CMH. Fluctuations in total antioxidant capacity, catalase activity, and hydrogen peroxide levels of follicular fluid during bovine folliculogenesis. *Reprod Fert Dev.* 2011; 23(5): 673-680. doi:10.1071/RD10270
38. Mortazavi SB, Khajehnasiri F, Allameh A, Akhondzadeh S, Hashemi H. (2013). Total Antioxidant Capacity and Malondialdehyde in Depressive Rotational Shift Workers. *J Env Pub Health.* 2013; 150693
39. Ugwu CE, Alumana EO, Ezeanyika LUS. Comparative effect of the leaves of *Gongronema latifolium* and *Vernonia amygdalina* incorporated diet on the lipid profile of rats. *Biokemistri.* 2009; 21: 59-65.

40. Nwanjo, H. U. Efficacy of aqueous leaf extract of *Vernonia amygdalina* on plasma lipoprotein and oxidative status in diabetic rat models. Nig J Physio Sci. 2005; 20: 39-42.
41. Adaramoye OA, Achem J, Akintayo OO, Fafunso MA. (2007) Hypolipidemic effect of *Telfairia occidentalis* (Fluted pumpkin) in rats fed a cholesterol rich diet. J Med Food. 2007;10: 330-336.
42. Yusuf, Hawken S, Öunpuu S (2004). Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. Lancet. 2004; 364: 937–952.
43. Ascaso J, González SP, Hernández MA. Management of dyslipidemia in the metabolic syndrome. Recommendations of the Spanish HDL. Form Amer J Cardiovas Drugs. 2007;7: 39–58.
44. Mudhaffar SK. Atherogenic Index of Plasma (AIP) As a Parameter in Predicting Cardiovascular Risk in Males Compared To the Conventional Dyslipidemic Indices (Cholesterol Ratios). Karbala J Med. 2013; 6(1):1506-1513
45. Nwagha UI, Ikepeazu EJ, Ejezie FE, Neboh EE, Maduka IC. Atheogenic index of plasma as useful predictor of cardiovascular risk among postmenopausal women in Enugu, Nigeria. Afri Health Sci. 2010; 10(3): 248-252
46. Dobiasova M, Frohlich J. The plasma parameter log (TG/HDL-C) as an atherogenic index: correlation with lipoprotein particle size and esterification rate in apo B-lipoprotein-depleted plasma. (FERHDL). Clin Biochem. 2001; 34:583- 88.