

# Integrability of very weak Solutions for Boundary value problems of Nonhomogeneous **p**-Harmonic equations

Yeqing Zhu, Yanxia Zhou, Yuxia Tong

College of Science

North China University of Science and Technology

Hebei Tangshan, China

**Abstract**—The paper deals with very weak solutions  $u$  to boundary value problems of the nonhomogeneous **p**-harmonic equation. We show that, any very weak solution  $u$  to the boundary value problem is integrable provided that  $r$  is sufficiently close to  $p$ .

**Keywords**—Integrability; Very weak solution; Boundary value problem; **p**-harmonic equation.

## I. INTRODUCTION

Let  $1 < p < n$ ,  $\theta(x) \in W^{1,q}(\Omega)$ ,  $q > r$ ,  $f(x) \in L^{\frac{nq}{n-(p-1)r}}(\Omega, \square^n)$ . We shall examine the boundary value problem of the **p**-harmonic equation

$$\begin{cases} -\operatorname{div}(|\nabla u(x)|^{p-2} \nabla u(x)) = f(x), & x \in \Omega, \\ u(x) = \theta(x), & x \in \partial\Omega, \end{cases} \quad (1.1)$$

Throughout this paper  $\Omega$  will stand for a bounded regular domain in  $\square^n$  ( $n \geq 2$ ). By a regular domain we understand any domain of finite measure for which the estimates (3.3) and (3.4) below for the Hodge decomposition are satisfied, see [1], [2]. A Lipschitz domain, for example, is regular.

**Definition 1.1.** A function  $u \in \theta + W_0^{1,r}(\Omega)$ ,  $\max\{1, p-1\} < r < p$ , is called a very weak solution to the boundary value problem (1.1), for all  $\Phi \in W_0^{1,r/(r-p+1)}(\Omega)$  with compact support sets in  $\Omega$ , there is

$$\int_{\Omega} \left( |\nabla u|^{p-2} \nabla u, \nabla \Phi \right) dx = \int_{\Omega} f(x) \Phi dx \quad (1.2)$$

where  $f(x) \in L^{\frac{nq}{n-(p-1)r}}(\Omega, \square^n)$ .

Recall that a function  $u \in \theta + W_0^{1,p}(\Omega)$  is called the weak solution of the boundary value problem (1.1) if (1.2) holds true for all  $\Phi \in W_0^{1,p}(\Omega)$ . The words very weak in Definition 1.1 mean that the Sobolev integrable exponent  $r$  of  $u$  can be smaller than the natural one  $p$ . see [1], Theorem 1, page 602.

In this paper we will need the definition of weak  $L^t$ -space (see [2]): for  $t > 0$ , the weak

$L^t$  - space,  $L^t_{weak}(\Omega)$ , consists of all measurable functions  $f$  such that

$$\left| \{x \in \Omega : |f(x)| > s\} \right| \leq \frac{k}{s^t}$$

for some positive constant  $k = k(f)$  and every  $s > 0$ , where  $|E|$  is the  $n$ -dimensional Lebesgue measure of  $E$ .

Integrability property is important in the regularity theories of nonlinear elliptic PDEs and systems. In [3], Zhu et al. studied the global integrability of nonhomogeneous quasilinear elliptic equations  $-\text{div}A(x, u, \nabla u) = f(x) + \text{div}(|\nabla u|^{p-2} \nabla u)$ . In [4], Guo et al. studied the higher order integrability of the divergence elliptic equation  $-\text{div}A(x, \nabla u) = -\text{div}f$ . In [5], Zhang et al. studied the global integrability of  $A$ -harmonic equation  $-\text{div}A(x, \nabla u) = -\text{div}f$ . In this paper, we consider the global integrability of the very weak solutions of the boundary value problem (1.1). The main result is the following theorem.

**Theorem 1.1.** Let  $\theta \in W^{1,q}(\Omega)$ ,  $q > r$ , There exists  $\varepsilon_0 = \varepsilon_0(n, p) > 0$ , such that for each very weak solution  $u \in \theta + W^{1,r}_0(\Omega)$ ,  $\max\{1, p-1\} < r < p < n$ , to the boundary value problem (1.1), we have

$$u \in \begin{cases} \theta + L^{q^*}_{weak}(\Omega) & \text{for } q < r, \\ \theta + L^{\tau}_{weak}(\Omega) & \text{for } q = r \text{ and } \tau < \infty, \\ \theta + L^\infty(\Omega) & \text{for } q > n, \end{cases} \quad (1.3)$$

provided that  $|p-r| < \varepsilon_0$ .

Note that we have restricted ourselves to the case  $r < n$  since otherwise any function in  $W^{1,r}(\Omega)$  is in the space  $L^t(\Omega)$  for any  $t < \infty$  by the Sobolev embedding theorem. At the same time, it is also noted that the very weak solution  $u$  to the boundary value problem (1.1) is taken from the Sobolev space  $W^{1,r}(\Omega)$ , and the embedding theorem ensures that the integrability of  $u$  reaches from  $r$  to  $r^*$ . And our result theorem 1.1 improves this integrability. Note that the key to proving the theorem 1.1 is to use Hodge decomposition<sup>[1][6]</sup> to construct the appropriate test function.

## II. PRELIMINARY LEMMAS

**Lemma 1.1.** For  $p \geq 2$  and any  $X, Y \in \square^n$ , one has

$$2^{2-p} |X - Y|^p \leq \langle |X|^{p-2} X - |Y|^{p-2} Y, X - Y \rangle.$$

**Lemma 1.2.** For any  $X, Y \in \square^n$  and  $\varepsilon > 0$ , one has

$$\begin{aligned} & \left| |X|^\varepsilon X - |Y|^\varepsilon Y \right| \\ & \leq \begin{cases} (1+\varepsilon)(|Y| + |X-Y|)^\varepsilon |X-Y|, & \varepsilon > 0, \\ \frac{1-\varepsilon}{2^\varepsilon(1+\varepsilon)} |X-Y|^{1+\varepsilon}, & -1 < \varepsilon \leq 0. \end{cases} \end{aligned}$$

**Lemma 1.3.** For  $1 < p < 2$  and any  $X, Y \in \square^n$ , one has

$$\begin{aligned} & \langle |X|^{p-2} X - |Y|^{p-2} Y, X - Y \rangle \\ & \geq |X - Y| ( (|X - Y| + |Y|)^{p-1} - |Y|^{p-1} ). \end{aligned}$$

**Lemma 1.4.** Let  $\varepsilon_0 > 0$ ,  $\phi: (s_0, \infty) \rightarrow [0, \infty)$  is a decrement function such that for each  $r, s$  ( $r > s > s_0$ ), if

$$\phi(r) \leq \frac{c}{(r-s)^\alpha} (\phi(s))^\beta$$

where  $c, \alpha, \beta$  are constants, we have

(1) if  $\beta > 1$  we have that  $\phi(s_0 + d) = 0$ , where  $d^\alpha = c 2^{\alpha\beta/(1-\beta)} (\phi(s_0))^{\beta-1}$ ;

(2) if  $\beta < 1$  we have that  $\phi(s) \leq 2^{\mu/(1-\beta)} (c^{1/(1-\beta)} + (2s_0)^\mu \phi(s_0)) s^{-\mu}$ , where  $\mu = \alpha/(1-\beta)$ .

### III. PROOF OF THEOREM 1.1

For any  $L > 0$ , let

$$v = \begin{cases} u - \theta + L & \text{for } u - \theta < -L, \\ 0 & \text{for } -L \leq u - \theta \leq L, \\ u - \theta - L & \text{for } u - \theta > L, \end{cases} \quad (3.1)$$

Then according to the hypothesis, we have  $v \in W_0^{1,r}(\Omega)$  and  $\nabla v = (\nabla u - \nabla \theta) \cdot 1_{\{|u-\theta|>L\}}$ , Where  $1_E$

is the characteristic function of the set  $E$ . We introduce the Hodge decomposition of vector field  $|\nabla v|^{p-2} \nabla v \in L^{r/(r-p+1)}(\Omega)$ . So that

$$|\nabla v|^{r-p} \nabla v = \nabla \Phi + h, \quad (3.2)$$

Here  $\Phi \in W_0^{1,r/(r-p+1)}$ ,  $h \in L^{r/(r-p+1)}(\Omega, \mathbb{R}^n)$  is a vector field with zero divergence, and satisfied

$$\|\nabla \Phi\|_{r/(r-p+1)} \leq C(n, p) \|\nabla v\|_r^{r-p+1} \quad (3.3)$$

and

$$\|h\|_{r/(r-p+1)} \leq C(n, p) |p-r| \|\nabla v\|_r^{r-p+1}. \quad (3.4)$$

From the counter-proof method, it is inevitable to exist  $\varphi$  such that  $\Phi = \varphi - \varphi_\Omega$ . Taken  $\Phi$  as a test function of the integral identity (1.2), that is

$$\int_{\{|u-\theta|>L\}} \langle |\nabla u|^{p-2} \nabla u, |\nabla u - \nabla \theta|^{r-p} (\nabla u - \nabla \theta) \rangle dx = \int_{\{|u-\theta|>L\}} \langle |\nabla u|^{p-2} \nabla u, h \rangle dx + \int_{\{|u-\theta|>L\}} f(x) \Phi dx.$$

This implies

$$\begin{aligned}
& \int_{\{|u-\theta|>L\}} \langle |\nabla u|^{p-2} \nabla u - |\nabla \theta|^{p-2} \nabla \theta, |\nabla u - \nabla \theta|^{r-p} (\nabla u - \nabla \theta) \rangle dx \\
&= \int_{\{|u-\theta|>L\}} \langle |\nabla u|^{p-2} \nabla u - |\nabla \theta|^{p-2} \nabla \theta, h \rangle dx \\
&\quad + \int_{\{|u-\theta|>L\}} \langle |\nabla \theta|^{p-2} \nabla \theta, h \rangle dx \\
&\quad - \int_{\{|u-\theta|>L\}} \langle |\nabla \theta|^{p-2} \nabla \theta, |\nabla u - \nabla \theta|^{r-p} (\nabla u - \nabla \theta) \rangle dx \\
&\quad + \int_{\{|u-\theta|>L\}} f(x) \Phi dx \\
&= I_1 + I_2 + I_3 + I_4.
\end{aligned} \tag{3.5}$$

Now we shall distinguish between two cases.

Case 1:  $p \geq 2$ . using Lemma 2.1, (3.5) can be estimated as

$$\begin{aligned}
& \int_{\{|u-\theta|>L\}} \langle |\nabla u|^{p-2} \nabla u - |\nabla \theta|^{p-2} \nabla \theta, |\nabla u - \nabla \theta|^{r-p} (\nabla u - \nabla \theta) \rangle dx \\
&\geq 2^{2-p} \int_{\{|u-\theta|>L\}} |\nabla u - \nabla \theta|^r dx.
\end{aligned} \tag{3.6}$$

Using the Lemma 2.2, Hölder inequality and Young inequality,  $|I_1|$  can be estimated as

$$\begin{aligned}
|I_1| &= \left| \int_{\{|u-\theta|>L\}} \langle |\nabla u|^{p-2} \nabla u - |\nabla \theta|^{p-2} \nabla \theta, h \rangle dx \right| \\
&\leq (p-1) \int_{\{|u-\theta|>L\}} (|\nabla \theta| + |\nabla u - \nabla \theta|)^{p-2} |\nabla u - \nabla \theta| |h| dx \\
&\leq 2^{p-2} (p-1) \left( \int_{\{|u-\theta|>L\}} |\nabla \theta|^{p-2} |\nabla u - \nabla \theta| |h| dx + \int_{\{|u-\theta|>L\}} |\nabla u - \nabla \theta|^{p-1} |h| dx \right) \\
&\leq 2^{p-2} (p-1) \left[ \left( \int_{\{|u-\theta|>L\}} |\nabla \theta|^r dx \right)^{\frac{p-2}{r}} \left( \int_{\{|u-\theta|>L\}} |\nabla u - \nabla \theta|^r dx \right)^{\frac{1}{r}} \right. \\
&\quad \cdot \left. \left( \int_{\{|u-\theta|>L\}} |h|^{\frac{r}{r-p+1}} dx \right)^{\frac{r-p+1}{r}} + \left( \int_{\{|u-\theta|>L\}} |\nabla u - \nabla \theta|^r dx \right)^{\frac{p-1}{r}} \cdot \left( \int_{\{|u-\theta|>L\}} |h|^{\frac{r}{r-p+1}} dx \right)^{\frac{r-p+1}{r}} \right] \\
&\leq 2^{p-2} (p-1) C(n, p) |p-r| \left[ \left( \int_{\{|u-\theta|>L\}} |\nabla \theta|^r dx \right)^{\frac{p-2}{r}} \right. \\
&\quad \cdot \left. \left( \int_{\{|u-\theta|>L\}} |\nabla u - \nabla \theta|^r dx \right)^{\frac{r-p+2}{r}} + \int_{\{|u-\theta|>L\}} |\nabla u - \nabla \theta|^r dx \right]
\end{aligned} \tag{3.7}$$

Using the Hölder inequality, (3.4) and Young inequality,  $|I_2|$  and  $|I_3|$  can be estimated as

$$\begin{aligned}
|I_2| &= \left| \int_{\{|u-\theta|>L\}} \langle |\nabla \theta|^{p-2} \nabla \theta, h \rangle dx \right| \\
&\leq \int_{\{|u-\theta|>L\}} |\nabla \theta|^{p-1} |h| dx \\
&\leq \left( \int_{\{|u-\theta|>L\}} |\nabla \theta|^r dx \right)^{\frac{p-1}{r}} \left( \int_{\{|u-\theta|>L\}} |h|^{\frac{r}{r-p+1}} dx \right)^{\frac{r-p+1}{r}} \\
&\leq C(n, p) |p-r| \left( \int_{\{|u-\theta|>L\}} |\nabla \theta|^r dx \right)^{\frac{p-1}{r}} \cdot \left( \int_{\{|u-\theta|>L\}} |\nabla u - \nabla \theta|^r dx \right)^{\frac{r-p+1}{r}} \\
&\leq C(n, p) |p-r| [C(\varepsilon) \int_{\{|u-\theta|>L\}} |\nabla \theta|^r dx + \varepsilon \int_{\{|u-\theta|>L\}} |\nabla u - \nabla \theta|^r dx],
\end{aligned} \tag{3.8}$$

$$\begin{aligned}
|I_3| &= \left| -\int_{\{|u-\theta|>L\}} \langle |\nabla\theta|^{p-2} \nabla\theta, |\nabla u - \nabla\theta|^{r-p} (\nabla u - \nabla\theta) \rangle dx \right| \\
&\leq \int_{\{|u-\theta|>L\}} |\nabla\theta|^{p-1} |\nabla u - \nabla\theta|^{r-p+1} dx \\
&\leq \left( \int_{\{|u-\theta|>L\}} |\nabla\theta|^r dx \right)^{\frac{p-1}{r}} \left( \int_{\{|u-\theta|>L\}} |\nabla u - \nabla\theta|^r dx \right)^{\frac{r-p+1}{r}} \\
&\leq C(\varepsilon) \int_{\{|u-\theta|>L\}} |\nabla\theta|^r dx + \varepsilon \int_{\{|u-\theta|>L\}} |\nabla u - \nabla\theta|^r dx.
\end{aligned} \tag{3.9}$$

Using the Hölder inequality, Sobolev-Poincaré inequality<sup>[7]</sup>,

$$\left( \int_{\Omega} |u - u_{\Omega}|^{pm/(n-p)} dx \right)^{(n-p)/pm} \leq C \left( \int_{\Omega} |\nabla u|^p dx \right)^{1/p}, (1 \leq p < n),$$

and using (3.3) and Young inequality,  $|I_4|$  can be estimated as

$$\begin{aligned}
|I_4| &= \left| \int_{\{|u-\theta|>L\}} f(x)\Phi dx \right| \\
&\leq \left( \int_{\{|u-\theta|>L\}} |f(x)|^{\frac{nr}{n(p-1)+r}} dx \right)^{\frac{n(p-1)+r}{nr}} \cdot \left( \int_{\{|u-\theta|>L\}} |\varphi - \varphi_{\Omega}|^{\frac{nr}{n(r-p+1)-r}} dx \right)^{\frac{n(r-p+1)-r}{nr}} \\
&\leq C(n, p) \left( \int_{\{|u-\theta|>L\}} |f(x)|^{\frac{nr}{n(p-1)+r}} dx \right)^{\frac{n(p-1)+r}{nr}} \cdot \left( \int_{\{|u-\theta|>L\}} |\nabla\Phi|^{\frac{r}{r-p+1}} dx \right)^{\frac{r-p+1}{r}} \\
&\leq C(n, p) \left( \int_{\{|u-\theta|>L\}} |f(x)|^{\frac{nr}{n(p-1)+r}} dx \right)^{\frac{n(p-1)+r}{nr}} \cdot \left( \int_{\{|u-\theta|>L\}} |\nabla v|^r dx \right)^{\frac{r-p+1}{r}} \\
&\leq C(n, p) [C(\varepsilon) \left( \int_{\{|u-\theta|>L\}} |f(x)|^{\frac{nr}{n(p-1)+r}} dx \right)^{\frac{n(p-1)+r}{nr}} \\
&\quad + \varepsilon \int_{\{|u-\theta|>L\}} |\nabla u - \nabla\theta|^r dx].
\end{aligned} \tag{3.10}$$

Combining (3.5)-(3.10), we arrive at

$$\begin{aligned}
&\int_{\{|u-\theta|>L\}} |\nabla u - \nabla\theta|^r dx \\
&\leq C(n, p, \varepsilon) \int_{\{|u-\theta|>L\}} |\nabla\theta|^r dx \\
&\quad + (C(n, p) |p-r| + \varepsilon) \int_{\{|u-\theta|>L\}} |\nabla u - \nabla\theta|^r dx \\
&\quad + C(n, p, \varepsilon) \left( \int_{\{|u-\theta|>L\}} |f(x)|^{\frac{nr}{n(p-1)+r}} dx \right)^{\frac{n(p-1)+r}{n(p-1)}},
\end{aligned} \tag{3.11}$$

Case 2:  $1 < p < 2$ . Lemma 2.3 yields

$$\begin{aligned}
&\int_{\{|u-\theta|>L\}} \langle |\nabla u|^{p-2} \nabla u - |\nabla\theta|^{p-2} \nabla\theta, |\nabla u - \nabla\theta|^{r-p} (\nabla u - \nabla\theta) \rangle dx \\
&\geq \int_{\{|u-\theta|>L\}} |\nabla u - \nabla\theta|^{r-p+1} \\
&\quad \cdot (|\nabla u - \nabla\theta| + |\nabla\theta|)^{p-1} - |\nabla\theta|^{p-1} dx.
\end{aligned}$$

This implies

$$\begin{aligned}
& \int_{\{|u-\theta|>L\}} |\nabla u - \nabla \theta|^r dx \\
& \leq \int_{\{|u-\theta|>L\}} |\nabla u - \nabla \theta|^{r-p+1} (|\nabla u - \nabla \theta| + |\nabla \theta|)^{p-1} dx \\
& \leq \int_{\{|u-\theta|>L\}} \langle |\nabla u|^{p-2} \nabla u - |\nabla \theta|^{p-2} \nabla \theta, |\nabla u - \nabla \theta|^{r-p} (\nabla u - \nabla \theta) \rangle dx \\
& \quad + \int_{\{|u-\theta|>L\}} |\nabla u - \nabla \theta|^{r-p+1} |\nabla \theta|^{p-1} dx \\
& \leq \int_{\{|u-\theta|>L\}} \langle |\nabla u|^{p-2} \nabla u - |\nabla \theta|^{p-2} \nabla \theta, |\nabla u - \nabla \theta|^{r-p} (\nabla u - \nabla \theta) \rangle dx \\
& \quad + \varepsilon \int_{\{|u-\theta|>L\}} |\nabla u - \nabla \theta|^r dx + C(\varepsilon) \int_{\{|u-\theta|>L\}} |\nabla \theta|^r dx.
\end{aligned} \tag{3.12}$$

Using Lemma 2.2 and (3.4),  $|I_1|$  can be estimated as

$$\begin{aligned}
|I_1| & = \left| \int_{\{|u-\theta|>L\}} \langle |\nabla u|^{p-2} \nabla u - |\nabla \theta|^{p-2} \nabla \theta, h \rangle dx \right| \\
& \leq \frac{3-p}{2^{p-2}(p-1)} \int_{\{|u-\theta|>L\}} |\nabla u - \nabla \theta|^{p-1} |h| dx \\
& \leq \frac{3-p}{2^{p-2}(p-1)} \left( \int_{\{|u-\theta|>L\}} |\nabla u - \nabla \theta|^r dx \right)^{\frac{p-1}{r}} \cdot \left( \int_{\{|u-\theta|>L\}} |h|^{\frac{r}{r-p+1}} dx \right)^{\frac{r-p+1}{r}} \\
& \leq \frac{3-p}{2^{p-2}(p-1)} C(n, p) |p-r| \int_{\{|u-\theta|>L\}} |\nabla u - \nabla \theta|^r dx.
\end{aligned} \tag{3.13}$$

For the case  $1 < p < 2$ ,  $|I_2| - |I_3|$  can also be estimated by (3.8)-(3.9). Combining (3.5), (3.12) and (3.13), we arrive at (3.11).

Let  $\varepsilon_0 = 1/C(n, p)$ , Then for  $|p-r| < \varepsilon_0$  we have  $C(n, p)|p-r| < 1$ , Taking  $\varepsilon$  small enough, such that  $C(n, p)|p-r| + \varepsilon < 1$ , then the second term on the right-hand side of (3.11) can be absorbed by the left-hand side; thus we obtain

$$\begin{aligned}
& \int_{\{|u-\theta|>L\}} |\nabla u - \nabla \theta|^r dx \\
& \leq C(n, p) \int_{\{|u-\theta|>L\}} |\nabla \theta|^r dx + C(n, p) \left( \int_{\{|u-\theta|>L\}} |f(x)|^{\frac{nr}{n(p-1)+r}} dx \right)^{\frac{n(p-1)+r}{n(p-1)}}.
\end{aligned} \tag{3.14}$$

Since  $\theta \in W^{1,q}(\Omega)$ ,  $q > r$ , using the Hölder inequality, we have

$$\begin{aligned}
& \int_{\{|u-\theta|>L\}} |\nabla \theta|^r dx \\
& \leq \left( \int_{\{|u-\theta|>L\}} |\nabla \theta|^q dx \right)^{r/q} |\{|u-\theta|>L\}|^{(q-r)/q} \\
& = \|\nabla \theta\|_q^r |\{|u-\theta|>L\}|^{(q-r)/q}.
\end{aligned} \tag{3.15}$$

By the proof idea of reference [9](Page 442), and the Hölder inequality, we get

$$\begin{aligned}
& \left( \int_{\{|u-\theta|>L\}} |f(x)|^{\frac{nr}{n(p-1)+r}} dx \right)^{\frac{n(p-1)+r}{n(p-1)}} \\
& \leq \left( \int_{\{|u-\theta|>L\}} |f(x)|^{\frac{nr}{n(p-1)+r}} dx \right)^{\frac{n(p-1)+r^2}{qn(p-1)}} |\{|u-\theta|>L\}|^{(q-r)/q} \\
& \leq M |\{|u-\theta|>L\}|^{(q-r)/q},
\end{aligned} \tag{3.16}$$

where  $M = \left( \int_{\{|u-\theta|>L\}} |f(x)|^{\frac{nq}{n(p-1)+r}} dx \right)^{\frac{nr(p-1)+r^2}{qn(p-1)}}$ ,  $M$  is bounded and is a constant dependent only on  $n$ ,  $p$ . Then (3.14) can be collated into the following results

$$\begin{aligned} & \int_{\{|u-\theta|>L\}} |\nabla u - \nabla \theta|^r dx \\ & \leq C(n, p) \left( \int_{\{|u-\theta|>L\}} |\nabla \theta|^q dx \right)^{r/q} |\{|u-\theta|>L\}|^{(q-r)/q} \\ & \quad + C(n, p) M |\{|u-\theta|>L\}|^{(q-r)/q} \\ & = C |\{|u-\theta|>L\}|^{(q-r)/q} (1 + \|\nabla \theta\|_q^r), \end{aligned} \quad (3.17)$$

where  $C = C(n, p, M)$ .

We now turn our attention back to the function  $v \in W_0^{1,r}(\Omega)$ . By the Sobolev embedding theorem, we have

$$\begin{aligned} \left( \int_{\Omega} |v|^{r^*} dx \right)^{1/r^*} & \leq C(n, r) \left( \int_{\Omega} |\nabla v|^r dx \right)^{1/r} \\ & = C(n, r) \left( \int_{\{|u-\theta|>L\}} |\nabla u - \nabla \theta|^r dx \right)^{1/r}, \end{aligned} \quad (3.18)$$

since  $|v| = (|u-\theta| - L) \cdot \mathbf{1}_{\{|u-\theta|>L\}}$ , we have

$$\left( \int_{\{|u-\theta|>L\}} (|\nabla u - \nabla \theta| - L)^{r^*} dx \right)^{1/r^*} = \left( \int_{\Omega} |v|^{r^*} dx \right)^{1/r^*}, \quad (3.19)$$

and for  $\tilde{L} > L$ ,

$$\begin{aligned} & (\tilde{L} - L)^{r^*} |\{|u-\theta|>\tilde{L}\}| \\ & = \int_{\{|u-\theta|>\tilde{L}\}} (\tilde{L} - L)^{r^*} dx \\ & \leq \int_{\{|u-\theta|>\tilde{L}\}} (|u-\theta| - L)^{r^*} dx \\ & \leq \int_{\{|u-\theta|>L\}} (|u-\theta| - L)^{r^*} dx. \end{aligned} \quad (3.20)$$

By collecting (3.17)-(3.20), we deduce that

$$\begin{aligned} & ((\tilde{L} - L)^{r^*} |\{|u-\theta|>\tilde{L}\}|)^{1/r^*} \\ & \leq C(n, r) (\|\nabla \theta\|_q + 1) |\{|u-\theta|>L\}|^{1/r-1/q} \end{aligned} \quad (3.21)$$

Thus

$$\begin{aligned} & |\{|u-\theta|>\tilde{L}\}| \\ & \leq \frac{1}{(\tilde{L} - L)^{r^*}} (C(n, r) (\|\nabla \theta\|_q + 1))^{r^*} |\{|u-\theta|>L\}|^{r^*(1/r-1/q)} \end{aligned} \quad (3.22)$$

Let  $\phi(s) = |\{|u-\theta|>s\}|$ ,  $\alpha = r^*$ ,  $c = (C(n, r) (\|\nabla \theta\|_q + 1))^{r^*}$ ,  $\beta = r^*(1/r-1/q)$ ,  $s_0 > 0$ , Then (3.22) become

$$\phi(\tilde{L}) \leq \frac{c}{(\tilde{L} - L)^\alpha} \phi(L)^\beta \quad (3.23)$$

for  $\tilde{L} > L > 0$ .

(1) For the case  $q < n$ , one has  $\beta < 1$ . In this case, if  $s \geq 1$ , we get from Lemma 2.3 that

$$|\{|u-\theta|>s\}| \leq c(\alpha, \beta, s_0) s^{-\tau},$$

where  $t = \alpha/(1-\beta) = q^*$ . For  $0 < s < 1$ , one has

$$|\{|u - \theta| > s\}| \leq |\Omega| = |\Omega| s^{q^*} s^{-q^*} \leq |\Omega| s^{-q^*}.$$

Thus

$$u \in \theta + L_{weak}^{q^*}(\Omega).$$

(2) For the case  $q = n$ , one has  $\beta = 1$ . For any  $\tau < \infty$ , (3.23) implies

$$\begin{aligned} \phi(\tilde{L}) &\leq \frac{c}{(\tilde{L}-L)^\alpha} \phi(L) = \frac{c}{(\tilde{L}-L)^\alpha} \phi(L)^{1-\alpha/\tau} \phi(L)^{\alpha/\tau} \\ &\leq \frac{c|\Omega|^{\alpha/\tau}}{(\tilde{L}-L)^\alpha} \phi(L)^{1-\alpha/\tau}. \end{aligned}$$

As about, we derive

$$u \in \theta + L_{weak}^r(\Omega).$$

(3) For the case  $q > n$ , one has  $\beta > 1$ . Lemma 2.3 implies  $\phi(d) = 0$  for some  $d = d(\alpha, \beta, s_0, r, (\|\nabla\theta\|_q + 1))$ . Thus  $|\{|u - \theta| > d\}| = 0$ , which means  $u - \theta \leq d$  a.e. in  $\Omega$ , Therefore

$$u \in \theta + L^\infty(\Omega),$$

completing the proof of Theorem 1.1.

## References

- [1] T. Iwaniec, "p-harmonic tensors and quasiregular mappings," Ann. Math, 1992, 136(2): 589-624.
- [2] Hongya Gao, Shuang Liang, Yi Cui, "Integrability for very weak solutions to boundary value problems of p-harmonic equation," 2016, 66(141): 101-110.
- [3] Kunjie Zhu, Shuhong Chen, "The properties of very weak solutions of nonhomogeneous A -harmonic equations," Fujian: Minnan normal University, 2017.
- [4] Kaili Guo, Hongya Gao, "Functional minima and integrability of solutions of elliptic differential equations," Baoding: Hebei University, 2017.
- [5] Shicong Zhang, Shenzhou Zheng, "Regularity of generalized solutions of Dirichlet boundary value problems for two classes of elliptic equations," Beijing: Beijing Jiaotong University, 2018.
- [6] L. Greco, T. Iwaniec, C. Sbordone, "Inverting the p-harmonic operator," Manuscr. Math. 1977, 92: 249-258.
- [7] T. Iwaniec, L. Migliaccio, L. Nania, C. Sbordone, "Integrability and removability results for quasiregular mappings in high dimensions," Math. Scand, 1994, 75: 263-279.
- [8] Reshetnyak Yu.G, "Space mappings with bounded distortion," Vol. 173, Trans. Math. Mokeygraphs, Amer. Soc., 1989.
- [9] Hongya Gao, Qinghua Di, Dongna Ma, "Integrability for solutions to some anisotropic obstacle problems," 2015, 146: 433-444