

A COMPARATIVE STUDY ON MEASUREMENT OF MAXILLARY OCCLUSAL CANT OBTAINED THROUGH FACEBOW TRANSFER AND USING LATERAL CEPH. : AN IN VIVO STUDY

Dr Rohit Saini^{1*}, Dr Rajeev Gupta², Dr Archana Nagpal³, Dr Vishal Katna⁴, Dr kusha Thakur⁵, Dr Payal kashyap⁶

¹ -Postgraduate Student, Department of Prosthodontics, Himachal Dental College, Himachal Pradesh

² - Dr Rajeev Gupta, MDS, Professor, Department of Prosthodontics, Himachal Dental College, Himachal Pradesh.

³ -Dr Archana Nagpal, MDS, Professor & Head, Department of Prosthodontics, Himachal Dental College, Himachal Pradesh.

⁴ -Dr Vishal Katna, MDS, Reader, Department of Prosthodontics, Himachal Dental College, Himachal Pradesh.

⁵ -Dr Kusha Thakur, Postgraduate Student, Department of Prosthodontics, Himachal Dental College, Himachal Pradesh.

⁶ -Dr Payal Kashyap, Postgraduate Student, Department of Prosthodontics, Himachal Dental College, Himachal Pradesh

ABSTRACT

Purpose: The purpose of the study was to compare measurements of maxillary occlusal cant obtained through facebow transfer and through lateral cephalogram.

Method: 40 subjects were included in this study according to inclusion and exclusion criteria and divided into two groups: dentulous and edentulous. For edentulous subjects, all the steps of complete denture fabrication were carried out and finished dentures were delivered. Alginate impressions were now made for all the subjects (with complete denture worn in edentulous cases) for both upper and lower arches. The study was conducted in two parts. Facebow transfer was done next and casts were mounted. In first part of the study, sagittal inclination was measured after facebow transfer. After

mounting of the casts, four points were marked to measure the inclination of the occlusal plane. In second part, cephalometric evaluation of occlusal plane and Frankfurt horizontal plane was carried out. Angle between Frankfurt horizontal plane and the occlusal plane was maxillary occlusal cant. which was evaluated by tracing. Paired t test was used to compare mean facebow values and lateral ceph values in edentulous subjects. Intergroup comparison between lateral ceph and mean facebow values between dentulous and edentulous subjects was evaluated using independent t test.

Results: Facebow measurements gave comparatively higher values in both dentulous and edentulous patients and are subjected to less variation as compared to the lateral cephalogram values $p < 0.0001$.

Conclusion: The occlusal plane angle of lateral cephalogram was found to be significantly different from angle obtained through facebow transfer.

KEYWORDS

Occlusal cant, facebow, frankfurt horizontal plane, lateral cephalogram

1. INTRODUCTION

In complete denture construction, the Prosthodontist is responsible for restoring the natural esthetics of the patient and for developing an occlusion that is compatible with functional movements of the mandible.^[1] One of the salient factor that help us in developing occlusion which is compatible with the functional movement of the stomatognathic system is the orientation of occlusal plane.^[2] Occlusal plane orientation is one of the most important clinical procedure in removable prosthodontic treatment for edentulous patients.^[3]

Ideally the occlusal plane should be located in a direction perpendicular to the occlusal bite force. This position provides stability to dentures supported by underlying resilient tissue. Functionally the occlusal table is a milling surface that is designed in such a manner so that the tongue and the buccinator muscle are able to position the food bolus onto it and hold it there during the process of mastication.

To orient the maxillary arch and dentition using a facebow, involves a plane of reference, ie, the Frankfurt horizontal plane (porion-orbitale), which appears horizontal when the head is placed in the

natural head position.^[2] A facebow is used to record the antero-posterior and vertical relationship of the maxilla to the hinge axis of the temporomandibular joints and to transfer this relationship to the opening axis of an articulator.^[4] The proper use of an anatomic articulator is dependent upon an accurate facebow transfer.^[5] The third point of reference recommended for the Hanau Wide-View model 183-2 semiadjustable articulator is, orbitale.^[6]

A lateral cephalogram reveals areas in a cranial base that are not subjected to alteration, it is used in identifying predictable relationships between the teeth and other cranial landmarks, henceforth it is considered as the gold standard.^[2] Cephalometric analysis is an important diagnostic tool in dentistry, in prosthodontics, the significance of cephalometrics lies in the ability to re-establish the spatial position of lost structures (such as the teeth).^[7] In complete denture fabrication, recording a correct jaw relationship is of utmost importance and occlusal plane record is a part of the same. Hence, the purpose of the study was to compare measurements of maxillary occlusal cant obtained through facebow transfer and through lateral cephalogram.

2. MATERIALS AND METHODS

The study included 20 dentulous and 20 edentulous subjects comprising both males and females randomly selected who visited the out-patient department of Prosthodontics. All the procedures were carried out in Department of Prosthodontics. All the subjects were informed about the study and institutional ethical clearance was also obtained.

Inclusion criteria (dentulous patients):

- Age group: 18-30 years with completed facial growth
- Full complement of healthy and natural teeth
- No history of orthodontic treatment

Exclusion criteria (dentulous patients):

- Periodontally compromised teeth
- Teeth grossly attrited or abraded
- Presence of fixed or removable partial dentures
- Gross malalignment of teeth

Inclusion criteria (edentulous patients):

- Normal ridge relationship
- Well-formed ridge

Exclusion criteria (edentulous patients):

- Resorbed ridge

Reference planes:

- Frankfurt horizontal plane.
- Occlusal plane: Plane touching mesiopalatal cusp of left maxillary first molar and left mesioincisal edge of central incisor.

2.1 METHODOLOGY:

Subjects, both dentulous as well as edentulous, were selected randomly keeping in mind the specified inclusion criteria. For edentulous subjects, all the steps of complete denture fabrication were carried out and finished dentures were delivered.

Following this, alginate impressions (Algitex, Mumbai) were now made for all the 40 subjects (with complete denture worn in edentulous cases) for both upper and lower arches followed by pouring of casts in Type III gypsum (Kalstone, Kalabhai Karson Pvt Ltd, Mumbai). Facebow transfer was done next and casts were mounted (Figure 1-3).




Figure 1 a -Facebow transfer in dentulous patient



Figure 1b- Facebow transfer in edentulous patient




Figure 2- Facebow with Bite transferred on to the articulator



Figure 3- Mounting of maxillary cast on articulator

Standard mounting procedure was followed as per the manufacturer's instructions. The study was done in two parts:

2.1.1. FIRST PART (Measurement of sagittal inclination after facebow transfer)

After mounting of the casts, four points were marked to measure the inclination of the occlusal plane. Two marks were marked on the U-shaped frame of facebow of Hanau articulators (Hanau Wide-View model 183-2 semi-adjustable articulator) (figure 4). These were as follows:



Figure 4- Points I and C marked on U shaped frame of facebow

- **Point C** – A point near the condylar axis on the upper surface of U frame. This was done by sticking surgical tape on the area and marking two lines which bisected each other at right angles; the point of intersection of these lines was taken as point C.
- **Point I** – A point close to third point reference i.e. orbitale on the upper surface of U frame on the left side of face. The markings were done in the same way as described for point C. The point of intersection of the two lines was taken as point I.

The plane formed by C and I was corresponded to Frankfurt horizontal plane. These two point marks were the stationary reference points from which all measurements were recorded.

On the articulated casts, two points were taken: one point on the mesiopalatal cusp of left upper molar (point M) and the other on the mesioincisal edge of the left upper central incisor (point A).

For the ease of measurement, a steel plate was fixed above the bite fork with an adhesive (Figure 5).



Figure 5- Steel plate

This represented the occlusal plane and placed below the maxillary cast touching the incisal edge and mesiopalatal cusp, followed by the marking of these points on a steel plate. Now the marked points were reproduced on the left border of the steel plate by drawing perpendicular lines extending to one side. The points were marked as A and M on steel plate placed right under the one side of the frame of the facebow. A and M points corresponded to the occlusal plane (Figure 6).



Figure 6- Points reproduced on the left border of the steel plate by drawing perpendicular lines

With the help of a pair of dividers, linear distances were measured as follows:

- Distance between the C (point near the condyle) and the I (point near orbitale) (CI);
- Distance between C and the point M (mesiopalatal cusp) on steel plate (CM);
- Distance between C and the point A (mesioincisal egde) on steel plate (CA);
- Distance between I and the point M (IM); and
- Distance between I and the point on mesioincisal edge A of central incisor (IA).

These values were then plotted on a graph paper (Figure 7 a-b).




Figure 7a- Markings plotted on graph paper of dentulous patient




Figure 7b- Markings plotted on graph paper of edentulous patient

The angle formed between lines CI and MA represented the horizontal plane and the occlusal plane respectively, therefore, an angle formed was maxillary occlusal cant obtained through facebow transfer.

2.1.2. SECOND PART (Procedure for cephalometric evaluation of occlusal plane and Frankfurt horizontal plane)

Before cephalometric evaluation, a piece of lead foil was placed (dentulous patients using composite and edentulous patients using adhesive) on the mesioincisal edge of central incisor and mesiopalatal cusp of molar (Figure 8).



Figure 8- Lead foil fixed on mesiopalatal cusp of maxillary left molar and mesioincisal edge of maxillary left central incisor using composite

Following lead foil placement, the lateral cephalogram was taken for all the subjects (patient wearing denture in edentulous patients) with Frankfurt horizontal plane parallel to the ground in a cephalostat (Planmeca X- ray machine, model 2002). Tracing was done to evaluate the angle between Frankfurt horizontal plane and the occlusal plane (formed by line joining the mesiopalatal cusp of upper molar and incisal edge of central incisor) (Figure 9 a-b).



Figure 9a- Tracing of lateral cephalogram of dentulous patient



Figure 9b- Tracing of lateral cephalogram of edentulous patient

Therefore, the angle formed was maxillary occlusal cant.

3. RESULTS

The study was conducted in the Department of Prosthodontics. 40 cases (20 dentulous and 20 edentulous) were selected keeping in view of inclusion criteria. The values of maxillary occlusal cant using facebow as well as lateral ceph for both the groups (dentulous and edentulous) were sent for statistical analysis. The results obtained are shown in Tables 1-5.

The descriptive statistics of lateral ceph and face bow values of Dentulous and Edentulous patients are presented in table 1 and table 3. For Dentulous patients the mean Face bow value was found to be significantly higher as compared to the lateral Ceph value ($P < .001$). This is evaluated by paired t test and the summary results of the significance level are presented in Table 2.

Even in the case of Edentulous patients the mean facebow value was significantly higher in comparison to the Lateral Ceph value ($P < 0.0001$) as observed by paired t test. The summary result of this significance test is presented in Table 4.

Table 1. Descriptive statistics of measurements in Dentulous patients (n = 20)

					95% Confidence Interval for Mean			
	N	Mean	Std. Deviation	Std. Error	Lower Bound	Upper Bound	Minimum	Max

LateralCeph	20	8.33	2.40	0.54	7.20	9.45	4	12.5
Facebow	20	10.48	3.39	0.76	8.89	12.06	5	16

Table 2. paired comparison between Lateral Ceph and Face bow values in dentulous patients (N= 20)

Paired Samples Statistics					
		Mean	N	Std. Deviation	Std. Error Mean
Pair 1	LateralCeph	8.3250	20	2.39668	.53591
	Facebow	10.4750	20	3.39301	.75870

Paired Samples Correlations				
		N	Correlation	Sig.
Pair 1	Lateral Ceph & Facebow	20	.719	.000

Paired t test			
Std. Error Mean	t	df	Sig (2-tailed)
.52703	-4.079	19	.001

Table 3. Descriptive statistics of measurements in Edentulous patients (n = 20)

					95% Confidenc e Interval for Mean			
	N	Mean	Std. Deviation	Std. Error	Lower Bound	Upper Bound	Minimum	Max
Facebow	20	11.30	2.34	0.52	10.20	12.40	7	16
Lat Ceph	20	9.70	2.32	0.52	8.61	10.79	6	15

Table 4. paired comparison between Lateral Ceph and Face bow values in Edentulous patients (N= 20)

Paired Samples Statistics					
		Mean	N	Std. Deviation	Std. Error Mean

Pair 1	LateralCeph	9.7000	20	2.31926	.51860				
	Facebow	11.3000	20	2.34184	.52365				
Paired Samples Correlations									
		N	Correlation	Sig.					
Pair 1	LateralCeph & Facebow	20	.778	.000					
Paired Samples Test									
		Mean	Std. Dev	Std. Error Mean	95% Confidence Interval of the Difference		T	d f	Sig (2-tiled)
					Lower	Upper			
Pair 1	LateralCeph - Facebow	-1.60	1.55	.34717	-2.33	-.87	- 4.61	19	.0001

The lateral cephalometric values obtained in dentulous and edentulous patients and facebow values obtained in dentulous and edentulous patients are further compared for inter group comparison, by independent t test method. The results are presented in Table 5. The results revealed that the mean of lateral cephalometric values no differ among two groups of patients as well as no such differences were observed for facebow values when compared among the dentulous and edentulous patients.

Table 5. Comparison of two parameters among dentulous and edentulous patients- results of independent sample t test

Parameters	Groups	Mean	Sd	t	df	Sig. (2-tailed)	Std. Error Difference
LateralCeph	Dentulous	8.32	2.40	-1.84	38	0.73	0.746
	Edentulous	9.70	2.32				
Facebow	Dentulous	10.5	3.39	0.895	38	0.380	1.04
	Edentulous	11.3	2.34				

The values for edentulous patient appeared to be lower than dentulous patients. The values are compared with independent sample t test. The two-tailed P value equals 0.3891. By conventional criteria, this difference is considered to be not statistically significant. The intermediate values used in calculations are $t = 0.871$, $df = 38$ and standard error of difference = 0.631. It appeared that facebow measurements gave comparatively higher values in both dentulous and edentulous patients and these values are subjected to less variation as compared to the lateral cephalometric values.

4. DISCUSSION

The present study was conducted in the Department of Prosthodontics, patients who met the needs of the inclusion criteria were randomly selected and divided into two groups i.e. a group of dentulous and other group of edentulous patients. Complete dentures were fabricated and delivered to patients in the edentulous group prior to the analysis and measurements.

In the study, sagittal inclination of the occlusal plane of articulated maxillary casts to the horizontal reference plane using facebow was evaluated and compared with the cephalometric occlusal cant for both the groups of patients

Maxillary models were mounted on a semi adjustable articulator following facebow transfer. This was followed by making physical measurements, to determine the inclination of the maxillary occlusal plane with respect to the horizontal reference line i.e Frankfurt horizontal plane.

Degree of occlusal cant on the lateral cephalograms was also evaluated. All lateral cephalometric films were placed on transparent cellulose acetate sheet of 54 μ thickness

The data obtained from the articulator and the lateral ceph were subjected to statistical analysis. Following were the main observations made:

The maximum angle measured on cephalogram for edentulous patients was 16°, whereas the minimum was 7°, with the mean angle evaluated $11.30^{\circ} \pm 2.34^{\circ}$.

The maximum angle measured on the articulated cast using facebow 15°, whereas the minimum angle was 6°, with the mean angle calculated was $9.70^{\circ} \pm 2.32^{\circ}$.

The maximum angle measured on cephalogram for dentulous patients was 12.5°, whereas the minimum was 4°, with the mean angle being $8.33^{\circ} \pm 2.40^{\circ}$ for this study.

In the study carried out by **Shetty et al., (2016)**^[2], the Frankfurt horizontal plane-occlusal plane angle for lateral cephalogram varied from a maximum of 13.3° to a minimum of 3.5° with a mean of $8.7^{\circ} \pm 2.24^{\circ}$ thereby showing similar results as shown in the current study.

According to the study by **Rupal J Shah et al., (2013)**^[8], minimum angle value for lateral ceph was 3° and maximum was 17° mean value was $9.13^{\circ} \pm 3.77$.

In another study conducted by **Nazir et al., (2012)^[9]**, the maximum angle measured on cephalogram was 15° , whereas the minimum was 6° , with the mean angle being $9.61^{\circ} \pm 2.55$.

The mean occlusal plane angle in cephalogram was $10.4^{\circ} \pm 4.3$, which was slightly higher in the study by **Kyung Suk Seo., (2003)^[10]** as compared to the present study.

On the casts that were mounted on hanau wide vue articulator using facebow for dentulous patients, the maximum angle measured was 16° and the minimum was 5° . The mean angle was calculated to be $10.48^{\circ} \pm 3.39$.

This result is in accordance with the study carried out by **Shetty et al., (2016)^[2]**, in which the Frankfurt horizontal plane - Occlusal plane angle using Hanau Wide-Vue group, varied from a maximum of 15° to a minimum of 5.1° with a mean of $10.69^{\circ} \pm 2.44^{\circ}$. The study by **Nazier et al., (2012)^[9]** also yielded similar result showing maximum angle of 15° and minimum of 6° . The average angle of sagittal inclination was calculated to be $10.77^{\circ} \pm 2.60^{\circ}$.

The mean angle of sagittal inclination of maxillary cast mounted on Hanau Wide-Vue articulator was, however, higher in the study conducted by **Mohammad Abdullah and Sherfudhin., (1994)^[4]** and a study by **Kyung Suk Seo., (2003)^[10]** who got a mean angle of 13.77° and $13.5^{\circ} \pm 5.4$ respectively. On the other hand, **Rupal J Shah et al., (2013)^[8]**, in their study, got a mean angle of $8.57^{\circ} \pm 3.45$ which was lower than the values in the current study.

The mean difference between the facebow and lateral ceph for dentulous patients in this study is 2.15°

This study showed a mean difference 2.15° between the sagittal inclination of maxillary cast mounted on Hanau wide Vue articulator and the value obtained using lateral ceph.

This result was similar to the results given by **Shetty et al., (2016)^[2]**, who after reported a mean difference of 1.9° between the occlusal cant measured on Hanau wide Vue articulator and lateral ceph.

Nazir et al., (2012)^[9] also showed a mean difference of 1.16° in their study.

Kyung Suk Seo., (2003)^[10] in his study, found a mean difference of $3.3^{\circ} \pm 4.6$ which was higher as compared to this study.

On the contrary, a mean difference of -0.567° was found in a study conducted by **Rupal J Shah et al., (2013)**^[8].

The results showed that the angle formed between the Frankfurt horizontal plane-Occlusal plane in a lateral ceph could be considered more reliable as compared to the measurements done with facebow transfer using articulator.

A lateral ceph is considered as the gold standard as it unveils hard tissue areas in a cranial base. It is used in assessing predictable relationships between the teeth and other cranial landmarks that remain unaffected even post extraction of teeth.

In reality, the Frankfurt horizontal plane is not transferred to the articulator by the use of orbitale pointer. This is because only the anterior point of reference for this plane is used; the orbitale. Porion does not come into play during the face-bow transfer.⁶ As the facebow transfer on articulator is an arbitrary process, there could be chances of errors due to soft tissue involvement, position of anterior reference, mounting of maxillary casts.

If there are errors during the facebow transfer using Hanau Wide-View articulator, it can further leave an impact of the procedures to follow and consequently lead to unreliable result after delivery of the prosthesis.

The various procedures that can get adversely affected due to these errors may range from full mouth rehabilitation procedures and fixed partial dentures to balanced complete denture prosthesis.

Thus, the present study confirms the importance of cephalometry in the field of Prosthodontics to establish plane of occlusion for proper functions of chewing, mastication and also to restore the esthetics of an individual.^[11]

5. CONCLUSION

The present study comprised of 40 patients, 20 dentulous and 20 edentulous who visited the out-patient department of Prosthodontics. The maxillary occlusal cant was evaluated through facebow transfer on semi adjustable articulator and through cephalometrically.

Study was divided into following groups:

- Occlusal cant of dentulous patients through facebow transfer.

- Occlusal cant of dentulous patients through lateral cephalogram.
- Occlusal cant of edentulous patients through facebow transfer.
- Occlusal cant of edentulous patients through lateral cephalogram.

After statistical analysis, the following conclusions were made:

- Within the limitations of this study, it was seen that reproducibility of the occlusal cant on an articulator by a facebow was not exact.
- The sagittal inclination of the mounted maxillary casts on the Hanau Wide-View semi adjustable articulator was closer to the individual's occlusal cant as measured on the cephalogram.
- The correlation value (Pearson's value) obtained between maxillary cast mounted on Hanau Wide-View articulator was greater as compared to the lateral cephalogram.
- Thus, it could be concluded that the occlusal plane angle of lateral cephalogram was significantly different from angle obtained through facebow transfer.

Based on the results of this study, the lateral cephalogram is more reliable as compared to the facebow transfer in determining the maxillary occlusal cant.

ACKNOWLEDGEMENT

I express my heartfelt gratitude to my guides, mentor and my fellow colleagues for their invaluable guidance and assistance through the course of this study.

COMPLETING INTEREST

Authors have no competing interests exists.

AUTHORS' CONTRIBUTIONS

Author 1- executed the study and performed statistical analysis

Author 2 and 3 - designed and planned the study

Author 4- Managed the analysis of study

Author 5 and 6 - Managed literature searches

All the authors have read and approved the final manuscript.

CONSENT

Informed consent to participate was obtained from each patient prior to their enrollment in the study.

ETHICAL APPROVAL

Ethical approval was obtained from institutional and university ethical research cell committee.

REFERENCES

- [1]D Souza NL and Bhargva K. A cephalometric study comparing the occlusal plane in dentulous and edentulous subjects in relation to the maxillomandibular space. J Prosthet Dent 1996; 75:177-82 [https://doi.org/10.1016/S0022-3913\(96\)90096-7](https://doi.org/10.1016/S0022-3913(96)90096-7)
- [2]Shetty S, Shenoy KK and Sabu A. Evaluation of accuracy of transfer of maxillary occlusal cant of two articulators using two facebow/ semi-adjustable articulator system: An in vivo study. J Indian Prosthodont Soc 2016; 16:248-52 <https://doi.org/10.4103/0972-4052.176525>
- [3]Quran FAM, AL Quran, Hazza'a A and Nahass N Al. The Position of the Occlusal Plane in Natural and Artificial Dentitions as Related to Other Craniofacial Planes. Journal of Prosthodontics (2010) 601–5 <https://doi.org/10.1111/j.1532-849X.2010.00643>
- [4]Abdullah MA, Sherfudhin H. A comparative study of Facebow transfer n Hanau and Whip-Mix articulators. Saudi Dent J 1994; 6:8-12
- [5]Teteruck WR, and Lundeen HC. The accuracy of an ear face-bow. J Prosthet Dent 1966;16: 1039-46 [https://doi.org/10.1016/0022-3913\(66\)90169-7](https://doi.org/10.1016/0022-3913(66)90169-7)
- [6]Bailey JO. Evaluation of the third point of reference for mounting maxillary casts on the Hanau articulator. J Prosthet dent 1984; 51:199-201 [https://doi.org/10.1016/0022-3913\(84\)90260-9](https://doi.org/10.1016/0022-3913(84)90260-9)
- [7]Hindocha AD, Vartak VN, Bhandari AJ and Dudani MT. A cephalometric study to determine the plane of occlusion in completely edentulous patients. Indian Journal of Dental Research 2013;24;669-73 <https://dx.doi.org/10.1007%2Fs13191-011-0049-x>
- [8]Shah RJ, Sharma S and Agrawal P. Evaluation of two facebow transfer systems in measuring occlusal cant. Guident research; oct 2013 64-68
- [9]Nazir N, Sujesh M, Kumar R and Sreenivas P. Accuracy of two face-bow/semi-adjustable articulator systems in transferring the maxillary occlusal cant. Indian Journal of Dental Research 2012;23: 437- 42 <https://doi.org/10.4103/0970-9290.104945>.

[10]Seo KS, Park MH, Lee JH, Kim CH and Chae JH. The comparative study for occlusal plane between articulated cast model and cephalogram in orthognathic surgery patients. J. Kor. Oral Maxillofac. Surg. 2003;29:239-44

[11]Mittal R. Comparison of the occlusal plane in dentulous and edentulous patients: A cephalometric study. The Journal of Indian Prosthodontic Society 2008; 8: 195-200 <https://doi.org/10.4103/0972-4052.49182>