

Original Research Article

Maize response to leguminous biomass composted with phosphate rocks in the Northern zone of Tanzania

Abstract

A field study was conducted in Babati District of Northern Tanzania to evaluate maize response to leguminous biomass composted with phosphate rocks (PRs). Three leguminous (*Crotalaria juncea*, *Lablab purpureus* and *Mucuna pruriens*) strips were cultivated in 2013/14 to produce a biomass which was harvested at flowering to early podding stage and air dried. Air-dry biomass was composted with PRs from Minjingu (medium reactive PR) and Panda Hill (low reactive PR). Maize response to different treatments in a split plot design was evaluated across the field strips in 2014/15 season. The strips were used as main plots and each strip was divided into seven subplots receiving different treatments at random. A medium term maize variety SC. 627 was used as a test crop. Average maize grain yields obtained from *Crotalaria*, *Lablab* and *Mucuna* strips reached 5.3, 4.5 and 4.0 t ha⁻¹, respectively and were statistically different (P=0.05). Application of Minjingu or Panda Hill PR alone didn't increase maize grain yield above the control while Minjingu PR applied with urea or composted with biomass increased maize grain yield by 2.40 and 1.58 t ha⁻¹, respectively above the control. Application of Panda Hill PR with urea or composted with biomass increased grain yield by 1.20 and 1.06 t ha⁻¹, respectively above the control. The observed differences (0.82 and 0.14 t ha⁻¹) were not statistically significant indicating that biomass composted with PR was as effective as the PR applied with urea.

Key words: *Crotalaria*, *Lablab* *Mucuna*, phosphate rocks, compost, maize yield

26

27 The present address of Mawazo Shitindi is Sokoine University of Agriculture, Department
28 of Soil and Geological Sciences, P.O. Box 3008 Morogoro Tanzania.

29

30 **1. Introduction**

31 Maize (*Zea mays*) is Tanzania's most important staple food with an estimated annual
32 per capita consumption of 113 kg, contributing about 60% of dietary calories [1] and [2].
33 According to [3], the crop also contributes about 50% of Tanzania's rural cash income.
34 However, current production of maize in Tanzania is far below the national average yield
35 potential of 4.8 t ha⁻¹, fluctuating between 1.0 and 1.5 t ha⁻¹ [4]. Continuous maize production
36 without or with limited fertilizer application coupled with crop residue removal have been
37 reported as major factors for soil fertility decline and low crop yields [5; 6; 7]. Limited
38 fertilizer use in most developing countries has been attributed to their high costs and limited
39 availability [8; 9].

40 While food production per unit land is declining because of soil fertility deterioration,
41 the population of Tanzania has more than tripled from 12.3 million to 44.9 million between
42 1967 and 2012. Based on 2012 census projections, the population was expected to reach
43 47.42 million people by the year 2016 [10]. This increase in the population will cause
44 additional pressure on arable land because more than 70% of Tanzanians depend entirely on
45 agriculture for their food and income [10]. This calls for integrated soil fertility management
46 programs based on locally available resources so as to improve soil fertility and reduce
47 smallholders' dependence on imported industrial fertilizers.

48 Phosphate rock (PR) deposits located in Tanzania could serve as alternative source of
49 phosphorus (P) for smallholders but P contained in the rocks is not readily available for plant
50 uptake. Upon decomposition, plant biomass releases low- molecular-weight organic acids

51 that may complex calcium and other metals in the rock to free P for plant uptake [11]. Thus,
52 composting the rocks with leguminous biomass may improve the availability of nitrogen (N)
53 and P for plant uptake. The objective of the field experiment was to investigate carbon (C),
54 N, and P content of three common leguminous plants (*Crotalaria juncea*, *Lablab*
55 *purpureus* and *Mucuna pruriens*) used in Tanzania and the effect of each leguminous
56 biomass when composted with PRs on maize yield. The PRs used were those of Mijingu (a
57 PR of medium reactivity) and Panda Hill (a PR of low reactivity).

58

59 **2. Materials and Methods**

60 **2.1 Site Description, Soil Characterization and Fertility Assessment**


61 This study was conducted at Wang'waray Farmers Training Center (F.T.C) located in
62 Babati District of Manyara region in the Northern zone of Tanzania. The site is about 167 km
63 from Arusha and 4.5 km to the South East of Babati town along the road to Mamire Ward.
64 The center is at 1410 m above sea level on the foot hills of mount Kwaraa, and receives a
65 bimodal rainfall with average precipitation around 700-900 mm year⁻¹. However, as with
66 other areas in Tanzania, rainfall distribution at Wang'waray F.T.C and Babati District as a
67 whole has been altered by climate change to such an extent that the two seasons are now not
68 very distinct and average precipitation is less than 700 mm year⁻¹. Crop production is a major
69 land use activity at Wang'waray F.T.C. dominated by maize-legume intercropping and
70 rotation systems. Because soils at Wang'waray FTC were not characterized before, a profile
71 was opened and described according to FAO guidelines [12]. Representative profile and
72 surface (0-15 cm) soil samples were collected and shipped to the Soil and Geological
73 Sciences (SGS) laboratory at Sokoine University of Agriculture (SUA) in Morogoro for
74 physical and chemical analyses (Table1). Based on morphological description of the site, and
75 laboratory analyses performed on the profile samples, the soil was classified down to sub

76 group as *Rhodic Eutrosto*x using the USDA-NRCS Keys to soil taxonomy [13]. Analyses of
77 representative surface (1-15cm) soil samples collected from the rest of the field were used for
78 assessment of general fertility status of soils.

79 2.2 Leguminous Biomass Production

80 Following soil characterization, two portions of the field separated by a contour band were
81 ploughed and harrowed. On each portion of the field, three strips of 5 m x 76 m each were
82 established and randomly assigned to one of the three legume crops (two strips for each cover
83 crop) as shown in Figure 1. *Mucuna pruriens* and *Lablab purpureus* were planted at 50 cm x
84 30 cm spacing, while *Crotalaria juncea* was drilled at 50 cm inter row spacing. The first
85 weeding was done two weeks after germination and weeding was repeated whenever weeds
86 emerged to keep the competition for moisture and nutrients to a minimum.

87






Figure 1. Layout of the field for leguminous crop biomass production at Wang'waray F.T.C

2.3 Carbon, nitrogen and phosphorus contents of the biomass

At flowering - podding initiation stage, the biomass was cut close to the soil surface and air dried by species for later composting with Minjingu or Panda Hill PR. Before composting, the air-dry biomass was chopped into small pieces to increase surface area and thoroughly mixed. Subsamples were collected, oven dried at 55°C for 72 hours, and finely ground to < 0.5mm using a CT 193 Cyclotec™ Sample Mill [Foss Allé 1 Post box 260 DK-3400 Hillerød Denmark] for chemical analyses. Organic carbon (OC) was determined following the Walkely Black procedure [14], while total N was determined following Kjeldahl procedures [15]. For the determination of P and sulfur (S) in the biomass, a 0.5 g sample < 0.5 mm was digested following the HNO₃ - H₂O₂ wet digestion procedure using a 40 space Foss Tecator block digester. Phosphorus content of the digest was determined by a procedure using ascorbic acid method [16], while S content was determined by a turbidity method [17].

2.4 Phosphate Rock Collection, Processing, and Chemical Analysis

Minjingu PR was collected from Minjingu Mines and Fertilizers Company in Manyara region while Panda Hill PR was obtained from a storage facility at SUA. Both PRs were ground to pass a 100-mesh sieve at the Geological Survey of Tanzania (GST) laboratory in Dodoma region. A representative sample was collected from each PR and shipped to the

111 Southern and Eastern Africa Mineral Center (SEAMIC) laboratory in Dar es Salaam for X-
112 ray fluorescence (XRF) analysis.

113 **2.5 Production of Biomass-PR Composts**

114 Previously chopped leguminous biomass (< 2 cm) and ground PRs (< 100 mesh) were
115 composted by the pit method [18] with some modifications. In the modifications, the size of
116 an individual pit was 2 m x 2 m x 1m; floor and walls of each pit were lined with a
117 polyethylene plastic sheet to avoid leaching losses during decomposition. The biomass was
118 composted with a PR in alternating layers (i.e. PR was applied over every layer of biomass)
119 followed by 500g of dried cattle manure to inoculate the biomass. The biomass:PR ratio
120 varied from 12:1 to 18:1 based on the biomass size and N contents. Following inoculation,
121 water was applied to bring the moisture content of the compost mixture to about 60%.

122

123 Three PVC aeration pipes were inserted into each compost mix at regular intervals
124 and the material was covered with polyethylene plastic sheets to protect it from rain water
125 and undesirable/ foreign materials. The compost material in each pit was turned into a
126 different pit every 30 days for 120 days to allow optimum decomposition and water was
127 sprinkled at every turn to maintain the moisture at 60%. After the last turn, representative
128 samples were collected from each pit for laboratory analysis and all composts were air dried
129 to around 20% moisture content and stored for later use as source of N and P for maize.
130 Representative samples taken from each pit were shipped to the SUA-SGS laboratory for
131 chemical analysis. In the laboratory, representative compost samples were dried and ground
132 to pass through 0.5 mm for total N, P and SO₄-S analysis as previously described.

133

134 **2.6 Evaluation of Maize Response to Treatments**

135 The field strips previously used for cover crop biomass production were used in the
136 next season to evaluate maize response to newly imposed treatments. The experiment was
137 designed as a split plot arranged in a randomized complete block design (RCBD). The field
138 was divided into four blocks where half of each strip initially used to produce the crop
139 biomass was used as a main plot within a block and each main plot was divided into seven
140 sub plots (16 m²) which received randomly assigned treatments (Figure 2).

141

142

143

144

145

146

147

148

149

150

151

152

153


154

155

156

157

158



159

160

Figure 2. Layout of maize field experiment at Wang'waray FTC

161 Letters represent legumme species preceding maize on each strip (M = Mucuna, L = Lablab,

162 C = Crotalaria) while numbers (1 - 7) represent treatments imposed on experimental units.

163 Seven treatments were evaluated on each main plot. These include a common control where

164 maize was grown without external inputs after removal of the crop biomass (1), Minjingu PR

165 alone applied (2), Minjingu PR + urea (3), composted Minjingu PR + biomass (4). Panda Hill
166 PR alone (5), Panda Hill PR + urea (6), and composted Panda hill PR + biomass (7). Thus,
167 treatment combinations were identified as C1 to C7, L1 to L7, and M1 to M7 where C, L, and
168 M stand for Crotalaria, Lablab, and Mucuna strip, respectively.

169
170 The composts were applied at a rate corresponding to 112 kg N ha⁻¹ recommended for
171 maize in the Northern Zone [19]. The PRs were applied at 45kg P ha⁻¹ with or without urea
172 while urea was applied at 112kg N ha⁻¹ (split application at planting and two weeks following
173 germination) on selected plots based on treatment scheme. A medium **term** hybrid maize
174 variety (SC.627) was planted at 90 x 30 cm spacing (five rows per plot). At tasselling stage,
175 nine representative ear leaf samples were collected from each plot for nutrient analysis. At
176 maturity stage, maize ears of the three inner rows in each plot were harvested for yield
177 determination. Maize grain yield was reported at 13% moisture content, while maize stover
178 yield from the three inner rows of each plot was reported on oven-dry basis. The data
179 collected were subjected to analysis of variance (ANOVA) using a mixed procedure of SAS
180 software version 9.4 (SAS Instit. Inc. Cary, NC) and the means were separated at P = .05 by
181 Tuckey-Kramer procedure.

182

183 **3. Results and Discussion**

184 **3.1 Fertility status of soil at Wang'waray FTC**

185 Selected physical-chemical analyses of soil at Wang'waray FTC were as presented in
186 table 1. The soil had a medium pH value suitable for production of most crops with a very
187 low electrical conductivity indicating that there were no limitations for crop production due
188 to salt accumulation.

189

190 Table 1. Selected chemical properties of surface (0 -15 cm) soil samples at Wang'waray
191 F.T.C

Soil property	Mean [†]	Rating	Reference
pH – H ₂ O	6.88	Medium	[20]
EC (MScm ⁻¹)	0.05	Very low	[20]
Organic Carbon (g kg ⁻¹)	14.3	Low	[20]
Total N (g kg ⁻¹)	1.03	Low	[21]
Bray 1 P (mg kg ⁻¹)	5.54	Low	[20]
SO ₄ – S (mg kg ⁻¹)	9.38	High	[22]
Exch. Ca (Cmol kg ⁻¹)	7.40	High	[20]
Exch. Mg (Cmol kg ⁻¹)	2.96	High	[20]
Exch. K (Cmol kg ⁻¹)	3.28	High	[20]
Exch. Na (Cmol kg ⁻¹)	0.27	Low	[20]
PBS (%)	70.9	High	[20]
DTPA Extract. Cu (mg kg ⁻¹)	3.6	High	[20]
DTPA Extract. Zn (mg kg ⁻¹)	0.5	Low/medium	[20]
DTPA Extract. Mn (mg kg ⁻¹)	116.5	High	[20]
DTPA Extract. Fe (mg kg ⁻¹)	22.0	High	[21]
Sand (g kg ⁻¹)	643		
Silt (g kg ⁻¹)	87		
Clay (g kg ⁻¹)	270		
Textural class	Sandy Clay Loam		[23]

192

193 †, Each reading is an average of six representative surface soil samples

194

195 Levels of extractable S, exchangeable bases and DTPA extractable Fe, Cu and Mn were all
 196 **high but** the levels of organic carbon, total N, Bray-1 extractable P were low, and therefore
 197 limiting. The low levels of organic carbon, N, and P have been reported in highly weathered
 198 tropical soils like those of Babati [24].

199 3.2 Carbon, Nitrogen, and Phosphorus Content of Leguminous Biomass Used

200 Carbon contents of the leguminous biomass used varied significantly (P=.05) **while P**
 201 **contents were not statistically different (P = .05).**

202

203

204 Table 2. Chemical composition of the leguminous biomass used

205

Crop species	C	N	P	C:N	C:P	N:P
	-----%-----					
<i>Crotalaria juncea</i>	48.7 a	2.44 a	0.37 a	20.1	136	6.74
<i>Lablab purpureus</i>	41.6 b	2.30 a	0.34 a	18.1	124	6.76
<i>Mucuna pruriens</i>	44.5 b	2.00 b	0.36 a	22.3	122	5.75
LSD	6.30	0.14	0.05	-	-	-

206 †, Values in the same column followed by the same letter are similar (P = .05)

207

208

209 Chemical composition of plant species grown for compost production is an important factor

210 to take into account because it has effect on the rate at which plant material is acted upon by

211 decomposers to release nutrients in plant available forms. On average, OC contents of the

212 biomass used were 48.7%, 41.6% and 44.5% for *Crotalaria*, *Lablab*, and *Mucuna* biomass,

213 respectively. On the other hand, total N content of the biomass used were 2.4%, 2.3% and

214 2.0%, while the C:N ratios were 20.1, 18.1, and 22.3 for *Crotalaria*, *Lablab* and *Mucuna*

215 biomass, respectively. The total N values determined in all leguminous crop biomass were

216 below 3.0 % which is considered as critical value for sufficiency in most legume plants.

217 However, the tropical soil biology and fertility program data base cited by [25] specified total

218 N in the range of 1.6-5.7%, 1.7-6.3% and 1.4-6.5%, as normal for of *Crotalaria*, *Lablab* and

219 *Mucuna* biomass respectively when harvested at flowering stage depending on soil

220 properties and environmental condition of a given area. The data base also specified the C:N

221 ratios in the range of 8.0-32.1, 7.4-29.1, and 9.8-30.8 for *Crotalaria*, *Lablab* and *Mucuna*

222 biomass, respectively when harvested at flowering stage. Based on these specifications, the

223 OC, total N and C:N ratios were all within the normal range for the crop species used.

224 Furthermore, the C:N ratios of the biomass used were below 30:1 which is the recommended

225 highest value acceptable for an effective decomposition and mineralization of plant biomass
226 [26].

227


228 **3.4 Selected Chemical Properties of PRs Used**

229 Selected chemical properties of PRs are as presented in Figures 3, 4, and 5. Solubility
230 of PR depends largely on soil moisture status, soil pH, exchangeable Ca, available P and P
231 adsorption capacity of a soil [27]. Composition of PRs also affects relationships between
232 concentrations of their dissolution products and their sinks in the soil hence affecting
233 dissolution reactions in equilibrium. Apart from affecting the nature and rates of dissolution
234 reactions, chemical constituents of the PRs also play different roles in plant nutrition hence
235 contributing to variations in crop responses following application of PRs of different
236 chemical compositions [28].

237

238 Minjingu PR as shown in figure 3, has higher concentrations of P_2O_5 , CaO, MgO_2 ,
239 K_2O and NaO than Panda Hill PR. The differences are characteristic of geological origin i.e.
240 dependent upon parent material and dictate the relative availability of P, Ca, Mg, K and Na
241 from the two PRs. Apart from Na which is only essential in some plants where it has been
242 reported to take over the function of K when the latter is not readily available; P, Ca, K and S
243 are essential elements for all plants and therefore contribute to the fertilizer value of Minjingu
244 PR. Furthermore, with the exception of Ca, most of the elements found in higher
245 concentrations in Minjingu PR have low affinity for P. This explains the reason for higher
246 reactivity and therefore positive crop response reported following applications of Minjingu
247 PR than that of Panda Hill PR [29; 30; 31; 32].


248



249

250 Figure 3. Concentrations of P₂O₅, K₂O, MgO, NaO and SO₃ in Minjingu and Panda Hill PRs.
251

252 High concentration of Ca in Minjingu PR is also in agreement with the liming effects
253 reported following application of Minjingu PR on acid soils [20; 33]. Apart from creating a
254 more favorable environment for plant root growth, the liming effect of Minjingu PR on acid
255 soils can also correct imbalance of exchangeable cations in the soil system. A combination of
256 these effects explains the reason for higher crop response reported following application of
257 Minjingu PR than Panda Hill PR. Figure 4 indicates that Panda Hill PR has higher
258 concentrations of FeO₃, SiO₂, and AlO₃ than Minjingu PR. Higher concentrations of these
259 oxides are undesirable as far as reactivity of the PR is concerned because Fe, Si, and Al have
260 high affinity for P and therefore tend to form complex compounds with P, making it difficult
261 to be released from the PR for plant uptake. High concentrations of these metal oxides
262 explains the reason for low reactivity of Panda Hill PR as compared with Minjingu PR and
263 associated differences in crop response following applications of the two PRs on soils with
264 similar characteristics.



265

266 Figure 4. Concentrations of FeO₃, SiO₂ and AlO₃ in Minjingu and Panda Hill PRs.
 267

268 With the exception of MnO₃ content of Panda Hill PR, all oxides determined in the
 269 two PRs indicate low concentrations of micronutrients Zn and Cu for the two PRs to be
 270 considered as promising source of micronutrients (Figure 5). This implies that direct
 271 application of Minjingu or Panda Hill PR as source of P for crops will require an alternative
 272 source of micronutrient for a balanced fertilization. Co-application of the PRs with manure or
 273 composts may benefit plants more than just PR application alone or with industrial N
 274 fertilizers because animal manures and composts contain most nutrients though in small
 275 amounts [18].




Figure 5. Concentrations of ZnO₃, CuO₃, and MnO₃ in the PRs

276

277

278

279 3.6 Chemical composition of the PR-biomass Composts

280 **Organic carbon, total N** and P content of the composts produced are presented in

281 Table 3. Chemical analysis results indicate that OC content of the composts produced from

282 Mucuna biomass mixed with either Minjingu or Panda Hill PR was different (P=.05) from

283 OC determined in the composts of Crotalaria and Lablab biomass mixed with the same PRs.

284 Panda Hill PR composted with Mucuna biomass was found to have the highest and

285 significant (P=.05) total N concentration, followed by Minjingu PR composted with

286 Crotalaria biomass. Lablab composted with Panda Hill PR had the lowest N content of all

287 composted materials (P=.05).

288

289

290

291

292 Table 3. Selected chemical properties of composts used

293

Compost composition	OC	N	P	C:N	C:P	N:P
	----- % -----					
Minjingu PR + <i>Crotalaria juncea</i>	22.4 ab	1.98 b	0.51 a	11.3	42.7	4.02
Minjingu PR + <i>Lalab purpureus</i>	24.0 a	1.64 c	0.52 a	14.7	46.4	3.18
Minjingu PR + <i>Mucuna pruriens</i>	22.1 b	1.69 c	0.55 a	13.1	40.9	3.12
Panda Hill PR + <i>Crotalaria juncea</i>	23.3 a	1.70 c	0.49 a	14.1	45.4	3.80
Panda Hill PR + <i>Lablab purpureus</i>	23.2 a	1.36 d	0.48 a	16.6	49.0	2.96
PandaHill PR + <i>Mucuna pruriens</i>	21.8 b	2.16 a	0.38 b	10.8	58.0	5.37
LSD (P=0.05)	0.87	0.15	0.07	-	-	-

294

295 †, Values in the same column followed by different letter(s) are statistically different (p=0.05)

296

297 In general, the composted materials showed $\frac{3}{2}$, $\frac{3}{2}$, and $\frac{1}{2}$ lower contents of OC, total
 298 N and C:N ratio as compared with the initial biomass. A decrease in OC, N and C:N ratio as
 299 shown in Table 3 for the composts as compared with the initial biomass (Table 2) was caused
 300 by oxidation of OC to produce carbon dioxide that was lost as CO₂ gas while portion of the
 301 OC is incorporated into microbial cells. Lower total N content in the compost than previously
 302 determined in the biomass was probably caused by a dilution effect due to addition of PR to
 303 the compost material. Similar trend of total N decrease was reported when coffee pulp was
 304 composted with Minjingu PR using surface soil for inoculation of the compost mix [34].

305

306 Other research findings [35] reported a slight increase in total N of the compost
 307 relative to N content of the raw material when coffee pulp and coffee husks were mixed with
 308 cow dung and composted with phosphate rock after inoculation with P-solubilizing bacteria
 309 (*Bacillus megatherium*). However, the increase in N content reported [35] could be due to
 310 relatively high amount of cow dung (12 kg) equivalent to 20% of total weight of the compost
 mix used to enrich the compost.

3.8 Effect of Leguminous Crop Strips on Maize Grain Yield

Leguminous crop strips had a significant effect on maize grain yield only when Panda Hill PR was used as P source and the yields under Crotalaria strip was significantly greater than those under Mucuna strip. (Figure 6). Maize grain yield obtained from the three leguminous crop strips were 5.3, 4.5, and 4.0 t ha⁻¹ from Crotalaria, Lablab and Mucuna strips, respectively.




Figure 6. Effect of cover crop strips (species) on maize grain yield
†. Values for the same PR type followed by the same letter(s) are statistically similar (p=0.05)

In Bukoba District of Tanzania, maize grain yield of 0.7 t ha⁻¹ was reported following incorporation of crotalaria residues while lablab residues increased maize grain yield by 57-103% above the control crop yield although the effect of lablab was below yield increase obtained from crotalaria strips [36]. Other studies conducted in Tanzania reported maize grain yield ranging from 1.2 to 4.0 t ha⁻¹ following incorporation of crotalaria as green manure [37]. In South Africa, maize grain yields ranging from 2.6 to 10.6 t ha⁻¹ were reported following incorporation of Crotalaria, Lablab, and Mucuna [38]. Among all the leguminous

330 crops tested, maize biomass and grain yields were highest on Crotalaria plots [38]. Superior
331 influence of Crotalaria on maize grain yields over Lablab and Mucuna was also reported in
332 Malawi [38]. Maize grain yield obtained in this work is therefore within the range reported by
333 other researchers in Sab Saharan Africa (SSA); suggesting that legume biomass composted
334 with PRs could effectively substitute for the application of PRs with urea. Superior
335 performance of Crotalaria over Lablab and Mucuna also agrees with majority of research
336 works conducted in Tanzania and neighbor countries using these leguminous crops as source
337 of N for maize.


338

339 Other studies [39; 40] obtained results showing that incorporation of Lablab produced
340 more maize grain yield than Crotalaria and Mucuna. Variations reported in different studies
341 could be attributed to differences in soil property, local climatic conditions, yield potentials
342 of maize varieties used and management practices such as timing of biomass incorporation as
343 green manure vs. composting.

344 **3.9 Effect of treatments on maize grain yield**

345 Figure 7 presents maize grain yield obtained following application of different
346 treatments. Application of Minjingu or Panda Hill PR alone failed to increase maize grain
347 yield above the control. This observation is in agreement with findings reported by other
348 researchers [29 and 41] following direct application of Minjingu and Panda Hill PRs on soils
349 with varying properties. Such observations were attributed to application of PRs on soils
350 where P is not the primary limiting factor for crop performance, as well as masking effect of
351 moisture stress, soil acidity and deficiencies of other nutrients which affect maize yield [29,
352 41].

353



354

355 Figure 7. Maize grain yield obtained with different treatment combinations following
 356 leguminous crop biomass removal. † Values for the same PR type followed by the same
 357 letter(s) are statistically similar (P=.05)

358


359 Addition of urea with Minjingu PR and Minjingu PR composted with leguminous
 360 crop biomass increased maize grain yield by 2.40 and 1.58 t ha⁻¹, respectively above the
 361 control while addition of urea to Panda Hill PR and Panda Hill PR composted with
 362 leguminous crop biomass increased grain yield only by 1.20 and 1.06 t ha⁻¹, respectively
 363 above the control. Difference observed in maize grain yields following the application of
 364 Minjingu PR or Panda Hill PR alone were not significant (P=.05) even though the two PRs
 365 have different reactivity and chemical composition.

366 Average maize grain yield produced when legume biomass was removed but
 367 Minjingu or Panda Hill PR + urea was applied reached 5.62 t ha⁻¹ compared with 5.15t ha⁻¹
 368 when biomass-PR compost was applied. However, the observed difference (0.47 t ha⁻¹) was
 369 also not statistically significant (P=.05) indicating that biomass composted with PR was as
 370 effective as the PR applied with urea. This suggests that legume biomass composted with PRs
 371 could effectively substitute for the application of PRs with urea at Wang'waray FTC and
 372 other areas with similar soil type and climatic conditions in the long run.

373

374 3.10 Effect of Treatments on Maize Stover Yield

375 Figure 8 indicates that stover yield was significantly different ($P=.05$) between PR
376 alone and PR + urea treatments.



377

378

379 Figure 8. Effects of treatments on maize stover yield following leguminous crop biomass
380 removal. Values for the same PR type followed by the same letter(s) are statistically similar
381 ($P=.05$)


382

383 Application of Minjingu or Panda Hill PR with Urea produced the highest (2.76 t ha^{-1}
384 and 2.54 t ha^{-1}) yield of maize stover, respectively as compared with Minjingu or Panda Hill
385 PR alone (1.92 and 1.66 t ha^{-1}). However, maize stover yield obtained following application
386 of PRs with urea and PRs composted with cover crop biomass were not statistically different
387 ($P=.05$) from stover yields obtained in the control plots. The lowest stover yield obtained
388 following application of PR alone could be due to limited supply of N and further distortion
389 of the balance between nutrient supply levels in the soil. This observation is in agreement
390 with the lowest maize grain yield obtained when PRs were applied alone and highest grain

391 yield following application of PR with urea. As we seek for alternatives of synthetic
 392 fertilizers, PRs + biomass composting makes a good case, better still, if a reactive PR is used.


393 3.11 Interaction of legume crop strips x fertilizer treatments effect on maize grain yield

394 With the exception of *Crotalaria* strips, when above ground crop biomass was
 395 removed and no external input was applied, maize grain yield was below 4 t/ha. (Figures 9
 396 and 10).



397
 398
 399 Figure 9. Interactional effect of leguminous crop strips and treatments with Minjingu PR on
 400 maize grain. MPR = Minjingu PR; Values followed by the same letter(s) are similar (P=.05)
 401

402 Following removal of *Mucuna pruriens* and *Lablab purpureus* above ground biomass, the
 403 application of PRs without urea or compost did not increase maize grain yield compared with
 404 the control plot. Although not significant (P=.05), higher maize yield was generally obtained
 405 on crotalaria strips. Superior performance of maize on crotalaria strips implies that crotalaria
 406 has additional positive effects on rhizosphere processes. This makes another case for our
 407 study though additional research is required to confirm such processes.



408

409 Figure 10. Interactional effect of cover crop strips and treatments with Panda Hill PR on
 410 maize grain yield. PPR = Panda Hill PR; Values followed by the same letter(s) are
 411 statistically similar (P=.05)

412

413

414 4.0 Conclusion

415 This study investigated the effect of three leguminous crops (*Crotalaria juncea*,
 416 *Lablab purpureus* and *Mucuna pruriens*) biomass composted with Minjingu (medium
 417 reactivity) or Panda Hill (low reactivity) PR on maize yield. The effect of each PR composted
 418 with leguminous crop biomass on maize grain and stover yield was found to be similar to that
 419 of the PRs applied with urea, while PRs applied alone failed to increase maize yield above
 420 the controls. Similar maize yields obtained with PR-urea and PR-biomass compost treatments
 421 imply that leguminous crop biomass composted with PRs was as effective as PRs applied
 422 with urea in terms of P and N supply for maize. Based on these results, it was concluded that
 423 leguminous crop biomass composted with PRs have a potential for improving maize yield
 424 and could replace the use of urea for maize production in the long run. Cost-benefit analysis
 425 is however required to justify substituting urea for PR – biomass composts in maize
 426 production.

427 Reference

- 428 1. Saidia PS, Chilagane DA, Alexander W, Janet FM. Evaluation of EM technology on
 429 maize (*Zea mays* L.) growth, development and yield in morogoro Tanzania. 2010;
 430 Accessed 10 Aug. 2015. Available:
 431 kilimo.org/.../wp.../Saidia-Paul-Sabas-Chilagane-Daudi-Amos-et-al.pdf
 432

- 433 2. Mboya R, Tongoona, P, Derera J, Mudhara M, Langyintuo A. The dietary importance of
434 maize in Katumba ward, Rungwe district, Tanzania, and its contribution to household
435 food security. *African journal of Agricultural Research*. 2011; Vol. 6 (11) 2617 –
436 2626
- 437
- 438 3. USAID. The legal, regulatory, and institutional constraints to the growth of maize and
439 rice in Tanzania. Tanzania MicroCLIR Report. 2010; Accessed 25 September 2015.
440 Available: egateg.usaid.gov/sites/default/files/Tanzania_MicroCLIR.pdf
- 441 4. Rowhani P, Lobel DB, Linderman M, Ramankutty N. Climate variability and crop
442 production in Tanzania. *Agric.Forest Meteor*. 2011; 151:449-460
- 443 5. Bekunda MA, Bationo A, Sali H. Soil fertility management in Africa. A review of selected
444 research trials. In: Buresh RJ, Sanchez PA, Calhoun F. Eds. Replenishing soil fertility
445 in Africa. Soil Science Society of America Special Publication No. 51. Madison,
446 Wisconsin; 1997.
- 447
- 448 6. Sizlas C, Semoka JMR, Borggard, OK. Establishment of an agronomic database for
449 Minjingu phosphate rock and examples of its potential use. *Nutr Cycl Agroecosyst*.
450 2007; 78:225-237.
- 451
- 452 7. Kamhabwa F, Consumption of Fertilizers and Fertilizer Use by Crop in Tanzania.
453 International Fertilizer Development Center (IFDC); 2014.
- 454
- 455 8. Kpombrekou-A K, Tabatabai MA. Effect of low-molecular weight organic acids on
456 phosphorus release and phytoavailability of phosphorus in phosphate rocks when
457 added to soils. *Agric. Ecosyst. Environ*; 2003; (100):275-284.
- 458
- 459 9. Mowo JG, Janssen BH, Oenema O, German LA, Mrema, JP, Shemdoe RS. Soil fertility
460 evaluation and management by smallholder farmer communities in the Northern
461 Tanzania. *Agriculture, Ecosystems & Environment*. 2006; 116 (1-2): 47-59
462 <http://www.wsi.nrcs.usda.gov/products/w2q/awm/docs/neh637c2.pdf>.
- 463
- 464 10. National Bureau of Statistics (NBS); Office of Chief Government Statistician (OCGS)
465 Zanzibar. 2012 Population and Housing Census: Population Distribution by
466 Administrative Units; Key Findings. Dar es Salaam, Tanzania; 2013.
- 467
- 468 11. Kpombrekou K, Tabatabai MA. Effect of organic acids on release of phosphorus from
469 phosphate rocks. *Soil Sci*. 1994; (158) 442–453.

- 470
- 471 12. FAO. Guide line for soil description. FAO, Rome, Italy; 2006
- 472
- 473 13. Soil survey staff. Keys to Soil Taxonomy 12th Edition. USDA-NRSA; 2014
- 474 14. Nelson DW and Sommers L E. Total carbon, organic carbon and organic matter. In:
 475 Sparks DL, editor. Methods of Soil Analysis Part 3. Chemical Methods. SSSA Book
 476 series No.5. SSSA and ASA, Madson, Wisconsin; 1996.
- 477 15. Bremner JM. Nitrogen Total. In: DL Sparks, editor. Methods of Soil Analysis Part 3.
 478 Chemical Methods. SSSA Book series No.5. SSSA and ASA, Madson, Wisconsin;
 479 1996.
- 480 16. Kuo S. Phosphorus. In: Sparks DL, editor. Methods of Soil Analysis Part 3. Chemical
 481 Methods. SSSA Book series No.5. SSSA, ASA. Madson, Wisconsin; 1996.
- 482 17. Okalebo JR, Gathua KW, Woomer PL. Laboratory methods of soil and plant analysis.
 483 Working manual. UNESCO Press, Nairobi, Kenya; 1993.
- 484 18. FAO. On farm composting methods. Land and water discussion paper 2. Rome; 2003.
- 485 19. Nkonya E, Xavery P, Ankomay H, Mwangi F, Ponia, A, Moshi A. Maize production
 486 technologies in the Northern Zone Tanzania. Mimeo. Kansas University; 1998.
- 487 20. Landon J R, editor. Booker Tropical Soil Manual. A handbook for soil survey and
 488 agricultural land evaluation in the tropics and subtropics. Addison Wesley Longman
 489 Limited, England; 1991.
- 490 21. Dierolf T S, Fairhurst TH, Mutert EW. Soil Fertility Kit: A Toolkit for Acid Upland Soil
 491 Fertility Evaluation and Management in South East Asia. Potash and Phosphate
 492 Institute, Canada; 2001.
- 493 22. Reisenauer HM, Walsh LM, Hoefl RG. Testing soil for S, B, Mo, and Cl. In: Walsh, M.L.
 494 and Beaton JD, editors. Soil testing and plant analysis, SSSA Inc. Madson,
 495 Wisconsin; 1973.
- 496 23. Gee GW, Bauder J.W. Particle-size analysis. In: Klute A, editor. Methods of soil analysis:
 497 Part 1—physical and mineralogical methods. SSSA Book Series 5.1. SSSA, ASA.
 498 Madson, Wisconsin; 1986.
- 499
- 500 24. Maranguit D, Nguillaume T, Kuzyakov Y. Land use change affects phosphorus fractions
 501 in highly weathered tropical soils. *Catena*. 2017; 149: 385-393.
- 502
- 503 25. Sakala WD, Kumwenda JDT, Saka AR. The potential of green manure to increase soil
 504 fertility and maize yield in Malawi. *Biol Agr & Hort*. 2002; 21(2):121-130.
- 505
- 506 26. Graves ER, Hattemer G M. Composting: National Engineering Handbook. 2000.
 507 Accessed 30 June 2009. Available:

508

www.springerlink.com/index/11P842T5VW685N40.pdf

509

27. Van Straaten P. Rocks for crops: Agrominerals of sub-Saharan Africa. ICRAF, Nairobi, Kenya; 2002.

511

512

28. FAO *Use of phosphate rock for sustainable agriculture*. FAO Fertilizer and Plant Nutrition Bulletin No. 13. Rome; 2004.

513

514

515

29. Semoka JMR, Mnkeni PNS, Ringo HD. Effectiveness of Tanzania phosphate rocks of igneous and sedimentary origin as source of phosphorus for maize. *Zimb Jour Agr Res*. 1992; 30 (2):127 - 136.

516

517

518

30. Van Straaten P, Editor. Rock for Crops. Agro minerals of sub-saharan Africa. Fidelity National Information Solutions Canada, Scarborough, ON, M1B. 3C3, Canada; 2002.

519

520

31. Mnkeni PNS, Semoka JMR, Kaitaba EG. Effects of Mapogoro phillipsite on availability of phosphorus in phosphate rocks. *Trop. Agric*. 1994;71:249-253.

521

522

523

32. Szilas C, Semoka JMR, Borggaard OK. Can local Minjingu phosphate rock replace super phosphate on acid soils in Tanzania? 2006. Accessed 8 June 201. Available:

524

525

526

33. Szilas C. The Tanzanian Minjingu phosphate rock - possibilities and limitations for direct application. PhD Thesis. Royal Veterinary and Agricultural University, Copenhagen, Denmark; 2002.

527

528

529

530

34. Shitindi MJ. Response of tomato (*Lycopersicon esculentum M.*) to coffee pulp compost, Minjingu phosphate rock and coffee pulp - minjingu phosphate rock compost applied to a Chromic Acrisol. MSc. Thesis. Sokoine University of Agriculture. Morogoro, Tanzania; 2011.

531

532

533

534

35. Preethu DC, Bhanu Prakash BNUH, Srinivasamurthy CA, Vasanthi BG. Maturity indices as an index to evaluate the quality of compost of coffee waste blended with other organic wastes. Proceedings of the International Conference on Sustainable Solid Waste Management. 5-7 September 2007, Chennai India. 2007.

535

536

537

538

539

36. Baijukya FP, de Ridder N, Giller KE. Nitrogen release from decomposing residues of leguminous cover crops and their effect on maize yield on depleted soils of Bukoba District, Tanzania. *Plant and Soil*. 2006; 279: 77-93.

540

541

542

37. Lupatu M, Kilimwiko L. Natural fertilizers: New life for tired soils. *Africa Farmer* No.6; December 1991. Hunger Project Savaccon gallery, 240 E. 13th street New York; 1991.

543

544

38 Odhiambo JJO, Ogola JBO, Madzivhandila T. Effect of green manure legume – maize rotation on maize yield and weed infestation levels. *Afr. J. Agric. Res*. 2010; 5(8): 618-625.

545

546

547

- 548 39. Mureiythi JG Gachene CKK, Ojien J. The role of green manure legumes in smallholder
549 farming systems in Kenya. The legume research network project. *Tropical and*
550 *Subtropical Agro ecosystems*. 2003; 1:57-70.
- 551 40. Odhiambo JJO. Potential use of green manure legume cover crops in smallholder maize
552 production systems in Limpopo province, South Africa. *Afr Jour Agr Res*. 2011; Vol.
553 6(1): 107-112.