

**INFLUENCE OF DIFFERENT ‘PRATA-ANÃ’ BANANA BUNCH AGES ON
POST-HARVEST QUALITY**

ABSTRACT

Objective: To determine the ideal harvest season of ‘Prata-Anã’ banana bunches by means of physical and chemical analyses of fruit cultivation conditions in the northern state of Minas Gerais.

Study Design: The employed experimental design was the completely randomized design was used in a 5x5 factorial scheme, with five bunch ages and five assessment days.

Study Location and Duration: The experiment was run in an area with banana trees planted 20 months beforehand, located at Unimontes’s Experimental Farm, at 530m of altitude, with coordinates being -15°43’46.99” south latitude and -43°19’17.61” west longitude, between April and November 2017.

Methodology: The bananas bunches they were marked weekly from April 14 to May 12, and week days were standardized for each marking. Five bunch ages were defined – 16, 17, 18, 19 and 20 weeks after inflorescence emission – for harvest. For differentiation of emerged bunches, tapes of different color were used. When the bunches marked in the first week completed 20 weeks, all bunches were harvested, which happened on September 1. After harvested the fruits were subjected to storage in refrigerated chamber at 10°C ± 1°C and relative humidity of 90% +5% for 25 days. After being stored for 25 days, the bananas were taken out of the chamber and exposed to a room temperature of 25°C, which analyzes were performed for 9 days, with a two-day interval in between, simulating the marketing period. The following analyses were carried out: firmness, peel color, soluble solids, pH, titratable acidity, amide, total sugars, reducing sugars and electrolyte extravasation.

Results: Lower hue, chroma, soluble solids, titratable acidity, total sugar, reducing sugar and electrolyte extravasation values were found for bananas harvested at 16 weeks.

Conclusion: Bunch harvest age had a direct influence on post-harvest quality of bananas ‘Prata-Anã’. Fruits from 16-week bunches were superior in physical and chemical characteristics compared to other ages, meaning a longer post-harvest life.

Keywords: storage, *musa* ssp, maturation stage

1. INTRODUCTION

Banana trees (*Musa* spp.) are the most relevant fruitful trees worldwide and its production is mostly concentrated in tropical countries, being the second most produced fruit in Brazil [1]. According to [2], in 2016, Brazil was the third among countries with the highest banana production, behind India and China only; besides, this fruit appears among the three most produced tropical fruits, alongside orange and pineapple.

The southwest region is the second greatest banana producer, with the north of Minas Gerais being a major producing pole in Brazil, with a high social and economic importance for the region. Banana ‘Prata-Anã’ (AAB) and its different clones are the most prevalent in cultivation, with good market acceptance due to their excellent quality attributes, being considered elementary in nutrition.

For being climacteric fruits classified as perishable, bananas require techniques that slow down their rapid ripening, preventing post-harvest losses, especially while being transported to more distant consuming markets.

Harvesting fruits at proper maturation stages is determinant to maintaining post-harvest quality. Maturation point is the ideal harvest moment without the occurrence of damages, which provides fruits with a longer preservation period. This point is usually reached when the fruit becomes physiologically mature, which corresponds to its maximum size and weight, but does not have desirable characteristics for marketing and consumption. Later, the fruit continues to go through transformations, ripens naturally and becomes suitable for consumption. However, the ideal harvest point depends on correlations between physiological characteristics inherent of each variety, the ideal maturation stage, and post-harvest preservation technologies applied. In banana trees, fruits harvested prematurely may not be physiologically developed, which hinders their ripening process and final quality [3]. Nevertheless, harvesting overripe fruits leads to rapid quality loss, reducing their marketing period.

35 In addition to defining the best harvest stage, another way of reducing damages
36 and prolonging storage period is to keep fruits refrigerated. Refrigeration is considered
37 one of the most efficient methods for fruit preservation, maintaining the fruit's desirable
38 characteristics, similar to those of its early stage, due to a delayed maturation process.

39 A fruit's external characteristics – which relate to its appearance as well as size,
40 shape, color, lightness, absence of imperfections – and internal characteristics –
41 perceived in how it tastes, smells and feels – are the main attributes evaluated by
42 consumers, who demand for quality during purchase [3].

43 Therefore, determining the ideal harvest point of bunches is imperative, when
44 the fruit reaches its physiological maturation, that is, its maximum size and weight,
45 which will influence its post-harvest quality and resistance for a longer preservation
46 period.

47 The present study aimed to determine the ideal harvest season of banana 'Prata-
48 Anã' bunches by means of physical and chemical analyses on fruits maturing under
49 cultivation conditions in the north of Minas Gerais, allowing for their maximum
50 utilization in order to provide the final consumer with quality, in accordance with their
51 demands and preferences.

52 **2. MATERIAL AND METHODS**

53 **2.1 Fruit Material, Post-harvest Treatment and Environmental Conditions**

54 The experiment was run in an area with banana trees planted 20 months
55 beforehand, located at Unimontes's Experimental Farm, at 530m of altitude, with
56 coordinates being -15°43'46.99" south latitude and -43°19'17.61" west longitude. After
57 inflorescence emission, the banana trees were randomly selected and marked through
58 criterion proposed by [4] for sourcing of bananas at different bunch ages. They were
59 marked weekly from April 14 to May 12, and week days were standardized for each
60 marking. Five bunch ages were defined – 16, 17, 18, 19 and 20 weeks after
61 inflorescence emission – for harvest. For differentiation of emerged bunches, tapes of
62 different color were used. When the bunches marked in the first week completed 20
63 weeks, all bunches were harvested, which happened on September 1. After harvest, the
64 bunches were separated into bouquets with four fruits each and washed in water and
65 neutral detergent at 0.2% for latex coagulation and superficial cleaning. The bouquets
66 were then immersed in a solution of Imazalil fungicide, at a dose of 2mL.100mL⁻¹ of
67 water at room temperature, and dried outdoors. Each bouquet was stored in low-density

68 polyethylene packs measuring 25µm in thickness and put inside a standard cardboard
69 box for export. The fruits were subjected to storage in refrigerated chamber at 10°C ±
70 1°C and relative humidity of 90% ±5% for 25 days. After being stored for 25 days, the
71 bananas were taken out of the chamber and packs and exposed to a room temperature of
72 25°C, after which they were analyzed for 9 days, with a two-day interval in between,
73 simulating the marketing period.

74

75 **2.2 Physical Quality Attributes**

76

77 Peel and pulp firmness: determined by maximum penetration strength with a flat
78 tip measuring 4 mm in diameter, placed 10mm away from the fruit, with the aid of a
79 Brookfield digital penetrometer, model CT3 10 KG; measures were taken from the
80 medium area of the four fruits of the bouquet with and without peel, and results were
81 expressed as Newton (N).

82 Peel color: Reading was carried out on the four fruits of the bouquet, using the
83 Color Flex digital colorimeter, model CT3 10 KG, which expresses color using three
84 parameters: L*(lightness), which ranges from 0 (black) to 100 (white); a* (transition
85 from green (-a*) to red (+a*)) and b* (transition from blue (-b*) to yellow (+b)). Based
86 on L*, a* and b* values, hue angle (°h) and chroma saturation index (C*) were
87 calculated.

88 The angle Hue (° h) represents a new coloration of fruits, which varies from 0 to
89 360 ° where 0 ° represents red color, 90 ° yellow color, 180 ° green color, 270 ° color
90 blue and the 360 ° red color again.

91 – ° h * = actg (a * / b *) (-1) 90 for a * negative;

92 – ° h * = 90- (actg (a * / b *)) for a * positive

93 Chroma saturation index (C *) is a saturation risk of color pigments,
94 consequently reducing color intensity, varying from 0 to 60°, where 0 are close to gray
95 and 60° pure, calculated from the following formula:

96 – $C * = \sqrt{(a *)^2 + (b *)^2}$

97 **2.3 Determination of Chemical Quality Parameters**

98 Soluble solids: analysis performed by means of Reichert digital refractometer,
99 using the four banana pulp kneaded in food processor, with results expressed as °Brix.

100 pH: determined by electrometric method in potentiometer using, 10g of mashed
101 sample composed of four fruits diluted in 90mL of distilled water, in accordance with
102 [5].

103 Titratable acidity (TA): determined using analyte (10g of the four-fruit pulp
104 homogenized and diluted in 90mL), titrated with sodium hydroxide standard solution
105 (NaOH) at 0.1N, having phenolphthalein as indicator. Results were expressed as malic
106 acid percentage. All methodology used complies with [5].

107 **2.4 Physiological Parameters**

108 Amide: chemically extracted and spectrophotometrically determined according
109 to a chemical method by [6]. It was determined at 510nm and results were expressed as
110 percentage.

111 Total sugars (TS): extracted with ethyl alcohol and determined by the Antrona
112 method [7]. The sample was subjected to reading on spectrophotometer at 620nm, and
113 results were expressed as percentage.

114 Reducing sugars (RS): determined by Nelson's methodology [6]. Reducing
115 sugar content was calculated by spectrophotometry at 510nm, and results were
116 expressed as percentage.

117 Non-reducing sugars (NRS): obtained by differences between total sugars and
118 reducing sugars, as per the formula below:

$$119 \text{ Non-reducing sugars} = \text{Total sugars} - \text{Reducing sugars} \times 0.95.$$

120 Electrolyte extravasation: it was determined according to [8]; a peel disc was
121 removed per damaged area from each fruit of the bouquet, measuring 1cm in diameter,
122 with the aid of a metal punch. This section was washed in distilled water and
123 superficially dried on absorbent paper, then incubated for 2 hours in a capped test tube
124 containing 18mL of distilled water, under ambient conditions. After this period,
125 electrical conductivity was measured on a SCHOT conductivity meter, model CG 853.
126 Later, the tubes containing the peel samples were autoclaved at 121°C and 1.5atm for 30
127 minutes. After autoclaving, electrical conductivity was read again. Results were
128 expressed as the ratio between values obtained in the first and second measurements
129 multiplied by 100.

130

131

132

2.5 Experimental Design and Statistical Analyses

133

134

135

136

137

138

139

140

141

142

143

144

145

146

3. RESULTS AND DISCUSSION

147

148

149


150

151

152

The experimental design employed was the completely randomized type (CRD), in a 5x5 factorial scheme, with five bunch ages (16, 17, 18, 19 and 20 weeks after inflorescence emission) and 5 assessment periods (1, 3, 5, 7 and 9 days after storage). Four repeats were used, and the experimental unit was composed of four fruits. Data on the variables were subjected to tests for analysis of homogeneity of variance [9], residue normality by the Shapiro-Wilk test [10] and model non-additivity [11]. Results were then subjected to analysis of variance (ANOVA), considering as sources of variation bunch ages, assessment days after storage, and interaction between bunch ages and days after storage, tested at 5% probability. Interaction was sliced or not, depending on significance; regression analysis was conducted, and models were chosen based on significance, coefficient of determination and potential to explain the biological phenomenon. The variables were studied using statistical program SISVAR.

Analyzing peel firmness, significant interaction was observed between bunch age and assessment day, factors that simulate the marketing period of fruits; fruits from bunches younger than 19 weeks were firmer on the 9th day after storage, presenting values of 17.39N, 21.36N, 20.86N and 15.77N, for bunch ages corresponding to 16, 17, 18 and 19 weeks, respectively; bunches with 17 weeks were superior to the other treatments, with longer shelf life (Figure 1).



153

154


155 **Figure 1:** Peel firmness of banana ‘Prata-Anã’ harvested at different bunch ages against days after
 156 storage

157

158 On the other hand, 20-week bunches were less firm, reaching 6.15N on the last
 159 assessment day.[12], while working with banana ‘BRS Tropical’, found that fruits with
 160 greater development at the harvest point had reduced firmness. According to [13] and
 161 [14], firmness reduction is related to amide hydrolysis and solubilization of pectic
 162 substances, as well as to water loss.

163 As for pulp firmness, significant interaction was observed between bunch ages
 164 and storage days for this variable, which reduced as bunch age increased; on the first
 165 assessment day, for the ages of 16, 17, 18, 19 and 20 weeks, the values found were
 166 19.36N, 18.27N, 17.18N, 16.10N and 15.01N, respectively (Figure 2). It was possible to
 167 observe on the days after storage a sharp firmness reduction in fruits harvested with 18,
 168 19 and 20 weeks, which showed respective values of 2.83N, 1.74N and 0.70N,
 169 compared to those harvested with 16 and 17 weeks, which presented values of 5.00N
 170 and 3.92N, respectively; fruits from the bunch with 16 weeks were firmer than the
 171 others. The results of this experiment corroborate with study by [15], which related
 172 firmness loss of banana ‘Prata-Anã’ pulp to older bunch harvest age, stressing that high
 173 storage temperatures also contribute to firmness loss, and it is possible to observe that

174 not even temperatures under 10°C were enough to prevent softening in fruits from 20-
175 week bunches, that is, these fruits had a greater firmness loss.




176

177

178 **Figure 2:** Pulp firmness of bananas ‘Prata-Anã’ harvested at different bunch ages against days after
179 storage

180

181 For color-describing variables, the hue angle parameter defines the basic color of
182 samples and represents the average hue of the banana samples; results were significant
183 for interaction. The hue angle values found in the banana peels at different bunch ages
184 (16, 17, 18, 19 and 20 weeks) dropped from 106.2°, 104.7°, 103.2°, 101.7° and 100.2°
185 to 85.3°, 83.8°, 82.4°, 80.9° and 79.4°, respectively, with this drop occurring from the
186 1st to the 9th day after storage, which varied by treatment (Figure 3). This behavior is
187 expected because hue angle values close to 100° presented a greenish color, and as
188 values move further or closer to 80° the color of the fruit turns yellowish, evidencing
189 ripening. Fruits harvested with a bunch age of 16 weeks had hue angle values higher
190 compared to other ages on the last assessment day, indicating their preservation and
191 allowing for them to be marketed for a longer period. According to [16], fruit color is an
192 important parameter to track the ripening process, which, in the case of bananas,
193 corresponds to yellow, due to chlorophyll degradation and carotenoid synthesis, besides
194 being a criterion used to characterize maturation stages.




195

196 **Figure 3:** Hue angle (a) of bananas ‘Prata-Anã’ harvested at different bunch ages

197

198 Figure 4 displays chromaticity values, which express color intensity, that is,
 199 saturation in terms of pigment. Significant difference is observed between bunch
 200 ages; values stood at 37.64, 38.84, 40.04, 41.25 and 42.45 with 16, 17, 18, 19 and 20
 201 weeks of age, respectively. These results are higher than those found by [3] while
 202 analyzing chromaticity in peels of banana ‘Prata-Anã’ stored at 10°C, with 18, 19
 203 and 20 weeks of development, finding estimated mean values of 33.67, 33.87 and
 204 34.61, respectively.

205




206

207 **Figure 4:** Chromaticity of bananas 'Prata-Anã' harvested at different bunch ages

208

209 The soluble solids variable presented significant interaction, being influenced by
 210 harvest ages and days after harvest. Figure 5 shows the behavior of soluble solid mean
 211 values, and it is possible to observe an increase in soluble solid content as fruits ripen;
 212 from the 1st to the 9th assessment day for all treatments (16, 17, 18, 19 and 20 weeks of
 213 bunch age), the values found were 4.08-16.73, 5.33-17.98, 6.88-19.53, 8.73-21.38, and
 214 10.88-23.53° Brix, respectively. Bananas have a high amide content when green and, as
 215 they ripe, amide is hydrolyzed into simple sugars for it to be used in fruit respiration,
 216 thus raising soluble solid content during maturation. The results observed on the last
 217 assessment day in the present experiment are similar to those found by [17] while
 218 working with fruits of banana trees 'Maravilha' and 'Preciosa', which showed values
 219 between 18.85 and 23.31 after cultivation during the 1st and 2nd production cycles, in the
 220 Upper Medium São Francisco River.




221

222 **Figure 5:** Soluble solid content in bananas ‘Prata-Anã’ harvested at different bunch ages against days
 223 after storage

224

225 There was significant interaction for pH between bunch ages and assessment
 226 days after storage. Figure 6 shows the behavior of values obtained for banana ‘Prata-
 227 Anã’ pH throughout the assessment days in relation to bunch ages. For all treatments, it
 228 is possible to observe a rapid decline in values obtained 9 days after storage, from 6.04
 229 to 5.10, 6.07 to 5.14, 6.04 to 5.11, 5.94 to 5.01 and 5.78 to 4.84 for bunches with 16, 17,
 230 18, 19 and 20 weeks, respectively. According to [18], while working with banana tree
 231 fruits, found pH values between 5.28 and 5.60, close to those found in the present study.
 232 [19], while working with bananas ‘Prata-Anã’ stored for 14 days under different
 233 controlled atmosphere conditions, found a mean pulp pH value of 4.25 after 3 days after
 234 atmosphere removal and maintenance in ambient atmosphere. According to [20],
 235 working with modified atmosphere associated with refrigeration in bananas, pH values
 236 in mature fruits varying from 4.2 to 5.0 were observed. According to [16], during the
 237 maturation phase of fruits there is an accumulation of soluble sugars, precursors of
 238 organic acids, with predominance of malic acid, which leads to a pH reduction
 239 throughout ripening.




240

241 **Figure 6:** pH values in bananas ‘Prata-Anã’ harvested at different bunch ages against days after storage

242

243 Titratable acidity showed significant interaction and associated with increased
 244 malic acid concentration. The graph reveals an increase in malic acid content in all
 245 fruits as they ripen and the bunches becomes older. For the bunch ages of 16 and 20
 246 weeks, 0.17g and 0.26g of malic acid were found per 100g of pulp, respectively, on day
 247 1. On the 9th assessment day, which simulates the marketing period, it is possible to
 248 observe that the bunch age of 16 weeks presented a value of 0.40g, whereas the bunch
 249 age of 20 weeks presented a value of 0.49g of malic acid per 100g of pulp, superior to
 250 the other bunch ages, evidencing faster ripening and shorter shelf life (Figure 7). The
 251 mean titratable acidity values observed in the present study 9 days after storage were
 252 similar to those found by [21] while working with ‘Prata-Anã’ in two production cycles.
 253 These results are in line with [22] in that unripe bananas have low acidity, and during
 254 ripening this acidity slowly increases until reaching a maximum value when the fruit is
 255 ripe and later falls with its senescence. [23] obtained results superior to those found in
 256 this study, with 0.54g of malic acid per 100g of "Prata anã" banana pulp throughout the
 257 storage period. Organic acid decline has been attributed to respiration or sugar
 258 conversion that occurs when banana tree fruits are ripening. These acids provide a
 259 sugar-acid balance, which results in a more tasteful fruit when ripe [24].




260

261 **Figure 7:** Titratable acidity content in bananas ‘Prata-Anã’ harvested at different bunch ages against days
 262 after storage

263

264 As for amide content, there was significant interaction as it was influenced by
 265 harvest seasons and days after storage in cold chamber (Figure 8). From the values
 266 obtained, a slight drop was seen in the amide content of the fruits as bunch age
 267 increased that is, bunches harvested later, because, as fruits ripen, amide rapidly
 268 degrades to be converted into sugars and may vary depending on bunch harvest season.
 269 According to [25], banana is a fruit with high amide content when unripe, and as it
 270 ripens, amide is broken into sugars for it to be used in the fruit’s respiration, raising the
 271 content of soluble solids. [17], while studying physical and chemical characteristics of
 272 different banana tree fruits, observed differences between AAB group fruits; unripe
 273 bananas ‘Prata-Anã’ showed a value of 29.68%, close to that found in fruits with bunch
 274 age of 16 weeks and superior to the other ages.




275

276 **Figure 8:** Amide content in bananas 'Prata-Anã' at different bunch ages against days after storage

277


278 For total sugar, significant difference was found between bunch ages and
 279 assessment days. Total sugar percentages were 6.60%, 7.95%, 9.29%, 10.64% and
 280 11.98% for the ages of 16, 17, 18, 19 and 20 weeks, respectively (Figure 9). Bunch
 281 harvest age had a significant influence on amide and sugar content during storage.
 282 Concerning different assessment seasons, on days 1, 3, 5, 7 and 9 days after storage,
 283 there is a linear increase in sugar percentage, with values at 1.61%, 5.45%, 9.29%,
 284 13.14% and 16.98% (Figure 10). The results found in this study are lower than those
 285 observed by [26], who also reported an increase in total sugar values over the evaluation
 286 days, ranging from 4.05% to 25%. According to [17], while amide is hydrolyzed, there
 287 is an increase in total sugar content, which makes fruits ripe and sweet. The main sugars
 288 found in ripe banana pulp are glucose, fructose and sucrose.



289

290 **Figure 9:** Total sugar in bananas 'Prata-Anã' harvested at different bunch ages

291




292

293 **Figure 10:** Total sugar in bananas 'Prata-Anã' on days after storage

294

295 Significant results were observed in interaction for the reducing sugar variable.
 296 As of the first assessment, it is possible to observe a slight percentage increase in
 297 reducing sugar for all treatments; however, concerning fruits from bunches with 16
 298 weeks, they reached a lower value than the other treatments – 9.62%. Treatment with the
 299 ages of 17, 18, 19 and 20 weeks presented higher values – 10.13%; 10.63%; 11.15%
 300 and 11.65%, respectively –; age increase resulted in sugar increase, and lower sugar
 301 percentage in the fruits was found in 16-week bunches (Figure 11). [27], while working

302 with climatization of banana ‘Prata-Anã’, also found increases in reducing sugar content
303 during ripening, arguing that such increase was due to insoluble molecule
304 interconversion, such as non-reducing sugars into depolymerized sugars and then
305 soluble sugars.




306

307 **Figure 11:** Variation of reducing sugar content in bananas ‘Prata-Anã’ harvested at different bunch ages
308 against days after storage

309

310 For non-reducing sugar, significant difference was found between assessment
311 periods, that is, 1, 3, 5, 7 and 9 days after storage, with values being 1.02%, 2.27%,
312 3.52%, 4.77% and 6.02%, respectively (Figure 12). According to [28], while working
313 with 10 banana genotypes (Pacovan, PV 04-44, PV 03-76, ‘Prata-Anã’, ‘Fhia-18’,
314 ‘Pioneira’, ‘Prata-graúda’, ‘Caipira’, ‘Nanica’, ‘Thapmaeo’), found non-reducing sugar
315 values for banana ‘Prata-Anã’ of $1.3 \pm 0.21\%$, which are lower than those found in the
316 present study.

317




318

319 **Figure 12:** Variation of non-reducing sugar content on days after storage

320


321 The fruits' electrolyte extravasation percentage had significant difference
 322 between bunch ages and assessment days. There was an increase on the days after
 323 storage, reaching values close to 37% on the 9th assessment day (Figure 13). Thus, fruits
 324 at a more advanced maturation stage tend to lose membrane integrity and have a faster
 325 electrolyte extravasation compared to those at an earlier maturation stage [29].
 326 According to [30], working with "Prata-Anã" banana submitted to hydrothermal
 327 treatment, the same behavior was observed, increased extravasation of electrolytes
 328 throughout the days after fruits were removed from the cold chamber, regardless of
 329 immersion temperature. Significant difference was found between treatments; although
 330 variation between some treatments was small, fruits from 16-week bunches presented
 331 lower electrolyte extravasation percentage compared to the other treatments, with a
 332 value of 17.72% (Figure 14). As shown by the results of this experiment, the fruits'
 333 greater resistance is due to lower cell membrane degradation, estimated by electrolyte
 334 extravasation percentage.



335

336 **Figure 13:** Electrolyte extravasation of bananas ‘Prata-Anã’ against bunch age

337



338

339 **Figure 14:** Electrolyte extravasation of bananas ‘Prata-Anã’ against days after storage

340

341 **4. CONCLUSIONS**

342 **Bunch harvest age had direct influence on post-harvest quality of ‘Prata-Anã’**
 343 **banana. Fruits from 16-week bunches stored at 10°C ± 1°C and relative humidity of**
 344 **90% +5% for 25 days were superior for physical and chemical characteristics compared**
 345 **to other ages, leading to longer post-harvest life.**

346

347

348 **5. REFERENCES**

349

350 1. Severo, F. E. ; Matos, M. C. P. ; Clauzet, M. (2017) Banana production in the coastal
351 region of São Paulo: the discourse of sustainability versus the management of solid
352 residues of agrochemicals. *UnisantBioScience*, 5 (5), 395-407. Retrieved from:
353 <http://wiki.unisant.br/index.php/bio/article/view/783>

354 2. Faostat - Food and Agriculture Organization of the United Nations. *Crops* (2008).
355 Retract: <http://www.fao.org/faostat/en/#data/QC>. (2008, March 23).

356 3. Ribeiro, D. M (2006). Evolution of Physical and Chemical Properties during the
357 Ripening of Banana "Prata-Anã. (ThesisDoctorado, Federal University of Viçosa,
358 Minas Gerais, Brazil). Retrieved from: <http://locus.ufv.br/handle/123456789/781>

359 4. Fortescue, J.A., & Turner, D. W. (2005). The association between low temperatures
360 and anatomical changes in preanthetic ovules of *Musa* (Musaceae). *Scientia*
361 *horticulturae*, 104 (4), 433-444. <https://doi.org/10.1016/j.scienta.2005.01.005>Get rights
362 and content

363 5. Adolfo Lutz Institute (2008). *Physical Chemistry Method for Food Analysis*. 6ª Ed,
364 1st Digital Edition, São Paulo, p.1020. Retrieved from:
365 http://www.ial.sp.gov.br/resources/editorinplace/ial/2016_3_19/analisedealimentosial_2
366 008.pdf

367 6. Nelson, N. (1944). A photometric adaptation of the Somogyi method for the
368 determination of glucose. *J. biol. Chem*, 153(2), 375-380. Retirado de:
369 <https://pdfs.semanticscholar.org/131b/e395d4d24726d051472489140cd543aa839e.pdf>

370 7. Dische, Z. (1955). New color reactions for determination of sugars in polysaccharides.
371 *Methods of biochemical analysis*, 313-358.
372 <https://doi.org/10.1002/9780470110188.ch11>

373 8. Whilton, T. H.; Bassuk, N. L.; Ranney, T. G.; Reichert, D. L. (1992). An improved
374 method for using electrolyte leakage to assess membrane competence in plant tissues.
375 *Plant physiology*, 98(1), 198-205. <https://doi.org/10.1104/pp.98.1.198>

376 9. Burr, I. W., & Foster, L. A. (1972). A test for equality of variances. *West Lafayette:*
377 *University of Purdue*, 26.

- 378 10. Shapiro, S. S., & Wilk, M. B. (1965). An analysis of variance test for normality
379 (complete samples). *Biometrika*, 52(3/4), 591-611. DOI: 10.2307/2333709
380 <https://www.jstor.org/stable/2333709>
- 381 11. Tukey, J. W. (1949). One degree of freedom for non-additivity. *Biometrics*, 5(3),
382 232-242. DOI: 10.2307/3001938 <https://www.jstor.org/stable/3001938>
- 383 12. Castricini, A., Dias, M. S. C., & Rodrigues, M. G. V. (2016). Post-harvest of banana
384 'Tropical BRS' in the function of technical management. *Caderno de Ciências Agrárias*,
385 8 (3), 52-62. Retrieved from
386 <https://seer.ufmg.br/index.php/ccaufmg/article/view/3123/2802>
387
- 388 13. Vilas Boas, E. V. B. ; Rodrigues, L. J. ; De Paula, N. R. F. (2003). Physical,
389 chemical-chemical and chemical modifications of the banana 'Maçã' during maturation.
390 Latin American Symposium on Food Science and Technology, 5.
391
- 392 14. Silva, E. A. ; Boliani, A.C. ; Souza Corrêa, L.S. (2006). Evaluation of banana
393 cultivars (*Musa* sp) in the region of Selvíria-MS. *Revista Brasileira de Fruticultura*, 101-
394 103. <http://dx.doi.org/10.1590/S0100-29452006000100028>
- 395 15. Martins, R. N., Dias, M. S. C., Boas, V., de Barros, E. V., & Santos, L. O. (2007).
396 Banana-stored store 'Ananas' comes from bunches at 16, 18 and 20 weeks. *Science and*
397 *Agrotechnology*, 1423-1429. <http://dx.doi.org/10.1590/S1413-70542007000500023>
398
- 399 16. Birth Junior, B. B. do; Ozorio, L. P. ; Rezende, C. M., Soares, A. G. ; Fonseca, M. J.
400 de O. (2008). Differences between bananas of cultivars Prata and Nanicão along of
401 ripening: physical-chemical characteristics and volatile compounds. *Food Science and*
402 *Technology*, 28 (3) Retrieved from:
403 <http://www.redalyc.org/articulo.oa?id=39594008802>
- 404 17. Silva, M.J. R. da; Jesus, Fr. R. R. de; Angels, J. M. C. dos; Machado, M., Ribeiro,
405 V. G. (2016). Adrenal and post-harvest characterization of 'Maravilha'e' Preciosa'
406 bananas in the São Francisco Valley. *Ceres*, 63 (1) .[http://doc.doi.org/10.1590/0034-](http://doc.doi.org/10.1590/0034-737X201663010007)
407 [737X201663010007](http://doc.doi.org/10.1590/0034-737X201663010007)

- 408 18. Souza, M. E. de; Leonel, S. ; Martins, R. L. ; Segtowick, E. C. dos S. (2013).
409 Physical-chemical characterization and sensorial evaluation of banana fruits. *Native*, 13-
410 17.<http://dx.doi.org/10.14583/2318-7670.v01n01a03>
- 411 19. Siqueira, M. de S. B. 2014. Characterization of the quality of silver-banana stored in
412 different atmospheric controlled conditions. (Master's Dissertation,
413 Universidade Estadual do Norte Fluminense Darcy Ribeiro, Rio de Janeiro, Brazil).
414 <http://locus.ufv.br/handle/123456789/781>
- 415 20. Siqueira, C., Almeida de, H. J., Serpa, M. F. P., Batista, P. S. C., Mizobutsi, G. P.
416 Modified atmosphere associated with refrigeration in the conservation of banana fruits
417 resistant to Black Sigatoka. *Journal of Agronomy*, vol. 48, n. 4, p. 614-624, Oct-10,
418 2017. doi: 10.5935 / 1806-6690.20170071.
- 419 21. Silva, M.J. R. da; Angels, J. M. C. dos; Jesus, Fr. R. R. de; Santos, G. S. dos; Lima,
420 F. B. F. ; Ribeiro, V. G. (2015). Production and characterization of banana 'PrataAnã'
421 (AAB) in production cycles (Juazeiro, Bahia). *Ceres*, 60 (1).
422 <http://dx.doi.org/10.1590/S0034-737X2013000100017>
- 423 22. Aquino, C.F., Solomon, C.C., C.Cecon, P. R., Siqueira, D.L., Ribeiro, S., & Rocha,
424 M. (2017). Physical, chemical and morphological characteristics of banana cultivars
425 depending on maturation stages. *Revista Caatinga*, 30 (1), 87-96.<http://dx.doi.org/10.1590/1983-21252017v30n110rc>
- 427 23. Oliveira, E.S. ; Viana, F.M.P. ; Martins, M.V.V. Alternatives to synthetic fungicides
428 in the control of banana anthracnose. *Summa Phytopathologica*, v.42, n.4, p.340-350,
429 2016.
- 430 24. Sakyi-Dawson, E., Asamoah-Bonti, P., & Amponsah Annor, G. (2008). Biochemical
431 changes in new plantain and cooking banana hybrids at various stages of ripening.
432 *Journal of the Science of Food and Agriculture*, 88 (15), 2724-2729
433 <https://doi.org/10.1002/jsfa.3399>
- 434 25. Pimentel, R. D. A. ; Guimarães, F. N. ; Santos, V. D. ; Resende, J. D. (2010). Post-
435 harvest quality of banana genotypes PA42-44 and Prata-Anãcultivados in northern
436 Minas Gerais. *Journal of Fruit Research*, 32 (2), 407-412.
- 437 26. Santos, T. C., Aguiar, F. S., Rodrigues, M. L. M., Mizobutsi, G. P., & Pinheiro, J.
438 M. D. S. (2018). Quality of bananas harvested at different development stages and
439 subjected to cold storage. *Pesquisa Agropecuária Tropical*, 48(2), 90-97.

- 440 27. Prill, M.A. de S .; Neves, L. C .; Silva, S .; Grigio, M.L., Chagas, E. A .; Campos,
441 A.J. (2011). Banana-based 'Silver-Banana' banana: methods and times for greening after
442 refrigerated storage. Journal of On-line Mentoring, 5 (2), 134-142.
443 <http://dx.doi.org/10.18227/1982-8470ragro.v5i2.500>
- 444 28. Jesus, S. C. D., Matsuura, F. C. A. U., Cardoso, R. L., & Folegatti, M. I. D. S.
445 (2004). Physical and chemical characterization of fruits of different genotypes of banana
446 tree. Embrapa Meio Ambiente-Elaborated newspaper article (ALICE) .<http://dx.doi.org/10.1590/S0006-87052004000300001>
- 448 29. Nolasco, C.D.A., Solomon, C.C., C. Cecon, P. R., Bruckner, C.H., & Rocha, A.
449 (2008). Postharvest quality of 'Prata' banana as affected by hot water treatment. Science
450 and Agrotechnology, 32 (5), 1575-1581. [http://dx.doi.org/10.1590/S1413-](http://dx.doi.org/10.1590/S1413-70542008000500033)
451 [70542008000500033](http://dx.doi.org/10.1590/S1413-70542008000500033)
- 452 30. Pinheiro, J. M. da S., Mizobutsi, G. P., Aguiar, F. S., Rodrigues, M. L. M., de Jesus,
453 M. O., de Queiroz, L. G. C., & Alves, P. F. S. (2019). Physiological and Biochemical
454 Responses of Banana 'Prata-Anã' Subjected to Hydrothermal Treatment. Journal of
455 Experimental Agriculture International, 1-10.
456 <https://doi.org/10.9734/jeai/2019/v33i230140>