

1 ORIGINAL RESEARCH ARTICLE  
2 **COMPONENTS OF TREE BIOMASS IN AN INTEGRATED CROP-LIVESTOCK-FOREST**  
3 **SYSTEM**  
4

5 **ABSTRACT**

6 **Aims:** This study performed the adjustment of volumetric models, and determined the  
7 biomass of *Eucalyptus grandis* x *Eucalyptus urophylla* hybrid cultivated in a crop-livestock-  
8 forest integration system (CLF).

9 **Study design:** The experimental area consists of a crop-livestock-forest integration system  
10 where trees are hybrids clones of seven year old *Eucalyptus grandis* x *Eucalyptus urophylla*.

11 **Place and duration of study:** This work was carried out at Fazenda Santa Brígida, Ipameri,  
12 Goiás (Brazil).

13 **Methodology:** A forest inventory of the area was carried out in October 2015 when the tree  
14 component was fully developed. Diameter at breast height (DBH) (at 1.30 m) and total  
15 height (H) of trees were measured in the field and categorized according to 4 classes.  
16 Afterwards, 12 trees were felled, which were cubed and compartmentalized to determine the  
17 volume and biomass of their components. The volumetric models developed by Schumacher  
18 & Hall and Ogaya were applied to obtain determination coefficients.

19 **Results:** The average DBH was 18.28 cm and the average H was 23.47 m. The highest  
20 volumes of wood were observed in the diametric classes that presented the largest number  
21 of individuals, however in the class of higher DBH an average individual volume of 0.36 m<sup>3</sup>  
22 of wood was observed. The total biomass of *Eucalyptus* was 56.64 Mg ha<sup>-1</sup>, being 83.70%  
23 wood, 6.52% in branches, 6.37% in bark and 3.40% in leaves.

24 **Conclusion:** The volumetric models developed by Schumacher and Hall as well as Ogaya  
25 were found to be applicable for estimating the volume of wood in CLF systems, where both  
26 showed a determination coefficient of 0.866.

27 *Keywords: compartmentalization; Eucalyptus; CLFS; volume, modeling*

28

29 **1. INTRODUCTION**

30 Crop-livestock-forest (CLF) integration has been proposed as an economically  
31 viable production technology for the recovery and renovation of degraded areas in the  
32 Cerrados, a vast tropical savanna ecoregion of Brazil [1]. The main habitat types of the  
33 Cerrado include: forest savanna, wooded savanna, park savanna and gramineous-woody  
34 savanna. Savanna wetlands and gallery forests are also included.

35 In addition to the formation or recovery of pastures, this technique favors the  
36 production of grain cultivars along with the exploitation of tree biomass production, either  
37 simultaneously, sequentially, or rotationally [2]. The intensification of the production has  
38 several benefits to the producer and the environment, such as: improving the physical,  
39 chemical, and biological conditions of the soil, increasing the cycling and efficiency for the  
40 use of nutrients, reducing production costs of agriculture and livestock, opening new areas  
41 for production, as well as diversifying and stabilizing the income of the producer [3]. The  
42 included tree component biomass promotes benefits ranging from soil protection to  
43 availability of nutrients and organic matter in the soil by the deposition of leaves and tree  
44 branches [4,5].

45 *Eucalyptus* has been presented as a good option in the integrated CLF due to  
46 its rustic nature, rapid growth, great utilization, and economic value in the market, being an  
47 alternative for farmers interested in wood production [6]. The rapid growth of the *Eucalyptus*  
48 in Brazil can be explained by the large investment of the companies in order to achieve the  
49 demand of the silvicultural products. The high productivity of stands planted by the Brazilian  
50 forest companies is recognized worldwide, due to the higher average productivity the  
51 minimum time required until harvesting, continuous investment in research on genetic  
52 improvement and silvicultural. The preservation and maintenance of native forests to  
53 preserve the biodiversity, makes planted forests even more important for preservation,  
54 because they have high productivity in a short period of time, avoid exploration of natural  
55 forests. These plantations, besides providing different products, also help with carbon

56 sequestration, while also maintaining animal biodiversity [7,8].

57 In the integration of CLF, one of the challenges lies in the careful planning of  
58 the system in defining short, medium, and long-term actions. The competition for light  
59 between forest species and agricultural and pastoral crops requires special attention, as this  
60 directly influences the productivity of the system. However, this competition can be reduced  
61 by selecting genetic material, adapting the planting arrangement of the tree component, and  
62 silvicultural treatments, which, in addition to adding value to the wood, also allows for greater  
63 light entry into the integration system that contributes to the maintenance or increase in the  
64 productivity of the other components, as pasture and agricultural culture [9,2].

65 The configuration of tree component arrangements may influence plant height,  
66 diameter of breast height (DBH), and volume of wood. Clemente [10] verified that integrated  
67 systems with single and double row arrangements provided higher volumes of wood. In their  
68 study, Oliveira et al. [11] verified higher volumes of *Eucalyptus* wood in integrated systems  
69 with forages, than in monoculture.

70 The balanced relationship between the integrated CLF components is important  
71 for the expression of the productive potential of the species involved. In the case of tree  
72 species, especially the fast-growing ones such as *Eucalyptus*, accumulation and biomass  
73 production are influenced by age of trees, among other factors. In the juvenile phase,  
74 accumulation is higher in the canopy components, whereas a greater increase of biomass in  
75 the trunk component is perceived over time [12].

76 However, this work had the aim of adjusting volumetric models and determining  
77 the biomass of the *Eucalyptus grandis* x *Eucalyptus urophylla* hybrid cultivated in an  
78 integrated crop-livestock-forest (CLF) system in Ipameri / Goiás (Brazil).

79

## 80 2. MATERIALS AND METHODS

This study was carried out at Fazenda Santa Brígida in the municipality of Ipameri – Goiás (Brazil), located at 17° 39'22" south latitude and longitude west of 48°

12'22", and at an altitude of 800 m a.s.l. [11]. According to the classification of Köppen-Geiger [13], the climate of the region is Aw (tropical savannah with dry season in winter), and the average temperature of the region varies between 22 and 23°C. The mean annual rainfall is between 1200 and 1400 mm, having a wet period of seven months from October to April, and the remaining five months characterize the dry season.

According to Embrapa [14], the soil of the experimental area is classified as red latosol, being naturally acidic and with low base saturation with good drainage and sand-clay texture.

The experimental area consists of an integrated crop-livestock-forest system that tends towards the east-west direction. Tree planting was carried out with clones of the hybrid *Eucalyptus grandis* x *Eucalyptus urophylla* (GG 100) in 2008 in an area of approximately four hectares (ha). The trees were arranged in double rows (1 m x 1 m x 26 m), occupying 1.4 ha of the total area of the system [11].

81 Before establishment of the seedlings, soil acidity was corrected with the use of  
82 two tons per hectare of dolomitic limestone and one ton of gypsum. At planting, the base  
83 fertilization used was 400 kg ha<sup>-1</sup> of yoorin thermophosphate and 180 grams (g ha<sup>-1</sup>) of NPK  
84 formulation 06-30-30, supplemented with 0.4 g ha<sup>-1</sup> of zinc, 0.2 g ha<sup>-1</sup> copper, and 0.2 g ha<sup>-1</sup>  
85 boron. The half of this composition was incorporated into the bottom of the planting pit, and  
86 the remainder was distributed 20 days after planting in two lateral holes located 10 cm from  
87 the seedlings.

88 15 months after of planting, a pruning was performed, and at 30 months, the  
89 third cover fertilization was applied using NPK formulation 00 - 00 - 36 with the addition of  
90 0.2 g ha<sup>-1</sup> of copper and 0.6 g ha<sup>-1</sup> of boron per plant, provided in continuous fillet in the  
91 crown projection.

92

### 93 **2.1 Determination of *Eucalyptus* biomass**

94 The forest inventory of the area was carried out in October 2015 when the tree

95 component was fully developed, seven years after planting. DBH (diameter at breast height  
 96 at 1.30 m above soil level) and H (total height of trees) were measured in the field with the  
 97 aid of a caliper and the use of a clinometer. For the of DBH and H measurements, a  
 98 systematic sampling was carried out with regular intervals on every sixth tree line in which  
 99 measurements were made on the two individuals that composed it.

100 Based on the data obtained from the forest inventory, the trees were distributed  
 101 in four classes of diameters (Table 1). Subsequently, three individuals were felled for  
 102 sampling in each diameter class, considering the lower, middle, and upper limits, totaling 12  
 103 trees.

104 **Table 1.** Diametric distribution (cm) of *Eucalyptus* in the integrated CLF system.

Class interval	Class center	Number of sampled individuals
9  — 14	11.5	16
14.1 –  19	16.5	59
19.1 –  24	21.5	80
24.1 –  29	26.5	6

105  
 106 After the trees were felled, they were subjected to rigorous sampling, according  
 107 to the method developed by Smalian and described by Finger [15]. The height points for  
 108 taking diameters with and without bark were: 0.10 m, 0.30 m, 1.30 m, 2.30 m, and so on, at  
 109 one meter intervals up to full height.

110 After measuring the diameters, the trunk was sectioned into 1 m long logs to  
 111 the point where the diameter was seven centimeters (commercial diameter). From there, up  
 112 to a diameter of three centimeters was considered as tip of the trees, and the remaining  
 113 portions to the apex were considered branches. For the determination of the dry weight of  
 114 wood, the methodology developed by Schumacher [16] was used, in which three samples  
 115 were taken along the trunk. The total height of the tree was divided into three sections, and

116 the midpoint of each third of the tree was taken to compose the sample. Each sampling point  
117 was composed of the complete disc of the tree cylinder that had a thickness of ten  
118 centimeters.

119 After sectioning, the logs were weighed both with and without bark to determine  
120 the wet weight of the wood and bark. The tree canopy, in turn, was divided into two  
121 components: leaves and branches. These components were also weighed in the field and  
122 properly sampled to determine the dry weight in the laboratory, as well as to determine wood  
123 biomass (WB), branch biomass (BB), and leaf biomass (LB).

124 The biomass samples were sent to the Forest Ecology Laboratory (ECOFLOR)  
125 of the Federal University of Goiás. They were placed in a force air circulation oven at 65°C  
126 for drying until the weight of the samples remained stable to obtain the dry mass of the  
127 components with a precision digital scale (0.01 g).

128

## 129 **2.2 Data analysis**

130 In order to relate the DBHs and biomass components of each tree, linear  
131 regressions were performed for each component: wood, bark, branches, and leaves. For the  
132 volumetric models, the DBH and the total height of the tree were considered the  
133 independent variables, and the total volumes and the trunk with the bark were dependent  
134 variables. Four volumetric models, one single-entry and three double-entry, were chosen  
135 because they were the most used for the quantification of the production in forest stands and  
136 have not yet been tested in integrated CLF systems. The models tested are described in  
137 Table 2.

138

139

**Table 2.** Volumetric models tested.

Author	Type	Model
--------	------	-------

Husch	Single entry	$V = \beta_0 + \beta_1 DAP$
Ogaya	Double entry	$V = DAP^2 (\beta_0 + \beta_1 H)$
Schumacher & Hall (log)	Double entry	$V = \beta_0 + \beta_1 \ln(DAP) + \beta_2 \ln(H)$
Spurr (log)	Double entry	$V = \beta_0 + \beta_1 \ln(DAP^2 H)$

DBH= diameter at breast height; H = total height;  $\beta_0$  = value of the height estimated when the diameter is zero;  $\beta_1$  = slope of the line, which corresponds to the value of the first derivative;  $\beta_2$  = rate of change in volume ( $m^3$ ) as height (m) variation occurs with constant DBH (cm);  $\beta_3$  = coefficient of the multivariate model.

The volumetric models were adjusted and evaluated by means of adjustment and precision statistics, following the importance proposed by Draper and Smith [17]: graphical analysis of the residues; estimate of the standard error in percentage (Syx%) that indicates the proximity between the estimated values and those observed and the closer to zero the model and the determination coefficient (R<sup>2</sup>) that shows how much the dependent variables are explained by the independents and, in this case, the closer to a better model.

140

### 141 3. RESULTS AND DISCUSSION

142

#### 143 3.1 Determination of *Eucalyptus* biomass

144

The integrated crop-livestock-forest (CLF) system evaluated presents a density of 303 trees per hectare. This occupied 33.65% of the area designated to the system and an average production of 0.18  $m^3$  of wood per tree, totaling a volume of wood without bark of 54.80  $m^3$  ha in the studied system. The remaining 66.35% were destined to other economic activities within the integration, such as agricultural and forage production.

149

After performing the forest inventory, the diameter distribution was analyzed, and four diameter classes were obtained. It is notable that the height, density, and volume of wood without bark were higher in class III, being 30.33 m, 151 trees  $ha^{-1}$ , and 40.06  $m^3$   $ha^{-1}$ ,

151

152 respectively (Table 3). Through the dendrometric characteristics of this integrated CLF  
 153 system, it is possible to verify a trend in relation to the height behavior of the plants and their  
 154 DBHs, being that the DBH tends to increase as the height linear increases . This is contrary  
 155 to the expected behavior in more homogeneous forest stands where trees with higher  
 156 heights and smaller diameters are observed. Similar results can be found in another studies  
 157 with *Eucalyptus*, with different clones and management, such as the one conducted by  
 158 Miguel et al [18], in Niquelandia, GO and another one by Lemos-Junior et al [19] in CFL with  
 159 same species in Cachoeira Dourada, GO.

160  
 161 **Table 3.** Dendrometric characteristics of *Eucalyptus grandis* x *Eucalyptus urophylla* grown in  
 162 the integrated CLF system at Fazenda Santa Brígida in the municipality of Ipameri / Goiás /  
 163 2015.

Diameters Classes (cm)	Mean height (m)	mean DBH (cm)	Density(tree. ha <sup>-1</sup> )	Volume of wood without bark (m <sup>3</sup> .ha <sup>-1</sup> )
I (9-14)	12.58	10.45	30	1.25
II (14,1-19)	20.73	14.53	111	9.45
III(19,1-24)	30.33	21.57	151	40.06
IV(24,1-29)	30.22	26.55	11	4.04
Total	23.47	18.28	303	54.80


164  
 165 Generally, resource availability tends to be higher in stands with less density of  
 166 less trees, reflecting higher growth in broader plantations [20]. This fact can be observed in  
 167 this study because in spite of the densification of the trees in the planting lines, the spacing  
 168 between the *Eucalyptus* ridges provides greater light availability in this integrated CLF  
 169 system. This causes the effect observed in the height as demonstrated by Miguel et al [18]  
 170 in homogeneous plantation of *E. urograndis* in Niquelandia, GO, DBH and wood volume as  
 171 demonstrated by Lemos-Junior et al [19] in CFL with same specie in Cachoeira Dourada,



172 GO, that can be attributed more to the lesser effect of resource competition than to  
173 continuous plantings where the height and DBH ratio are inversely related. Different spacing  
174 and thinning regimes late in the life of the stand presented the highest values of basal area  
175 production. The choice of the best thinning regime for *Eucalyptus* clonal material will vary  
176 according to the plantation objective [21].

177 The maximum and minimum diameter found in this integrated CLF ranged from  
178 9.4 to 28.25 cm, and the highest tree density were located in classes II and III, which  
179 consequently contributed with a higher volume of wood within the CLF system (Figure 1). In  
180 its study with *Eucalyptus* clones GG100 (*E. grandis* x *E. urophylla*) of 4.5 years, Cerdeira  
181 [22] observed a diametric variation between 5.0 and 17.1 cm. Cerdeira [22] also reported  
182 that the classes of greater diameter were those that presented the greater number of  
183 individuals, a DBH variation close to that of the present study, but the central classes were  
184 those with the highest number of individuals. Thus, we highlight that CLF present higher  
185 production of trees with higher diameter class, being relevant for the production of wood with  
186 noble purpose and greater value added by planted individuals.

187 Although Class IV presents the highest average wood volume per tree of 0.36  
188 m<sup>3</sup>, its contribution to the system is around 10%, among the individuals of lower number  
189 present in this class. However, classes III and II were the ones that concentrated the largest  
190 number of individuals, being responsible for 86.4% of the wood produced in this area of the  
191 integrated CLF with an average volume of 0.26 and 0.08 m<sup>3</sup>, respectively, per individual  
192 within the classes.



193

194 Figure 1. Volume of wood ( $\text{m}^3 \text{ha}^{-1}$ ) in different diametric classes of *Eucalyptus* trees in an  
 195 integrated crop-livestock-forest system in Ipameri / Goiás.

196 When evaluating three 32-month-old *Eucalyptus* trees (*E. urophylla* x *E.*  
 197 *grandis*), Torres et al [6] found volumes of wood without bark ranging from 0.01 to 0.24  $\text{m}^3$   
 198 for DBHs between 6.79 and 20.8 cm. These results are similar to those verified in the  
 199 present work in the corresponding diametric classes.

200 The total biomass produced by the hybrid GG 100 in this integrated CLF was  
 201 56642.76  $\text{kg ha}^{-1}$ , and the trees belonging to class III contributed the most in this production.  
 202 In general, 38303.00  $\text{kg ha}^{-1}$  of biomass were quantified through individuals with DBH  
 203 between 19.1 and 24 cm 24.1 and 29 cm, contributing 67.62% of the total produced (Table  
 204 2.4).

205 Among the components analyzed in the *Eucalyptus*, the wood biomass  
 206 contributed the most to the total biomass of the integrated CLF trees with 83.70%, followed  
 207 by the branches with 6.52%. Considering the trunk biomass (wood + bark), this was 90.07%  
 208 and the contribution of the canopy (leaves + branches) was 9.92% (Table.4).

209


210 **Table 4.** *Eucalyptus* wood biomass (WB), bark biomass (KB), branch biomass (BB), and leaf  
211 biomass (LF) with seven years of integrated CLF cultivation in Ipameri / Goiás.

Class	Class I (9-14 cm)	Class II (14.1-19 cm)	Class III (19.1-24 cm)	Class IV (24.1-29 cm)	Total of components
LB (kg ha <sup>-1</sup> )	169.80	615.93	992.19	149.23	1927.15 (3.40)*
KB (kg ha <sup>-1</sup> )	393.60	1008.29	1974.81	318.41	3695.11 (6.52)
BB (kg ha <sup>-1</sup> )	139.31	859.96	2363.32	245.73	3608.32 (6.37)
WB (kg ha <sup>-1</sup> )	1540.81	9918.34	32972.68	2980.35	47412.18 (83.70)
Total Biomass	2243.52 (3.96)	12402.52 (21.90)	38303.00 (67.62)	3693.72 (6.52)	56642.76

212 \* Values in parentheses refer to the percentage of component contribution in relation to total  
213 biomass.

214 Evaluating biomass components in *Eucalyptus* stands with different ages, [16]  
215 verified a trunk biomass around 80.3% for plantations with 8 years, a result that is consistent  
216 with the present study.

217 In the present work, it was verified that the order of contribution of biomasses in  
218 the different components was Wood>Branches>Bark>Leaves (Figure 2). These results were  
219 similar to those verified by Giumarães et al [23] in Allegrete//RS in homogeneous plantation  
220 of *E. dunni* with four years of age and those reported by Benatti [24] in Campos das  
221 Vertentes/MG using *Eucalyptus* clones I-144 with 6.5 years of age. Regarding biomass, it  
222 should be pointed out that the 8-year-old crops present about 80% of the biomass in the tree  
223 trunks shows the potential of planting, at this age, already for biomass for energy production,  
224 since most of the biomass of the planting may be removed and used as fuel.



225

226 Figure 2. Contribution of *Eucalyptus* biomass from different components and diametric  
 227 classes in the integrated CLF system in the Ipameri / Goiás municipality.

228


229 Different results to the present work were observed by [25] when working with  
 230 *E. benthamii* in an CLF at 12 months of age, [26] with *Eucalyptus* sp. of three years of age  
 231 and [27] in *Eucalyptus* in the agroforestry system of 18 months, where they verified the  
 232 inversion in the production of leaves and bark. This explains the effect of biomass  
 233 distribution during the different tree development phases, as the first one focused on leaf  
 234 expansion and the second on the development of trunks and leaf area limitation [28,16].

235 In his study comparing different commercial clones of *Eucalyptus* in integrated  
 236 CLF systems in the municipality of Juara/MT, [9] verified that GG100 *Eucalyptus* was the one  
 237 that allocated the largest biomass in the trunk when planted in double lines corresponding to  
 238 62.6% of the total biomass of the trees at 15 months of age, Moreover, this behavior was  
 239 maintained at seven years, as verified in the present work with the same clone in which this  
 240 percentage reached 90.07% as predicted by [29].

241                    From a commercial and structural point of view, the objective of the cultivator is  
242 to increase the volume of the trunk and to improve the quality of the wood. Less biomass in  
243 the branches is desirable since the primary product is the wood for commercialized [9].

244                    In Figure 3, it can be verified through the regression analysis that the linear  
245 model was adequate to explain the increase of the biomass of the different components in  
246 relation to the DBHs. One can observe an intense relation between these, mainly for wood  
247 and bark, and with lower intensity with the branches.

UNDER PEER REVIEW



248

249 Figure 3. Production of the different components of biomass of *Eucalyptus grandis* x  
 250 *Eucalyptus urophylla* in relation to DBH in an integrated CLF system in the city  
 251 of Ipameri / Goiás. ( $p < 0.05$  for the  $R^2$  values)

252 The biomass gains of wood and bark due to the increase of the DBH were  
 253 homogeneous. In other words, as the DAP increased, the biomass of these components  
 254 also increased, which is justified by the high values of the coefficient of determination  
 255 presented in the respective regressions. While the biomass of leaves and branches did not  
 256 present significant increases with the increase of DBH, which evidences the accumulation of  
 257 biomass as a function of age. In more developed plantations, the biomass of the leaves and  
 258 branches decrease [16].

### 3.2 Adjustments of volumetric models

259 Table 5 shows the adjustments for the different models tested as a function of  
 260 height and DBH. Considering the graphical analysis of the residues, the standard error and  
 261 the determination coefficient, the double entry models of Schumacher & Hall (log) (0.866 and  
 262 21.33%) and Ogaya (0.866 and 20.78%) can be considered the most efficient to predict the  
 263 volume of wood for an integrated CLF system in this spatial arrangement (Figure 4).


264

265 **Table 5.** Adjustments of volumetric models attributed to the *Eucalyptus* plantation used in  
 266 the integrated crop-livestock-forest system and their estimated coefficients ( $\beta$ ),  
 267 coefficient of determination (R<sup>2</sup>), and standard error (Syx %).

Models	$\beta$ 0	$\beta$ 1	$\beta$ 2	R <sup>2</sup>	Syx(%)
Husch	-0.18775	0.031469	-	0.681	35.10
Ogaya	-0.27662	0.00045	0.019777	0.886	20.78
Schumacher & Hall (log)	-8.8478	0.617035	1.848882	0.886	21.33
Spurr (log)	-1.45173	0.207884	-	0.748	31.21

268

269



270

271 Figure 4. Waste residue distribution of the volumetric models tested for clones of *Eucalyptus*

272 *grandis* x *Eucalyptus urophylla* in integrated CLF system in the city of Ipameri / Goiás.

273



274                    However, all models tested had a satisfactory distribution of residues, an R2  
275 ranging from 0.681 to 0.866 and a standard error of less than 35%, which makes it possible  
276 to use these models to estimate the volume of wood in the integrated CLF system.

277                    Lemes Junior et al [19] consider the Näslund and Ogaya models as the most  
278 efficient to determine the volume of wood in the Integrated CLF system with *Eucalyptus* of  
279 six years of age in Cachoeira Dourada / Goiás. These presented coefficients of  
280 determination of 99.5 and 99.1%, respectively. However, in spite of verifying a higher  
281 coefficient of determination for the Schumacher & Hall model, Miguel [30] observed a  
282 standard error that was considered high, another criterion used to indicate the volumetric  
283 model was the graphical distribution of the residues. In this scenario, the Takata model was  
284 the most suitable for estimating the volume of wood in a settlement of seven years of *E.*  
285 *urophylla* in Niquelândia, north of Goiás.

                    In their study with a silvipastoril system in the region of Coronel  
Pacheco/MG, Müller et al [31] tested different volumetric models to estimate the volume of  
*Eucalyptus* trees, and they found that the Schumacher & Hall model presented the best fit  
for those conditions, as also verified in the present work. This demonstrated that the  
Schumacher & Hall model has also been used for the integrated CLF system, since its  
statistical properties almost always result in non-biased estimates.

286                    With regards to the tree component of the integrated CLF system evaluated, a  
287 forest inventory was carried out to verify that at seven years after planting, the total biomass  
288 produced by the *Eucalyptus grandis* x *Eucalyptus urophylla* hybrid presented 56.64 Mg ha<sup>-1</sup>,  
289 a mean tree height of 25 m, and a chest height of 18. 222 cm. This biomass presented a  
290 distribution with greater quantity in the wood component, followed by the branches, bark,  
291 and leaves.

292                    Although crop-livestock-forest integration presents limitations in its operation,  
293 this system becomes feasible from an adequate planning that meets the production  
294 demands of the property in the short, medium, and long term. Although it is a complex

295 system because of the need to optimize the production conditions of each component, it is  
296 necessary to know the ecophysiology of the plants that will make up the integration. Besides  
297 the aggregate environmental benefits, this is important to determine if the productivity of the  
298 system is satisfactory to meet the social and economic demands and, thus, achieve the  
299 precepts of sustainability.

300 The environmental and productive importance of the integrated CLF system can  
301 be considered for the need to deepen the knowledge of the behavior of each component of  
302 the integration and prompted the interest in carrying out this research. It can be concluded  
303 that, finally, the initial objectives were reached, and it is, therefore, time for these results to  
304 be released.

305 Another aspect to be considered is the need for continuation of this research,  
306 both for this region of the Cerrado of Goiás and for the other regions of the country. It is  
307 known that many agricultural systems, conducted in an inadequate way, have contributed to  
308 the degradation of environmental quality and, due to this condition, seek to maintain  
309 production through the opening of new arable areas. Knowledge of crop-livestock-forest  
310 integration, as well as studies on the various possibilities of system implementation, are  
311 important factors for the productivity of agroecosystem and reduction of negative impacts on  
312 the Cerrado and other biomes.

313

#### 314 **4. CONCLUSIONS**

315 The highest average volume of wood per tree was verified in the highest DBH  
316 class; The volumetric models of Schumacher & Hall and Ogaya were efficient to estimate the  
317 volume of wood in the integrated CLF system; The biomass of *Eucalyptus grandis* x  
318 *Eucalyptus urophylla* was 56.64 Mg ha<sup>-1</sup>, and 90.07% was present in the components of the  
319 trunk, while the others allocated in the canopy. Adequate cultural (debris and thinning)  
320 treatment throughout the crop cycle has negatively influenced the development of culture.

321

322 **REFERENCES**

323

324 1. VASCONCELOS VV, VASCONCELOS CV, VASCONCELOS DM. Phyto-Environmental  
325 Characterization of Brazilian Savanna (Cerrado) and Brazilian Atlantic Forest, with the  
326 Research of Stone Lines and Paleosols Geografia. Ensino & Pesquisa (UFSM), 2010; 14: 6-  
327 17. Portuguese.

328

329 2. VIANA MCM, ALVARENGA RC, MASCARENHAS MHT, MACEDO GAR, SILVA EA, SILVA  
330 KT, RIBEIRO PCO. Consorciação de Culturas com o Eucalipto no Sistema de Integração  
331 Lavoura-Pecuária Floresta. In: XXIX CONGRESSO NACIONAL DE MILHO E SORGO, 29,  
332 Águas de Lindóia. SP: Associação Brasileira de Milho e Sorgo (ABMS), 2012. Portuguese.

333

334 3. GONTIJO NETO MM, VIANA MCM, ALVARENGA RC, SANTOS EA, SIMÃO EP,  
335 CAMPANHA MM. Sistemas de integração Lavoura Pecuária Floresta em Minas Gerais.  
336 Boletim de Industria Animal, 2014; 71: 183-191. Portuguese

337

338 4. XAVIER DF, LÉDO FJS, PACIULLO DSC, PIRES MFÁ, BODDEY RM. Dinâmica da  
339 serapilheira em pastagens de braquiária em sistema silvipastoril e monocultura. Pesquisa  
340 Agropecuária Brasileira, 2011; 46: 1214-1219. Portuguese.

341

342 5. GATTO A, BARROS NF, NOVAIS RF, SILVA FR, LEITE FP, VILLANI EMA. Estoques de  
343 carbono no solo e na biomassa em plantações de eucalipto. Revista Brasileira de Ciências  
344 do Solo, 2010; 34: 1069-1079. Portuguese.

345

346 6. TORRES CMME, OLIVEIRA AC, PEREIRA BLC, JACOVINE LAG, OLIVEIRA NETO SN,  
347 CARNEIRO ACO. Estimativas da produção e propriedades da madeira de eucalipto em  
348 Sistemas Agroflorestais. Scientia Forestalis, 2016; 44: 137-148. Portuguese.

349

350 7. IBA. Relatório IBÁ. Brasília. Indústria Brasileira de Árvores 2016. 6-15 . Portuguese.

351

352 8. ORTIGOZA FDSS, SENNA RJ. Caracterização do segmento de florestas plantadas de  
353 eucalipto em Mato Grosso do Sul: Revista de Política Agrícola. Brasília, 2016; 74-89, 2016.  
354 Portuguese.

355

356 9. TONINI H, MORALES MM, MENEGUCI JLP, ANTONIO DBA, WRUCK FJ. Biomassa e  
357 área foliar de clones de eucalipto para a desrama. Nativa, 2016; 4: 271-276. Portuguese.

358

359 10. CLEMENTE MA. Características agronômicas do sorgo e eucalipto em diferentes  
360 arranjos espaciais. 51 f. Dissertation (Master Degree) – Federal University of Uberlandia,  
361 2015. Portuguese.

362

363 11. OLIVEIRA P, FREITAS RJ, KLUTHCOUSKI J, RIBEIRO AA, CORDEIRO LAM,  
364 TEIXEIRA LP, MELO RAC, SILVA A, VILELA L, BALBINO LC. Evolução de sistemas de  
365 integração lavoura-pecuária-floresta (ILPF): estudo de caso da fazenda Santa Brígida,  
366 Ipameri, Go. 2 ed. Planaltina: Embrapa Cerrados, 2015. 50 p. Portuguese.

367

368 12. SAIDELLES FLF, KÖNIG FG, SCHUMACHER MV. Avaliação da biomassa e dos  
369 nutrientes em espécies florestais de rápido crescimento. In: 1 Simpósio Brasileiro de Pós-  
370 Graduação em Engenharia Florestal. Anais...Santa Maria: Perspectivas e Tendência da  
371 Pesquisa Florestal, 2001; 1: 134-144. Portuguese.

372

373 13. CARDOSO MRD, MARCUZZO FFN, BARROS JR. Classificação Climática de Köppen-  
374 Geiger para o Estado de Goiás e o Distrito Federal. Acta Geográfica. 2014; 8: 44-55.  
375 Portuguese.

376

377 14. EMBRAPA. Sistema brasileiro de classificação de solos. 3. ed. Rio de Janeiro-RJ:  
378 Embrapa Solos, 2013. 306p. Portuguese.

379

380 15. FINGER CAG. Fundamentos de biometria florestal. Santa Maria: UFSM/ CEPEF/  
381 FATEC, 1992. 269 p. Portuguese.

382

383 16. SCHUMACHER MV, WITSCHORECK R, CALIL FN. Biomassa em povoamentos de  
384 *Eucalyptus spp.* de pequenas propriedades rurais em Vera Cruz, RS. Ciência Florestal  
385 (UFSM. Impresso), 2011; 21: 17-22. Portuguese.

386

387 17. DRAPER NR, SMITH H. Applied regression analysis. New York: John Wiley & Sons,  
388 407 p. 1966.

389

390 18. MIGUEL EP, LEAL FA, ONO HA, LEAL, UAS. Modelagem na predição do volume  
391 individual em plantio de *Eucalyptus urograndis*. Revista Brasileira de Biometria, 2014; 32(4):  
392 584-598. Portuguese

393

394 19. LEMES JUNIOR JM, SILVA NETO CM, SOUZA KR, GUIMARÃES LE, OLIVEIRA FD,  
395 GONÇALVES RA, MONTEIRO MM, LIMA NL, VENTUROLI F, CALIL FN. Volumetric models  
396 for *Eucalyptus grandis* x *urophylla* in a crop-livestock-forest integration (CLFI) system in the  
397 Brazilian cerrado. African Journal of Agricultural Research, 2016; 11(15): 1336-1343.

398

399 20. REINER DA, SILVEIRA ER, SZABO MS. O uso do eucalipto em diferentes  
400 espaçamentos como alternativa de renda e suprimento da pequena propriedade na região  
401 sudoeste do Paraná. Synergismus scyentifica. 2011; 6 (1). Portuguese.

402

- 403 21. FERRAZ FILHO AC, MOLA-YUDEGO B, GONZALEZ-OLABARRIA JR, SCOLFORO  
404 JRS. (2018). Thinning regimes and initial spacing for *Eucalyptus* plantations in Brazil. *Anais*  
405 *da Academia Brasileira de Ciências*, 2018; 90(1): 255-265. Portuguese.
- 406
- 407 22. CERDEIRA ALN. Modelos para quantificação do volume de diferentes sortimentos em  
408 plantio de *Eucalyptus urophylla* x *Eucalyptus grandis*. 66 f. Monography (Forest  
409 Engineering), University of Brasília, 2012. Portuguese.
- 410
- 411 23. GUIMARAES CC, SCHUMACHER MV, WITSHORECK R, SOUZA HP, SANTOS JC.  
412 Biomassa e nutriente em povoamento de *Eucalyptus dunnii* Maiden no pampa gaúcho.  
413 *Revista Árvore*. 2015; 39 (5): 873-882. Portuguese.
- 414
- 415 24. BENATTI BP. Compartimentalização de biomassa e nutrientes em estruturas de plantas  
416 de eucalipto cultivadas em solos distintos. 114 f. Dissertation (Master Degree) – Federal  
417 University of Lavras. 2013; Portuguese.
- 418
- 419 25. UTIMA, A. Y. Crescimento e rendimento dos componentes agrícola e arbóreo de um  
420 sistema integrado de produção agropecuária no ano de implantação em área de proteção  
421 ambiental. 78 f. Dissertation (Master Degree in Agronomy), Federal University of Paraná,  
422 2015. Portuguese.
- 423
- 424 26. CARON BO, ELOY E, SOUZA VQ, SCHMIAT D, BALBINOT R, BEHILNG A,  
425 MONTEIRO GC. Quantificação da Biomassa florestal em plantios de curta rotação com  
426 diferentes espaçamentos. *Comunicata Scientiae*. 2015; 6 (1): 106-112. Portuguese.
- 427
- 428 27. CALIL FN, VIERA M, SCHUMACHER MV, LOPES VG, WITSCHORECK R. Biomassa e  
429 nutrientes em Sistema agrossilvicultural no extremo sul do Brasil. *Ecologia e Nutrição*

430 Florestal. 2013; 1 (2): 80-88. Portuguese.

431

432 28. GONÇALVES JLM, STAPE JL, BENEDETTI V, FESSEL VAG, GAVA JL. Reflexos do  
433 cultivo mínimo e intensivo do solo em sua fertilidade e na nutrição das árvores. In:  
434 GONÇALVES, J. L. M.; BENEDETTI, V. (eds.) Nutrição e fertilização florestal. Piracicaba:  
435 IPEF, 2005. 1-57. Portuguese.

436

437 29. MUÑOZ F, RUBILAR R, ESPINOSA M, CANCINO J, HERRERA M. The effect of pruning  
438 and thinning on above ground aerial biomass off *Eucalyptus nitens* (Deane & Maiden)  
439 Maiden. Forest Ecology and Management. 2008; 255 (3): 365-373.

440

441 30. MIGUEL, E. P. Avaliação biométrica e prognose da produção de *Eucalyptus urophylla*  
442 (s.t. blake) na região norte do estado de Goiás. 166 f. Dissertation (Mester Degree in Forest  
443 Engeneering), Federal University of Paraná. 2009. Portuguese.

444

445 31. MÜLLER MD, FERNANDES EM, CASTRO CRT, PACIULLO DSC, ALVES FF. Estimativa  
446 de acúmulo de biomassa e carbono em sistema agrossilvipastoril na zona da mata mineira.  
447 Pesquisa Florestal Brasileira, 2009; 60: 11-17. Colombian.

448

449

450

451

452

453

454

455