

## Effect of salinity on germination of lettuce cultivars produced in Brazil

## ABSTRACT

The challenge of recovering degraded soils due to salinity excess leads to the search for more effective strategies that can overcome this problem. Among these, one of the approaches is the use of resistant plant varieties in affected areas. This study aimed to evaluate the influence of different doses of salts on seed germination and seedling formation of two lettuce cultivars (Hanson and H121) and to verify the existence of tolerance among the cultivars. A completely randomized design was used in a 2x5x4 factorial scheme, where the cultivars were evaluated under five distinct salt doses (0, 25, 50, 75 and 100 mol.m<sup>-3</sup>, conductivities of 0.0, 2.8, 5.4, 8.0 and 10.6 dS.m<sup>-1</sup>, respectively), with four replicates. In this sense, the following variables were evaluated: germination rate (GR), germination speed index (GSI), seedling height (SH), root length (RL) and percentage of dry matter in relation to fresh matter (DM%). As a result, the Hanson cultivar presented better performance than the H121, under all the different salt doses, in all the studied variables. Also, the EC of 2.8 dS.m<sup>-1</sup> did not affect any of the studied variables, including both cultivars. However, EC above 2.8 dS.m<sup>-1</sup> had, significantly, reduced the development of the cultivars. The Hanson cultivar was influenced only in the variables SH, RL and DM%, where and DM% were influenced by EC values above 8.0 and 10.6 dS.m<sup>-1</sup>, respectively. The cultivar H121 was significantly influenced by all evaluated variables, which demonstrates its greater susceptibility to salinity.

*Keywords: Electrical conductivity, osmotic potential, water consumption, tolerance.*

## 1. INTRODUCTION

Population growth, food security and the scarcity of nonrenewable natural resources are some of the challenging themes that are present in the main discussions about the transformations in contemporary agriculture [1,2]. In a more rational model, priority is given to respect for the environment, where there is a growing concern with the management of natural resources, and at the same time, with the economic viability of the agricultural sector for its professionals, in order to guarantee the sustainability of current and future generations [3,4].

The development of sustainable agriculture is intrinsically related to the possibility of adopting measures that overcome the main challenges of agricultural production, in a rational and environmentally friendly way [5]. Many are the adversities found in agricultural fields throughout the world, but in the past few years, one has been gaining prominence, the soil salinity [6]. Soil salinity can occur naturally due to the characteristics of the source material [7]. However, the most damaging salinity is that induced by modern agricultural systems, where the irrational use of fertilizers and the use of brackish water for irrigation are the main agents that cause this adversity [8,9].

Soil salinity can make it unproductive, indirectly reducing the income of the farmers who cultivate the soil [10]. In Brazil, small farmers are the class that suffers the most from soil salinization, since the low income of these producers does not allow them to adopt

technologies to reverse it. Thus, technologies that allow cultivation on lands with an excess of salts are indispensable for the success of these farmers in the field. Among the available technologies, there is the selection of cultivars that tolerate high salinity concentrations.

Among the crops produced by small farmers, which suffer from the elevation of salinity levels, there is the lettuce (*Lactuca sativa* L.) [12]. The lettuce stands out because of its low cost of production, market acceptability, and its cultivation requires small areas, which makes it suitable for small farmers [12,13]. Lettuce cultivation is strongly affected by salinity, and about 13% of the yield declines by each unit of electrical conductivity (EC) increased above 1.3 dS m<sup>-1</sup> [13]. However, there is variation among cultivars, where some differ from the others regarding their tolerance to high levels of salinity [14,13].

The tolerance to salinity is a polygenic characteristic, which makes it difficult to identify tolerant genotypes, due to a large number of genes involved in the trait's control [15,16]. Identifying tolerant materials among the cultivars on the market has become an important strategy for expansion of lettuce crops in places where salinity is a limiting factor. Since it reduces the time and cost to obtain superior lineages in breeding programs. Given the above, this study aimed to evaluate the influence of different salt concentrations on germination of two lettuce cultivars, as well as to verify the existence of tolerance to salinity in the cultivars.

## 2. MATERIAL AND METHODS

The study was conducted in a completely randomized design in a 2x5x4 factorial scheme. The Hanson (cultivar 1) and H121 (cultivar 2) cultivars were used. To evaluate the tolerance to salinity, five distinct concentrations of sodium chloride (NaCl) were tested: 0, 25, 50, 75 and 100 mol.m<sup>-3</sup>. All concentrations of NaCl were diluted in distilled water, whose initial EC was zero. These concentrations corresponded to solutions with the following electrical conductivities: 0.0, 2.8, 5.4, 8.0 and 10.6 dS.m<sup>-1</sup>, respectively, measured with a conductivity / TDS and salinity meter.

To evaluate the germination, four replicates of 25 seeds were made in each treatment, wherein previous tests all the seed lots used in the study presented a germination rate (GR) higher than 90%. The test was performed in a 10 x 1.5 cm petri dish with the use of greatest paper. The paper was accommodated in double sheets at the bottom of the petri dish, and the seeds were evenly distributed over it. The saline solution was added after the seed distribution at 2.5 times the dry paper weight.

In order to favour germination, the seeds were kept in a Biochemical Oxygen Demand incubator (BOD) for 7 days, allowing total control of lighting and temperature that was alternated to maintain 12 hours of light and 12 hours of dark at a temperature of 25 ° C.

The germination evaluation occurred on the seventh day with the seedling count as proposed by the authors [17]. Also, only seeds with a primary root greater than 2 mm, as established by the authors [18], were considered as germinated. Through the counting data, it was possible to determine the GR in each treatment. Along with the germination test, daily germinated seedlings were counted from the beginning of the sowing until the seventh day, when the germination was stabilized. The germination speed index (GSI) was obtained using the equation (1) proposed by the author [19].

$$GSI = \sum (n/t)(1)$$

Where:

80 n = number of normal seedlings computed on the day of the count;

81 t = number of days from sowing until the day of the count.

82 After the germination stabilization on the seventh day, the variables seedling height (SH),  
83 root length (RL) and percentage of dry matter in relation to fresh matter (DM%) were  
84 determined. To do so, ten normal seedlings of each replicate were selected. The  
85 determination of the SH and RL was performed using a graduated ruler (centimetres). SH  
86 was measured from the neck to the apical meristem, while the RL was measured from the  
87 neck to the root cap.

88 The DM% was obtained using the equation (2):

89  $DM\% = mf/mi$  (2)

90 Where:

91 DM%: percentage of dry matter in relation to fresh matter;

92 mf: Total dry mass of the seedling;

93 mi: Total fresh mass of the seedling.

94 The seedlings had their fresh and dried masses measured by a precision scale. In order to  
95 obtain the dry mass values, the seedlings were left in an oven at 75° C until they reach a  
96 constant weight. Lastly, the data were submitted to the Kolmogorov Smirnov normality test,  
97 and then the variance analysis was performed. The means between treatments within the  
98 same cultivar were compared by the Tukey test, and the means of treatments between the  
99 cultivars were compared by the t-test. A simple linear regression analysis was used to infer  
100 the behaviour of the different cultivars under the effect of different salt concentrations in the  
101 studied variables. All statistical analyses were performed at a 5% probability level.

### 102 3. RESULTS AND DISCUSSION

103 The results presented in table 1 demonstrate that cultivar 1 obtained superior performance in  
104 relation to cultivar 2, for all the different tested. Cultivar 2 was significantly influenced (p-  
105 value <0.005) by the EC in all evaluated variables, while for cultivar 1, only SH, RL and  
106 DM% variables were influenced.

107 The EC of 2.8 dS.m<sup>-1</sup> did not affect any of the studied variables in both cultivars (Table 1).  
108 Similar results were found by the author [14] when studying salinity levels in the germination  
109 of the Elba cultivar. Conversely, higher than 2.8 dS.m<sup>-1</sup> negatively affected the root growth of  
110 both cultivars. The authors [20] emphasizes the importance of root formation in this  
111 phenological phase of the vegetables. The low root development in this phase culminates in  
112 fragile seedlings that tend to suffer more from environmental and biological stresses.

113 **Table 1. A comparative test of averages for germination rate (GR), germination speed**  
114 **index (GSI), seedling height (SH), root length (RL) and percentage of dry matter in**  
115 **relation to fresh matter (DM%), for the cultivars 1 and 2 submitted to different**  
116 **electrical conductivities.**

EC (dS.m <sup>-1</sup> )					
0	2.8	5.4	8	10.6	

Cultivar		GR (%)			
Cultivar 1	99.67±0.33	99.67±0.33 Aa	99.0±1 Aa	98.0±1.15 Aa	98.0±1.15 Aa
	Aa				
Cultivar 2	91.0±1 Ab	89.0±4.43 Aa	74.0±6.63 ABb	58.0±4.76 Bb	37.0±1.91 Cb
GSI					
Cultivar 1	39.49±0.24	39.34±0.34 Aa	39.54±0.17 Aa	38.41±0.51 Aa	38.0±0.74 Aa
	Aa				
Cultivar 2	24.34±0.94	23.54±1.62	18.29±1.94	13.49±0.71	8.3±0.39 Db
	Ab	ABb	BCb	CDb	
SH (cm)					
Cultivar 1	2.97±0.08	3.18±0.04 Aa	2.91±0.06 ABa	2.63±0.1 Ba	2.02±0.12 Ca
	Aba				
Cultivar 2	1.62±0.07 Ab	1.46±0.05 Ab	1.42±0.05 Ab	1.06±0.03 Bb	0.79±0.06 Cb
RL (cm)					
Cultivar 1	1.64±0.15 Aa	1.40±0.08 ABa	1.07±0.06 BCa	0.82±0.02	0.66±0.01 Da
				CDa	
Cultivar 2	0.45±0.07 Ab	0.37±0.01 ABb	0.34±0.02 ABb	0.3±0.03 Bb	0.24±0.01 Bb
DM (%)					
Cultivar 1	9.93±0.4 Aa	8.28±0.7 Aa	9.05±0.51 Aa	9.39±0.32 Aa	12.91±0.41
					Ba
Cultivar 2	16.37±0.92	14.68±0.96 Ab	14.7±1.04 Ab	20.13±2.70 Ab	30.32±3.48
	Ab				Bb

117 Means followed by the same capital letter, in the same line, do not differ by the Tukey test ( $P$   
118  $<0.05$ ) and means followed by the same lowercase letter, in the same column, do not differ  
119 from each other by the  $t$ -test ( $P <0.05$ ).

120 Figure 1A illustrates the influence of EC on the GR of seedlings from both cultivars. Cultivar  
121 1 did not suffer influence in the number of germinated seedlings, unlike cultivar 2, which can  
122 also be observed in table 1. In cultivar 2, the germination decreased from 91%, in the control  
123 group, to 37% for the highest EC tested, demonstrating its sensitivity to high levels of


124 salinity. Studies conducted by the authors [21] demonstrate that may exist different levels of  
125 salinity tolerance during the germination process among commercial lettuce cultivars.

126 The absence of influence on the seedlings GR of cultivar 1 may indicate that it tolerates high  
127 saline concentrations in this phenological phase. Salinity tolerance in the germination  
128 process is associated with the activation of mechanisms that lead to changes in cellular  
129 metabolism. Among these mechanisms, there is the ability to accumulate organic ions and  
130 low molecular weight solutes in cell vacuoles, to maintain water absorption, even in  
131 unfavourable osmotic conditions [22].

132 According to Figure 1B, GSI was different between the evaluated cultivars. For cultivar 1, it  
133 was practically constant, not differing between the treatments (Table 1). Conversely, a  
134 contrary effect can be observed for cultivar 2, where the decreasing curve in Figure 1B  
135 demonstrates how affected it was. These results reinforce the argument previously  
136 mentioned, regarding the cultivar 1 tolerance and the cultivar 2 susceptibility when submitted  
137 to saline environments. Studies carried out by the authors [23] confirm that the seeds tend to  
138 be vulnerable when subjected to the effects of salinity. The GSI of the seeds can be altered  
139 by a forced dormancy. The dormancy occurs due to the decrease in the water absorption by  
140 the seeds, which negatively affects the imbibition and, consequently, reduces the elongation  
141 and divisions of the cells, thus preventing the mobilization of indispensable reserves in the  
142 germination process [24,23].


143 The adjusted curves in Figure 1C show the behaviour of the DM%. This variable is  
144 completely associated with the water intake of the seedlings, since the higher dry mass/fresh  
145 mass ratio, the lower water content in plant tissues [25]. Results obtained by the authors [26]  
146 showed that the water consumption by lettuce plants is reduced linearly by the increase of  
147 the EC induced by the addition of salts, and it can be reduced by up to 45% when EC is  
148 raised to  $10.4 \text{ dS m}^{-1}$ . In the present study, from the values of the DM% variable, it can be  
149 inferred that the water consumption of the cultivars was differently affected. The adjusted  
150 curves for the DM% variable show that the water consumption in cultivar 1 was not  
151 influenced by the EC increase, while in cultivar 2 it was reduced when the EC reached  $5.0$   
152  $\text{dS m}^{-1}$ .

153 The growth of the seedlings' shoot and the root system was reduced with the progressive  
154 increase of the salinity (Figure 1D and 1E). Similar results were found by the authors [21]  
155 when studying the influence of salinity on two lettuce cultivars. Seedling development is  
156 affected by the decline of phytohormones levels, such as auxins, gibberellins, jasmonic acid  
157 and salicylic acid in plant tissues. This decrease is associated with the toxic effect of NaCl  
158 excess on plants [27,28]. The reduction of these plant hormones along with a decrease in  
159 the osmotic potential, caused by saline stress, results in a smaller number of cell divisions.  
160 Consequently, lower vegetative growth of the seedlings occurs [29,27].



**Fig. 1. Germination (A), germination speed index(B), the percentage of dry matter in relation to fresh matter (C), seedling height (D) and root length (E) in function of different levels of salinity.**

The linear correlation analysis (Figure 2A and 2B) showed that there is a positive correlation between root growth and shoot growth in both cultivars. It shows that, despite the negative influence of salinity on seedling development, the balance between these variables was not affected. Studies carried out by the authors [30,31, 32] in coffee, eucalyptus and potato cultivation, respectively, demonstrate the importance of good aerial and root system relationship for plants. Regarding lettuce cultivation, the relation of plant's root-shoot plays a fundamental role in the seedlings' formation. According to the authors [33], the increase in this ratio produces seedlings with superior quality, due to the better adhesion rates after transplanting.



**Fig. 2. Pearson's linear correlation between seedling height and root length for cultivar 1 (A) and cultivar 2 (B).** \*  $P < 0.05$ ; \*\*  $P < 0.001$ .

#### 4. CONCLUSION

Salinity affected root growth for both cultivars when EC was greater than  $2.8 \text{ ds m}^{-1}$ . For all the studied variables, cultivar 1 has shown better performance when compared to cultivar 2, allowing to affirm that it tolerates higher salinity rates in this phenological phase. The cultivar 2 was more susceptible to salinity since all the studied variables were statistically affected by different salt concentrations.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### AUTHORS' CONTRIBUTIONS

This work was carried out in collaboration AMONG ALL authors. All authors read and approved the final manuscript.

#### REFERENCES

- Boström M. A missing pillar? Challenges in theorizing and practising social sustainability: an introduction to the special issue. *Sustainability: Science, practice and policy*. 2012;8(1):3-14. DOI: 10.1080 / 15487733.2012.11908080
- Bretschger L. Is the Environment Compatible with Growth? Adopting an Integrated Framework for Sustainability. *Annual Review of Resource Economics*. 2017;9:185-207. DOI: 10.1146 / annurev-resource-100516-053332.
- Robertson MJ, Llewellyn RS, Mandel R, Lawes R, Bramley RGV, Swift L, O'Callaghan C. Adoption of variable rate fertilizer application in the Australian grains industry: status, issues and prospects. *Precision Agriculture*. 2012;13(2):181-199. DOI: 10.1007 / s11119-011-9236-3.
- Basso B, Cammarano D, Fiorentino D, Ritchie JT. Wheat yield response to spatially variable nitrogen fertilizer in Mediterranean environment. *EUR. J. Agron*. 2013;51:65-70. DOI: 10.1016 / j.eja.2013.06.007

- 201 5. Bretschger L, Smulders S. Sustainability and substitution of exhaustible natural  
202 resources: How structural change affects long-term R&D-investments. *Journal of*  
203 *Economic Dynamics and Control*. 2012;36(4):536-549. DOI: 10.1016 /  
204 j.jedc.2011.11.003.
- 205 6. Andrade EM. Irrigation and its implications on natural capital in arid and semi-arid regions:  
206 a review. *Ceres*. 2015; 56 (4): 390-398.
- 207 7. Oliveira JJ. et al. Spatial variability of chemical properties in a saline-sodium soil. *Brazilian*  
208 *Journal of Soil Science*. 1999; 23 (4): 783-789. DOI: 10.1590 / s0100-  
209 06831999000400004.
- 210 8. de Medeiros JF, Neto CPCT, Silva Dias N, Gheyi HR, Silva MVT, Loiola AT. Salinity and  
211 pH of an irrigated argisol with saline water under management strategies. *Brazilian*  
212 *Journal of Irrigated Agriculture*. 2017; 11 (3): 1407-1419. DOI: 10.7127 /  
213 rbai.v11n300560.
- 214 9. Azevedo LC, Martins ICS, Silva VL, Ribeiro CS, Cardoso L. Soil salinity in protected  
215 environment. *Digital Field*. 2018; 13 (1): 52-69.
- 216 10. Schossler TR, Machado DM, Zuffo AM, Andrade FD, Piauilino AC. Salinity: effects on  
217 physiology and mineral nutrition of plants. *Encyclopedia Biosphere*. 2012; 8 (15): 1563-  
218 1578.
- 219 11. Vital T, Sampaio Y. Family farming and irrigated fruit growing: case studies in the  
220 Northeast. *Annals of the Pernambuco Academy of Agronomic Science*. 2014; 4: 275-290.
- 221 12. Dias NS, Sousa Neto ON, Cosme CROJ, Rebouças JR, Oliveira AM. Response of  
222 lettuce cultivars to the salinity of the nutrient solution with saline waste in hydroponics.  
223 *Brazilian Journal of Agricultural and Environmental Engineering-Agriambi*. 2011; 15 (10):  
224 991-995. DOI: 10.1590 / s1415-43662011001000001.
- 225 13. Oliveira FA, Carrilho MJS, Medeiros JF, Maracajá PB, Oliveira MKT. Performance of  
226 lettuce cultivars submitted to different levels of irrigation water salinity. *Brazilian Journal*  
227 *of Agricultural and Environmental Engineering*. 2011; 15 (8): 771-777. DOI: 10.1590 /  
228 s1415-43662011000800002.
- 229 14. Viana SB, Fernandes PD, Gheyi HR. Germination and formation of lettuce seedlings at  
230 different levels of water salinity. *Brazilian Journal of Agricultural and Environmental*  
231 *Engineering*. 2001; 5 (2): 259-264. DOI: 10.1590 / s1415-43662001000200014
- 232 15. Zhu JK. Genetic analysis of plant salt tolerance using *Arabidopsis*. *Plant Physiology*.  
233 Rockville. 2000;124(3):941- 957. DOI: 10.1104 / pp.124.3.941
- 234 16. Ferreira LE, Medeiros JF, Silva NKC, Linhares PSF, Alves RC. Salinity and its effect on  
235 the production of grains of okra Santa Cruz 47. *Revista Verde*. 2012; 7 (4): 108-113.
- 236 17. Brazil. Rules for Seed Analysis. 1. ed. Brasília: Ministry of Agriculture, Livestock and  
237 Supply. 2009.
- 238 18. Oliveira AB, Gomes Filho E. Germination and vigor of forage sorghum seeds under  
239 water and saline stress. *Brazilian Journal of Seeds*. 2009; 31 (3): 48-56. DOI: 10.1590 /  
240 S0101-31222009000300005



- 241 19. Maguire JD. Speed of germination and in selection and evaluation for seedling  
242 emergence and vigor. *Crop Science*, 1962; 2 (1): 176-177. DOI: 10.2135 /  
243 cropsci1962.0011183x000200020033x.
- 244 20. Birth WM, Dias DCFS, Silva PP. Physiological quality of seed and establishment of  
245 vegetables in the field. *Embrapa Hortaliças-Chapter in scientific book (ALICE)*. 2011.
- 246 21. Nasri N, Saïdi I, Kaddour R, Lachaâl M. Effect of salinity on germination, seedling growth  
247 and acid phosphatase activity in lettuce. *American Journal of Plant Sciences*. 2015; 20  
248 (6): 57. DOI: 10.1007 / s11738-010-0625-4.
- 249 22. Munns R, Tester M. Mechanisms of salinity tolerance. *The Annual Review of Plant*  
250 *Biology*. 2008; 59: 651-681. DOI: 10.1146 / annurev.arplant.59.032607.092911
- 251 23. Lima BG, Torres SB. Hydric and saline stress on seed germination of *Zizyphus joazeiro*  
252 Mart. (Rhamnaceae). *Caatinga Magazine*. 2009; 22 (4): 93-99.
- 253 24. Ferreira LGR, Rebouças MAA. Influence of the hydration / dehydration of cotton seeds  
254 on the overcoming of salinity effects on germination. *Brazilian Agricultural Research*.  
255 1992; 27 (4): 609-616.
- 256 25. Chaves MM, Pereira JS, Maroco J, Rodrigues ML, Ricardo CPP, Osório ML, Pinheiro C.  
257 How do plants cope with water stress in the field? Photosynthesis and growth. *Annals of*  
258 *botany*. 2002; 89 (7): 907-916. DOI: 10.1093 / aob / mcf105.
- 259 26. Paulus D, Paulus E, Nava GA, Moura CA. Growth, water consumption and mineral  
260 composition of lettuce cultivated in hydroponics with salt water. *Ceres*.2015; 59 (1): 110-  
261 117. DOI: 10.1590 / S0034-737X2012000100016.
- 262 27.Egamberdieva D, Shurigin V, Gopalakrishnan S, Sharma R. Growth and symbiotic  
263 performance of chickpea (*Cicer arietinum*) cultivars under saline soil conditions. *Journal*  
264 *of Biological and Chemical Research*. 2014:1-10.
- 265 28.LlanesA, Andrade A, Masciarelli O, Alemano S, Luna V. Drought and salinity alter  
266 endogenous hormonal profiles at the seed germination phase. *Seed Science Research*.  
267 2016;26(1):1-13.DOI: <https://doi.org/10.1017/S0960258515000331>
- 268 29.Mauromicale G, Licandro P. Salinity and temperature effects on germination, emergence  
269 and seedling growth of globe artichoke. *Agronomie*.2002;22:443-450.  
270 DOI:<https://doi.org/10.1051/agro:2002011>
- 271 30. Ramos LCS, Lima MMA, Carvalho A. Growth of root system and aerial part in young  
272 plants of coffee trees. 1982.41 (1): 93-99. DOI: 10.1590 / s0006-87051982000100009.
- 273 31. Reis GG, Ferreira Reis MDG, Costa IFI, Monte MA, Birth GA, Oliveira CHR. Root and  
274 shoot growth of clones of hybrids of *Eucalyptus grandis* x *Eucalyptus urophylla* and  
275 *Eucalyptus camaldulensis* x *Eucalyptus* spp submitted to two irrigation regimes in the  
276 field. *Revista Árvore*.2006; 30 (6): 921-931. DOI: 10.1590 / S0100-67622006000600007.
- 277 32. Aguiar Netto AO, Rodrigues JD, Nascimento Júnior, NA. Growth Analysis in Potato  
278 Culture (*Solanum Tuberosum* Ssp *Tuberosum*), Subjected to Different Irrigation Blades:  
279 Tuber-Aerial Reason, Specific Foliar Area, Foliar Area Ratio, and Foliar Mass Ratio.  
280 *Irriga*. 2018; 4 (1): 1-9.DOI: <https://doi.org/10.15809/irriga.1999v4n1p1-9>

281 33. Echer MM, Guimarães VF., Krieser CR, Abucarma VM, Klein J, Santos L, Dallabrida  
282 WR. Use of biostimulant in the formation of yellow passion fruit seedlings. Semina:  
283 Agrarian Sciences. 2006; 27 (3): 351-317. DOI: 10.5433 / 1679-0359.2006v27n3p351.  
284

UNDER PEER REVIEW