

1
2
3
4
5
6
7
8
9
10

Original Research Article

Reduction of the forage seeds physiological quality by the contact with fertilizer in the crop-livestock integration

ABSTRACT

The aim of this study was to evaluate the effects of *Brachiaria* seed contact time with 05-25-15 fertilizer (N-P₂O₅-K₂O) on the seed's physiological quality. The experimental design was a completely randomized design with four replicates. Treatments were arranged in a 5x4 factorial, being: five seeds contact times with fertilizer (0, 24, 48, 72 and 96 hours); four forages: Marandu palisade grass, Piatã palisade grass, Xaraés palisade grass and Ruziziensis grass. The performed tests were: water content, germination (first count and germination percentage), tetrazolium (viability), electrical conductivity, accelerated aging, sand emergency and emergence speed index. Contact time of 05-25-15 fertilizer with *Brachiaria* seeds reduces linearly: germination, emergence and emergence speed index of *Brachiaria* seeds; Piatã and Xaraés palisade grass seeds viability; vigor (electrical conductivity), except in Ruziziensis grass seeds; and seeds germination of Piatã palisade grass submitted to accelerated aging test (vigor). Piatã palisade grass is the least recommended, among the analyzed grasses, for intercropping in simultaneous sowing.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

Keywords: *Brachiaria*; emergence; germination; grass; vigor.

1. INTRODUCTION

The Central-West region has maintained monoculture and conventional agriculture for decades, as well as livestock activity with pastures in an advanced degradation stage. Reduction in fertility and low soil water retention in degraded pasture areas reduce the quantity and quality of feed offered to animals [1], compromising cattle potential productive, as well as negatively impacting the environment. In addition to the problems of monoculture and area degradation, the cattle potential productivities is reduced due to the forage production seasonality.

Also, there is a need to increase the income of agriculture, due to the increasing food demand, the need to reduce opening of new areas by existing laws, and reduce the greenhouse gases emission [2,3]. An alternative to mitigate these impacts and promote socioeconomic and sustainable development is the integrated crop-livestock system (ICLS), through the consortium of grain crops, such as maize, with forage plants, mainly the *Brachiaria* (Syn. *Urochloa*) genus, increasing the production system efficiency [3].

In the ICLS, in case of the consortium of grain crops, such as maize, with forage plants, such as *Brachiaria*, the simultaneous sowing can be used to minimize problems such as lack of uniformity and reduced initial forage stand due to the reduced size and low weight of *Brachiaria* seeds. *Brachiaria* seeds are mixed with fertilizer recommended for sowing the

33 grain crop, being distributed at the same depth. Thus, the fertilizer will be used partially by
34 the forage, which will show slow development until the grain harvest.

35

36 However, there's a possibility of damage to the forage seeds physiological quality[4],
37 resulting in economics losses due the lower initial stand and consequent lower forage
38 production, due to the fertilizer harmful effects on seeds. Results shows that *Brachiaria*
39 *brizantha* [5] and *Brachiaria ruziziensis* [6] contact time with fertilizer, decreased the seeds
40 germination and vigor. Therefore, it was aimed to evaluate the effects of *Brachiaria* seed
41 contact time with 05-25-15 (N-P₂O₅-K₂O) fertilizer on the forage seed physiological quality.

42

43 2. MATERIAL AND METHODS

44

45 The experimental design was a completely randomized design with four replicates. The
46 treatments were arranged in a 5x4 factorial, being: five *Brachiaria* seeds contact times (0,
47 24, 48, 72 and 96 hours) with 05-25-15 (N-P₂O₅-K₂O) fertilizer, commonly used in maize
48 crop sowing; four forages: *Brachiaria brizantha* (cv. Marandu, BRS Piatã e Xaraés),
49 (Marandu palisade grass, piatã palisade grass and xaraés palisade grass, respectively) and
50 *Brachiaria ruziziensis* (cv. Kennedy), (Ruziziensis grass).

51 Based on the cultural value of the forage seed lots, the sowing rate of 4 kg of pure viable
52 seeds per hectare and the fertilizer quantity of 100 kg P₂O₅ ha⁻¹, required for sowing corn,
53 the ratio between seeds and fertilizer was adjusted.

54 Elapsed pre-established times, fertilizer and seeds were separated, and the following tests
55 were carried out: water content, germination (first germination count and germination
56 percentage), tetrazolium (viability), electrical conductivity, accelerated aging, emergence in
57 sand and emergence speed index.

58 To quantify water content, three samples of 4.0 g were used for each treatment, which were
59 placed in an oven for 24 hours at a temperature of 105 ± 1°C. After the drying process the
60 samples were placed in desiccators to promote the cooling and then the weight was
61 obtained using an analytical balance (0.0001 g) [7].

62 The methodology used for the germination test was described to the methodology of the
63 authors cited in the reference [7], using four subsamples of 50 seeds of each forage species
64 and fertilizer contact time with the forage seeds. The forage seeds were equidistantly
65 distributed in "gerbox" type germination boxes, on two sheets of blotting paper as substrate,
66 and moistened with distilled water in the proportion of two and a half times of the paper dry
67 mass.

68 Subsequently, they were taken to the biochemical oxygen demand (BOD) chamber with
69 photoperiod regulation of 12 hours and alternating temperature of 35/20 °C. Forage seeds
70 were analyzed at seven and 21 days, considering that the seeds that had emitted 2 mm of
71 primary root germinated.

72 For tetrazolium test, four subsamples of 50 seeds were used for each forage species and
73 the fertilizer contact time with forage seeds. The seeds were pre-wetted between paper
74 sheet for germination, conditioned in a BOD chamber with a temperature of 30 °C for 18
75 hours without light. Then, the longitudinal sectioning of the seeds was performed, and one
76 part of the seeds was immersed in 0.5% tetrazolium salt solution (2,3,5-triphenyl chloride-
77 tetrazolium). It was placed in a BOD chamber at a temperature of 30 °C for 3 hours for
78 staining the living tissue of forage seeds. In the sequence it was classified into viable and
79 non-viable seeds, according to the methodology of the authors cited in the reference [7].

80 In electrical conductivity analysis, four subsamples of 50 seeds were used for each
81 treatment, which were weighed in an analytical balance (0.0001 g), placed in a container
82 containing 75 mL of distilled water and conditioned in a BOD chamber with a temperature of
83 25 ° C for 24 hours, without light. Then, the exudates released through the conductivity
84 meter were measured, according to the methodology of the authors cited in the reference
85 [8].

86 The methodology of the authors cited in the reference [9] was adopted for accelerated aging
87 analysis, in which the seeds of each treatment were distributed on aluminum screen coupled
88 to gerboxes containing 40 mL of distilled water. The boxes were then capped, forming a wet
89 chamber, and placed in a BOD chamber for 36 hours at 42 ° C, without light. Finally, the
90 germination test was carried out with the aged forage seeds, according to the methodology
91 of the authors cited in the reference [7].

92 For emergency sand test, sowing was carried out in trays (29.1 x 23.0 x 5.3 cm) with 2.5 kg
93 of sterilized sand at 105 ± 1 °C for 24 hours. Seeds were seeded for each treatment, in 10
94 mm depth, maintained at 12 hours of light incidence and humidity around 65% of the field
95 capacity, for daily maintenance of the initial weight [9]. Daily analysis of the seeds was
96 performed up to 21 days, considering seedlings with 2 mm above the substrate level
97 emerged at least. Subsequently, emergency speed index (ESI) was calculated using to the
98 methodology of the authors cited in the reference [10].
99

100 In variance analysis was considered the forage species effect, contact time and the
101 interaction between them. For contact time analysis, a linear regression test was performed
102 for each forage, and the coefficients were submitted to the t test. For all tests it was
103 considered a 5% probability level. Was used the SANEST software [11].
104

105 3. RESULTS AND DISCUSSION

106
107 It was verified that there was contact time effect and no interaction between forages and
108 contact times, which shows that seeds contact time with 05-25-15 fertilizer does not interfere
109 in water content. An average water content of 11.42% was observed.

110 The absence of a change in water content may have been due to the fact that the seed
111 mixture with fertilizer was stored in semi permeable packaging (plastic bags), making it
112 difficult to absorb air moisture by both components, as well as the establishment of
113 hygroscopic balance inside the package.
114

115 Another factor is that the seeds hygroscopicity varies according to the chemical constitution,
116 especially the proteins, which have high affinity for water [12], and fatty acids that present a
117 hydrophobic characteristic, with an antagonistic relation to the seed's contents [13], besides
118 to being influenced by genetics, environmental conditions and plant traits [14]. Based on this,
119 it is inferred that the hydrophilic substances content of the species studied in this work may
120 be low.
121

122 In contrast to what was observed in this work, the authors cited in the reference [5] observed
123 an increase in water content of marandu palisade grass with contact time with 04-14-
124 08fertilizer ($N_2-P_2O_5-K_2O$; urea, triple superphosphate and potassium chloride), attributing
125 this effect to the high hygroscopicity of urea, which may have transferred air moisture to
126 seeds. One of the negative effects on the water content increase is the increase in seed's
127 metabolic activity, providing lower physiological quality [6].
128

129 Corroborating the above results, the authors cited in the references [15, 16, 17] verified an
 130 increase in seeds water content of marandu palisade grass as the contact time with triple
 131 superphosphate, single superphosphate and potassium chloride, respectively. The authors
 132 justify these results due to the seed coat rupture and absorption of environment moisture, as
 133 a consequence of obtaining the acid phosphate fertilizers, and the high salinity index of
 134 potassium chloride. Although the fertilizer used in this work consists of single and triple
 135 superphosphate and potassium chloride, which constitutes high salt content and acid pH, the
 136 effect described above was not evidenced.

137
 138 There was no effect of contact time and forage interaction on the first and final evaluation of
 139 the germination percentage of *Brachiaria* seeds ($P = .05$), (Table 1).


140
 141 **Table 1. First (1E) and final evaluation (FE) of seeds germination (%) of Marandu**
 142 **palisade grass, Piatã palisade grass, Xaraés palisade grass and Ruziziensis grass**
 143 **submitted to contact with fertilizer and Linear Regression (L).**

Analysis	Contact time (hours)					
	0	24	48	72	96	L
1E	55.43	57.48	50.94	51.13	44.97	0.011
FE	59.80	64.21	54.96	55.83	47.46	0.001

144 * Significant at 5% probability.

145

146 There was a linear decreasing effect of forage **seeds contact** time with fertilizer on the first
 147 and final evaluation of seed's germination, regardless of the forage (Figure 1). Comparing
 148 the 96-hour treatment with the absence of contact, there was an average reduction of 18.87
 149 and 20.63% in the first and final evaluation of the germination, respectively.



150

151 **Figure 1. First and final **evaluation of** seeds germination (%) of Marandu palisade**
 152 **grass, Piatã palisade grass, Xaraés palisade grass and Ruziziensis grass submitted to**
 153 **contact with fertilizer and Linear Regression (L). *Significant at 5% probability.**

154

155 Probably, the effect was justified by the tegument rupture and electrolytes extravasation,
 156 since the contact with fertilizer can act negatively on the seed coat and consequently inhibit
 157 the germination by the saline effect [18], besides the acid residues from the process of

158 obtaining phosphate fertilizers [5]. Another influencing factor is probably the rapid water
 159 absorption by the seed during the imbibition phase, causing damage to seed tissues [19,20].
 160

161 The decrease in germination percentages obtained in first and final evaluation of germination
 162 test of forage seeds was also verified by the authors cited in the references [15, 5], indicating
 163 deleterious effect of marandu palisade grass seeds contact time with superphosphate
 164 fertilizers triple and formulated 04-14-08 on seeds vigor, with reductions of 95.74 and
 165 82.41%, respectively, in germination obtained with the first evaluation of the germination
 166 test. In addition, the authors also observed a reduction in seed germination percentage in
 167 the final evaluation of the germination test after being conditioned for a period superior to
 168 three and 36 hours, respectively.
 169


170 There was interaction effect of contact time and forage on viability ($P = .05$), (Table 2).
 171

172 **Table 2. Seeds viability (%) of Marandu palisade grass, Piatã palisade grass, Xaraés**
 173 **palisade grass and Ruzizensis grass submitted to contact with fertilizer and Linear**
 174 **Regression (L).**

Forages	Contact time (hours)					L
	0	24	48	72	96	
Marandu	76.00	70.00	70.50	73.50	72.00	0.390 ^{ns}
Piatã	80.00	62.5	57.50	71.00	57.00	0.000*
Xaraés	73.00	56.00	59.50	55.50	36.50	0.000*
Ruzizensis	62.83	65.33	65.00	65.17	62.83	0.978 ^{ns}

175 * Significant at 5% probability ($P = .05$); ns: not significant at 5% probability ($P = .05$).
 176

177 There was a linear decreasing effect of seeds contact time with fertilizer on the seed's
 178 viability of the Piatã and Xaraés palisade grasses (Figure 2).



179
 180
 181 **Figure 2. Seeds viability (%) of Piatã and Xaraés palisade grass submitted to contact**
 182 **with fertilizer and Linear Regression (L). *Significant at 5% probability ($P = .05$)**
 183

184 Comparing the 96-hour treatment with the absence of contact, there was an average reduction
 185 of 28.75 and 50% in the seed's viability of the species, respectively. In addition, when
 186 analyzing the germination percentage at zero contact time and viability at the same time, the
 187 dormancy phenomenon is observed, as reported in studies of the authors cited in the
 188 references [21, 22] (Figure 2).

189

190 In Piatã and Xaraés palisade grasses the seeds contact with fertilizer caused damage to seed
191 embryos. Probably, due to the intrinsic factors of the fertilizer (high salinity, pH and electrical
192 conductivity) causing tegument ruptured and injured the seed embryo. Based on the eminent
193 effect only on the forage described previously, it is inferred that they have a tegument more
194 susceptible to damages than the other forages studied.

195

196 More vigorous seeds have higher soluble proteins amounts than those with less vigor [23].
197 Therefore, it is necessary to study the constitution of seed cover of *Brachiaria* seeds and the
198 use of Piatã and Xaraés palisade grasses in integrated systems under a simultaneous sowing
199 system.

200

201 In a tetrazolium test carried out on marandu palisade grass seeds, submitted to contact with
202 granulated monoammonium phosphate, granular superphosphate and superphosphate
203 powder, the authors reported in the reference [16] observed a linear decreasing effect of the
204 contact time of all fertilizers with the seeds on viability.

205

206 Interaction ($P = .05$) was observed between the forages and the contact time on the electrical
207 conductivity, which shows that there was an effect of the fertilizer contact time for each
208 *Brachiaria* cultivars. There was a linear effect of contact time with fertilizer on the electrical
209 conductivity of all cultivars, except for Ruziziensis grass (Table 3).

210

211 **Tabela 3. Electrical conductivity ($\mu^{-1} \text{ cm}^{-1} \text{ g}^{-1}$) of seeds of Marandu palisade grass,**
212 **Piatã palisade grass, Xaraés palisade grass and Ruziziensis grass submitted to**
213 **contact with fertilizer and Linear Regression (L).**

Forages	Contact time (hours)					L
	0	24	48	72	96	
Marandu	22.74	70.30	77.92	104.57	66.42	< 0.0001
Piatã	51.53	75.02	79.13	102.76	99.47	< 0.0001
Xaraés	63.78	87.72	89.16	95.21	82.39	< 0.0001
Ruziziensis	57.44	91.76	53.51	55.31	67.64	0.284 ^{ns}


214

* Significant at 5% probability ($P = .05$); ns: not significant at 5% probability ($P = .05$).

215

216 As the seeds of Marandu, Piatã and Xaraés palisade grasses contact time with fertilizer
217 increased, there was an increase in the electrical conductivity (Figure 3), which shows a
218 reduction in seed vigor. When comparing the absence of contact with the maximum time
219 studied in this work, it was verified that there was an average increase of 192, 93 and 29% in
220 the electrical conductivity of the cultivars Marandu, Piatã, and Xaraés palisade grass,
221 respectively.

222



223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253

Figura 3. Electrical conductivity (EC; $\mu^{-1} \text{ cm}^{-1} \text{ g}^{-1}$) of seeds of Marandu, Piatã and Xaraés palisade grass and Linear Regression (L). *Significant at 5% probability (P = .05).

The results were because the 05-25-15 fertilizer consisted of monoammonium phosphate, single superphosphate, triple superphosphate and potassium chloride, and some of these fertilizers present acid residues or high salt content, which probably caused seed coat rupture and electrolytes release contained in the reserve substances [16] and is indicative of reduced vigor. In addition, in the electrical conductivity results, the fertilizer residue effect on the seeds outer layer can occur, since they were not washed before the beginning of the test.

Similar results were evidenced by authors cited in the references [16, 17], who studied the contact of marandu palisade grass seeds with simple superphosphate and ammonium monophosphate in granulated, ground and powder forms; and potassium chloride, respectively, observed a proportional increase in the seeds electrical conductivity values.

The absence of effect of 05-25-15 fertilizer contact time on the electrical conductivity of Ruzizensis grass seeds should be studied, since this indicates a higher resistance of this species to physiological damages. Thus, it can be seen that up to 96 contact hours, there was no reduction in Ruzizensis grass seeds vigor, which allows the producer greater flexibility in sowing activities, being able to carry out a mixture of seeds and fertilizers in advance.

There was an interaction effect (contact time versus forage grass) on seeds germination submitted to accelerated aging, therefore, an individual effect was evidenced for each forage tested. There was no effect of the contact time with fertilizer for the forage tested, except for Piatã palisade grass, in which a significant linear effect was observed (Table 4).

Table 4. Seeds germination (%) of Marandu palisade grass, Piatã palisade grass, Xaraés palisade grass and Ruzizensis grass submitted to contact with fertilizer after accelerated aging and Linear Regression (L).

Forages	Contact time (hours)					L
	0	24	48	72	96	
Marandu	76,00	70,00	70,50	73,50	72,00	0,390 ^{ns}
Piatã	80,00	62,5	57,50	71,00	57,00	0,000 [*]
Xaraés	73,00	56,00	59,50	55,50	36,50	0,000 [*]
Ruzizensis	62,83	65,33	65,00	65,17	62,83	0,978 ^{ns}

254 * Significant at 5% probability ($P = .05$); ns: not significant at 5% probability ($P = .05$).

255

256 Comparing the 96-hour treatment with the absence of contact, there was an average reduction
257 of 38.46% in seeds germination of Piatã palisade grass submitted to aging. The effect was not
258 evidenced in the other studied species studied (Figure 4).


259

260 The high temperature at which the seeds were submitted during aging may accelerate
261 deterioration process, as it causes degenerative changes in seed metabolism, such as protein
262 denaturation, besides to causing rapid dry seeds imbibition, which favors injuries, and / or
263 mobilization of reserves and energy release through respiration, culminating in reduced
264 germination after accelerated aging [24,25].

265

266 Conversely, in some cases, there is an increase in germination after accelerated aging, which
267 may be associated with dormancy [26] or even with pathogen control.

268



269

270

271 **Figure 4. Seeds germination (%) of Marandu palisade grass, Piatã palisade grass,**
272 **Xaraés palisade grass and Ruziziensis grass submitted to contact with fertilizer after**
273 **accelerated aging and Linear Regression (L). *Significant at 5% probability ($P = .05$)**

274

275 The authors cited in the reference [26] evaluating seeds of *Brachiaria brizantha* cv. MG-5
276 Vitória, concluded that the accelerated aging method overcomes dormancy and favors seed
277 germination. However, the effect was only evidenced when accelerated aging was carried out
278 at 41 °C for a 96 hours period.

279

280 From the viability results (Table 2), electrical conductivity (Table 3) and accelerated aging
281 (Table 4), it is observed that Piatã palisade grass stands out from the other forages studied to
282 the effect of contact with fertilizer, presenting physiological damage more severe, intensified
283 by the increase in fertilizer contact time with the seeds. Therefore, precaution is recommended
284 in the use of Piatã palisade grass in detriment of the other forages studied for the crop-
285 livestock integration in simultaneous sowing system.

286

287 For the first and final evaluation of the emergence percentage, the effect of the seeds contact
288 time with fertilizer was observed. There was no interaction effect between forages and contact
289 time, therefore, for all the forage species analyzed, a similar effect was observed on the first
290 count and emergence percentage (Table 5).

291
292
293
294


Tabela 5. First (1E) and final evaluation (FE) of seeds emergence (%) of Marandu palisade grass, Piatã palisade grass, Xaraés palisade grass and Ruziziensis grass and Linear Regression (L)

Analysis	Contact time (hours)					L
	0	24	48	72	96	
1E	50.00	48.41	47.90	48.00	42.20	0.006*
FE	63.53	64.41	61.43	62.97	54.77	0.001*

295
296
297
298
299

* Significant at 5% probability (P = .05).

It was observed a reduction in the first and final evaluation of the *Brachiaria* seeds emergence with the increase of the contact time with the 05-25-15 fertilizer (Figure 5).



300
301
302
303

Figura 5. First and final evaluation of seeds emergence (%) of Marandu palisade grass, Piatã palisade grass, Xaraés palisade grass and Ruziziensis grass and Linear Regression (L). *Significant at 5% probability (P = .05).

304
305
306
307
308

Comparing the 96-hour treatment with the absence of contact, there was a mean reduction of 15.60 and 13.79% in the first and final evaluation of the emergence percentage, respectively (Figure 5).

309
310
311
312
313

Studying emerged seedlings percentage of *Brachiaria brizantha* cv. MG-5 and the emergence speed, the authors cited in the reference [4] observed damages by application of the 08-28-16fertilizer next to the seed, regardless of the seeding depth. In this study, the authors reported a decrease in emergence percentage in Marandu palisade grass subjected to contact with 04-14-08 fertilizer and potassium chloride, respectively.

314
315
316
317
318
319

Besides to direct damage caused by the contact of the fertilizer with the seeds, substances exuded from the seeds (mainly sugars) due to the tegument rupture (caused by high acidity, electrical conductivity and fertilizer pH) may have stimulated the development of microorganisms causing damage to seedlings establishment.

320
321
322

There was no interaction effect of contact time and forage on emergency speed index (ESI), (P = .05), (Table 6).


323 **Table 6. Emergence speed index (ESI) of Marandu palisade grass, Piatã palisade grass,**
 324 **Xaraés palisade grass and Ruzizensis grass and Linear Regression (L).**

Analysis	Contact time (hours)					L
	0	24	48	72	96	
ESI	46.03	45.69	43.21	44.06	37.15	0.001*

325 * Significant at 5% probability (P = .05).

326
 327 There was a linear decreasing effect of the seeds contact time with fertilizer on the ESI,
 328 independent of the forage species/cultivar (Figure 6). Comparing the 96-hour treatment with
 329 the absence of contact, there was an average reduction of 19.29% in ESI.

330
 331



332 **Figure 6. Emergence speed index of Marandu, Piatã, Xaraés and Ruzizensis grasses**
 333 **and Linear Regression (L). *Significant at 5% probability (P = .05).**

334
 335 The ESI reduction is a consequence of the interaction of seeds physiological potential with
 336 environmental conditions. Seed lots of forage species may have declined in physiological
 337 potential due to being subject to environmental changes, under conditions of temperature far
 338 from ideal, not withstand stress.

339
 340 Besides that, substances exuded from seeds (mainly sugars) due to tegument rupture (high
 341 acidity, electrical conductivity and fertilizer pH) may have stimulated the development of
 342 microorganisms, causing damage to seedling establishment.

343
 344 The ESI decrease of marandu palisade grass seeds with the advance of the fertilizer contact
 345 time was also verified by the authors cited in the references [15, 5] when using the
 346 superphosphate triple fertilizers and 04-14-08, respectively. The authors verified effect from
 347 12 and 3 hours of contact with fertilizers, respectively.

348
 349

350 4. CONCLUSION

351
 352 Water content of *Brachiaria* seeds is not altered by the contact time with fertilizer 05-25-15.

353
 354 Contact time of 05-25-15 fertilizer with *Brachiaria* seeds reduces linearly: germination,
 355 emergence and emergence speed index of *brachiaria* seeds; Piatã and Xaraés palisade

356 grass **seeds viability**; vigor (electrical conductivity), except in Ruziziensis grass seeds; and
357 seeds germination of Piatã palisade grass submitted to accelerated aging test (vigor).

358

359 Piatã palisade grass is the least recommended, among the analyzed grasses, for
360 intercropping in simultaneous sowing.

361

362

363 **COMPETING INTERESTS**

364

365 We declare that no competing interests exist.

366

367

368 **REFERENCES**

369

370 1. Macedo, M.C.M; Zimmer, A.H. Integrated crop-livestock systems in the cerrado region of
371 Brazil. International Symposium on Livestock-Livestock Integration. Curitiba: Embrapa Gado
372 de Corte.2007. **Accessed 02 Jan 2019.**

373 Available:[http://www.fcav.unesp.br/Home/departamentos/zootecnia/anaclaudiaruggieri/12.-
374 sistemas-integrados-de-lavoura-pecuaria---cerrado.pdf](http://www.fcav.unesp.br/Home/departamentos/zootecnia/anaclaudiaruggieri/12.-sistemas-integrados-de-lavoura-pecuaria---cerrado.pdf)

375

376 2. Balbino Junior AA, Moraes A, Veiga M, Pelissar A, Dieckow J. Crop-livestock integration:
377 intensification of the use of agricultural areas. English. Rural Science. 2009; 39 (6): 1925-
378 1933. DOI: 10.1590 / S0103-84782009005000107

379

380 3. Vilela L et al. Integration systems for livestock farming in the Cerrado region. Agricultural
381 Research. 2011; 46 (10): 1127-1138. English. DOI: 10.1590 / S0100-204X2011001000003

382

383 4. Foloni JSS, Custódio CC, Pompei FP, Vivan MR. Planting of forage species due to soil
384 depth and contact with formulated fertilizer NPK. Tropical Agriculture Research. 2009; 39
385 (1): 7-12. English.

386

387 5. Lima EDV, Tavares JC, Azevedo VR, Leitão-Lima PS. Mixture of Brachiariabrizanth seeds
388 with NPK fertilizer. Rural Science. 2010; 40 (2): 471-474. DOI: 10.1590 / S0103-
389 84782010005000003

390

391 6. Dan HA, Dan LGM, Barroso ALL, Braccini AL, Puccinin GG. Mixture of seeds of
392 Brachiaria ruziziensisG.et E with urea aiming at the implantation of the crop-livestock
393 integration system. Caatinga Magazine. 2011; 24 (4): 68-73. English.

394

395 7. Brazil. Ministry of Agriculture, Livestock and Supply. Rules for seed analysis.Brasília:
396 MAPA / ACS; 2009.

397

398 8. AOSA. Association of Official Seed Analysts. Seed Vigour Testing Handbook.
399 Washington: AOSA, 1983.

400

401 9. Marcos Filho J. Accelerated aging test. In: Krzyzanowski FC, Vieira RD, France Neto JB.
402 Seed vigor: concepts and tests. Londrina: Abrates; 1999.

403

404 10. Maguire LD. Speed of germination aid in selection and evolution forseedling emergence
405 and vigor. Crop Science. 1962, 2: 176-177. DOI: 10.2135 /
406 cropsci1962.0011183X000200020033x

407

- 408 11. Zonta EP, Machado AD. Statistical analysis system for microcomputers - SANEST.
409 Pelotas: Federal University of Pelotas; 1984. 150p. Portuguese.
410
- 411 12. Marcos Filho J. Physiology of seeds of cultivated plants. Londrina: ABRATES; 2015.
412
- 413 13. Delarmelino-Ferraresi LM, Villela FA, Aumonde TZ. Physiological performance and
414 chemical composition of soybean seeds. Brazilian Journal of Agricultural Sciences. 2014; 9
415 (1): 14-18. English. DOI: 10.5039 / agraria.v9i1a2864
416
- 417 14. Martins MTCS, Pôrto NA, Canuto MFS, Bruno RLA. Chemical Composition of
418 ManihotMill Species Seeds. (Euphorbiaceae). Brazilian Journal of Biosciences. 2007; 5 (1):
419 621-623. Portuguese.
420
- 421 15. Lima EDV, Tavares JC, Silva EC, Leitão-Lima P. Triple superphosphate as a way of
422 distribution of Brachiariabrizantha seeds for renewal of pastures in the Amazon. Revista
423 Brasileira de Zootecnia.2009; 38 (5): 796-800. English. DOI: 10.1590 / S1516-
424 35982009000500003
425
- 426 16. Peres AR, Vazquez GH, Cardoso RD. Physiological potential of Brachiariabrizantha cv.
427 Maranduseedskept in contactwithphosphaticfertilizers. Brazilian Journal of Seeds. 2012; 34
428 (3): 424-432. DOI: 10.1590 / S0101-31222012000300009
429
- 430 17. Lima EV, Tavares JCS, Leitão-Lima PS, Pinheiro DP. Contact periods of the KCl
431 fertilizer on the physiological quality of BrachiariabrizanthaStapf seeds. Amazon: Science &
432 Development. 2013; 8: 53-64. English.
433
- 434 18. Mateus GP, Borghi E, Marques RR, Bôas RLV, Crusciol CAC. Fertilizer sources and
435 contact periods and seed germination of Brachiariabrizantha. Brazilian Journal of Soil
436 Science. 2007; 31 (1): 177-183. English. DOI: 10.1590 / S0100-06832007000100018
437
- 438 19. Derré LO, Custódio CC, Agostini EAT, WEX War. Obtaining the embedding curves of
439 coated and uncoated seeds of Urochloabrizanthae Urochloa ruziziensis.
440 ColloquiumAgrariae. 2013; 9 (2): 103-111. English. DOI: 10.5747 / ca.2013.v09.n2.a094
441
- 442 20. Toledo LF, Carvalho-E-Silva SP, Sánchez C, Almeida MA, Haddad CFB. The review of
443 the Brazilian Forest Act: harmful effects on amphibian conservation. Biota Neotropica. 2010;
444 10 (4): 35-38. DOI: 10.1590 / S1676-06032010000400003
445
- 446 21. Dias MCLL, Alves SJ. Evaluation of the viability of Brachiariabrizantha (Hochst. Ex A.
447 Rich) Stapf seeds by the tetrazolium test. Brazilian Journal of Seeds. 2008; 30 (3): 145-151.
448 English. DOI: 10.1590 / S0101-31222008000300019
449
- 450 22. Pariz CM, Ferreira RL, Sá ME, Andreotti M, Chioderoli CA, Ribeiro AP. Physiological
451 quality of Brachiaria seeds and evaluation of dry mass yield in different systems of crop-
452 livestock integration under irrigation. Tropical Agriculture Research. 2010; 40 (3): 330-340.
453 DOI: 10.5216 / pat.v40i3.6590
454
- 455 23. Henning FA, Mertz LM, Jacob Junior EA, Machado RD, Fiss G, Zimmer PD. Chemical
456 composition and mobilization of reserves in high and low vigor soybean seeds. Bragantia.
457 2010; 69 (3): 727-734. Portuguese. DOI: 10.1590 / S0006-87052010000300026
458
- 459 24. Binotti FFS, Haga KI, Cardoso ED, Alves CZ, Sá ME, Arf O. Effect of the accelerated
460 aging period on the electrical conductivity test and on the physiological quality of bean

- 461 seeds. *Acta Scientiarum-agronomy*. 2008; 30 (2): 247-254. English. DOI: 10.1590 / S1807-
462 86212008000200014
- 463
- 464 25. Emer AA, Oliveira MC, Passos AI, Júnior Bertoncelli D. Biochemical composition and
465 vigor of soybean seeds under different conditions of accelerated aging. *Brazilian Journal of*
466 *Applied Technology for Agricultural Science*. 2015; 8 (3): 7-15. English. DOI: 10.5935 /
467 PAeT.V8.N3.01
- 468
- 469 26. Cardoso ED, Sá ME, Haga KI, Binotti FFS, Nogueira DC, Valério Filho WV.
470 Physiological performance and dormancy overcoming of *Brachiariabrizantha* seeds
471 submitted to chemical treatment and artificial aging. *Semina: Academic Sciences*. 2014; 35
472 (1): 21-38. DOI: 10.5433 / 1679-0359.2014v35n1p21

UNDER PEER REVIEW