

Biomass production and nutritional characterization of *Eucalyptus benthamii* in the Pampa Biome, Brazil

ABSTRACT

The objective of this study was to evaluate the biomass production and to characterize a 7-year-old *Eucalyptus benthamii* stands in the Pampa-RS Biome. Initially, a sample inventory was performed for the dendrometric characterization of the stand. For the determination of the biomass, nine trees were felled and fractionated in wood, bark, branch and leaves. Soil samples and plant tissues were collected and analyzed for nutritional characterization and determination of biological utilization coefficient (BUC). The average annual increment (AAI) with bark was $49.87 \text{ m}^3 \text{ ha}^{-1}$. The biomass production was 192 Mg ha^{-1} , distributed in wood (81.2%)> branches (11%)> bark (6,5%)> leaves (1,3%). The leaves component presented the highest nutrient concentration and the wood the highest amounts of nutrients allocated in the biomass, except for Ca and Mg, observed in the bark. The highest BUC was observed in the wood. Mg was the nutrient that provided the best efficiency with a yield of 6,014 kg of wood per kg of Mg used, followed by S, P, Ca, K and N.

Keywords: forest soil; productivity; harvest; forest nutrition; nutrient cycling.

1. INTRODUCTION

The natural population of *Eucalyptus benthamii* occurs to Australia, distributed along the eastern coast of New South Wales, southwest of the city of Sydney on alluvial plains on the banks of the Nepean River and its tributaries [1]. In this region, the species is distributed among only four populations, the largest of which consists of 6,550 trees and the other three with less than 340 trees. This condition places *E. benthamii* in extinction threat, with the following main factors: low seed viability due to the high degree of inbreeding and self-fertilization, low natural regeneration, competition with introduced species, changes in water regimes, fires, increase from urban areas and intense agricultural activity in the area of natural occurrence [2].

In relation to the silvicultural aspects, according to a study carried in the Colombo-PR region, the high frost tolerance stands out, supporting absolute minimum temperatures of up to -10°C , fast growth, uniform stem and high homogeneity of the plot [3]. These characteristics indicate that *Eucalyptus benthamii*, as a good alternative for silvicultural use in cold climate regions, especially where there is frequent and severe frost occurrence, as in southern Brazil [4].

The evaluation of the productivity, biomass and nutrient production at the end of the rotation of the forest plantation, can help in the decision making of the forester in relation to the choice of the species to be implanted and the nutritional replacement of the new production cycle. According to Viera et al. [5], the choice of efficient genotypes to absorb and use nutrients must be performed in order to improve applied fertilization.

In addition, the applied harvest intensity, removal of one or more components of the tree, directly affects the export of nutrients from the forest site. In order to maintain the soil productive capacity [5], the nutritional replacement cost would be increased from corrective fertilization and maintenance. According to Achat et al [6], the removal of the residues causes a decrease in the biological activity and an increase in soil compaction, as a result there is a decrease in growth.

Due to this, the objective of this work was to evaluate the biomass and nutrient production and determine the nutrient utilization efficiency of an experimental plantation of *Eucalyptus benthamii* Maiden & Cambage in the Pampa Biome of Rio Grande do Sul.

2. MATERIAL AND METHODS

2.1 Characterization of the experimental area

The present study was carried out in an area of 10 hectares of *E. benthamii*, seven years old, in the municipality of Alegrete-RS, with central geographical coordinates: 55° 32 '53' west longitude and 29° 47'60° south latitude.

The climate of the region is classified as humid sub-temperate, with frequent frosts from May to August, and intense heat in summer, mainly in the months of January and February, with the average temperature of the month being warmer > 22°C and average annual temperature > 18 °C. Annual precipitation presents rainfall indexes ranging from 1,250 to 1,500 mm [7]. The soil of the study area is classified as typical Distrophic Red Argisol[8].

The planting of the seedlings was done manually and without irrigation, using seminal seedlings, with initial density of 1428 plants ha⁻¹ (3.5 m x 2.0 m). Subsoiling was performed 30 days before planting, using a subsoiler with three stems, incorporating 300 kg ha⁻¹ of reactive natural phosphate (GAFSA, 12% P₂O₅ soluble in citric acid) followed by light harrowing.

Fertilization was carried out 15 days after planting, using the formula N-P₂O₅-K₂O from 06-30-06 + 0.6% B, 110 g plant⁻¹, divided into two sub-doses of 55 g incorporated at 15 cm distance on each side of the seedling. The second fertilization was carried out at 90 days post-planting, using fertilizer with formulation N-P₂O₅-K₂O from 20-05-20 + 0.2% B + 0.4% Zn, 122 g plant⁻¹, applied manually for the canopy projection. The third fertilization, at 270 days, was used the formula N-P-K₂O of 22-00-18 + 1% S + 0.3% B, 122 g plant⁻¹ applied mechanically in the interlining. At no time was the liming performed.

2.2 Experimental design and data collection

Through the forest inventory the growth variables were obtained. Four plots (35 m x 20 m) were randomly distributed, all diameters at breast height were measured (DBH) with diametric tape and all tree heights (m) with Vertex hypsometer. After the measurements were made the distribution of the trees by diameter class, where three classes were determined, the first class being from 10 to 16 cm, the second from 16.1 to 22 cm and the third class from 22.1 to 28 cm.

For the determination of above-ground biomass (Mg ha⁻¹), three trees were selected by diameter class, with a tree at the lower limit, a tree at the central limit and a tree at the upper limit of each class, totaling nine individuals slaughtered. The selected trees were sectioned at ground level, cubed by the Smalian methodology, as described by Finger [9]. After the

canopy, each felled tree was fractionated in the components: leaves, branches, bark and wood. Each component had its biomass measured in the field by weighing with hook scale, with accuracy of 50g.

A sample of leaves and branches was collected per evaluated tree. For wood and bark, three samples per tree distributed along the commercial shaft with a minimum diameter of 8 cm were collected, in the median positions of the sections resulting from the division into three equal parts of the same. All the samples were weighed in the field, later they were properly packed, identified and sent for analysis in laboratory.

The estimated tree biomass per hectare was determined by regression analysis applied to the inventory data and extrapolation based on the sample unit area. The amount of macronutrients (kg ha^{-1}) and micronutrients (g ha^{-1}), allocated to tree components, was obtained by multiplying the content of each nutrient by biomass.

For soil chemical analysis and density, samples were collected at depths of 0-20, 20-40 and 40-100 cm. Density determination followed the methodology proposed by Embrapa[10]. Plant tissue and soil to analyze were performed following the methodology described by Tedesco et al. [11] and Miyazawa et al. [12].

2.3 Statistics and Data Analysis

The Berkhout, Schumacher-Hall, Hohenadl-Kreen and Spurr models were used to determine the equations for height and volume estimation. The modeling of the equations for the individual biomass of the trees and their respective components was processed using the program "proc stepwise" - "forward" option of the statistical program SAS [13].

The biomass of each component and its arithmetic and logarithmic variants (natural logarithm) were considered as dependent variables. The independent variables were DBH (cm), height (m) and volume ($\text{m}^3 \text{ tree}^{-1}$) and their respective arithmetic and logarithmic variants (natural logarithm). The quality of the adjustment and selection of the equations considered as the main statistics, the highest adjusted coefficient of determination and the lowest relative standard error of the estimate ($\text{Syx}\%$).

The contrast of the averages of the chemical and physical attributes of the soil between the different depths and, for the nutrient contents in the components of the biomass (leaves, branches, bark and wood) was evaluated by the Tukey test at the level of 5% probability of error. A completely randomized design was used for the statistical analysis, where the treatments were the soil depths and the biomass components above the soil.

The biological utilization coefficient (BUC) was evaluated. The BUC can be described as the biomass nutrient conversion rate, obtained through the ratio between the biomass and the nutrient quantity, both with the same unit [14].

3. RESULTS AND DISCUSSION

3.1 Soil fertility

According to the Soil Chemistry and Fertility Commission - RS / SC [15], the soil of the experimental area presents: textural class 4 (clay content $\leq 20\%$); low organic matter content (≤ 2.5); pH in very low water (≤ 5.0); low exchangeable Ca content ($<2.0 \text{ cmolc dm}^{-3}$); low

exchangeable Mg content (≤ 0.5 cmol_c dm⁻³); high S content (> 5 mg dm⁻³); the very low available P content (≤ 7.0 mg dm⁻³); the exchangeable K content is considered to be very low (≤ 15 mg dm⁻³); content of B, Cu and Zn is considered high (> 0.3 , > 0.4 and > 0.5 mg dm⁻³ respectively), the saturation by Al is very high ($> 40\%$) and the saturation by bases is very ($< 45\%$).

In relation to the physical attributes, Lemos and Santos [16] classify the soil as sand-free surface texture and sandy loam clay texture in depth. According to Reinert and Reichert [17], the bulk density found is considered adequate for most crops (Table 1).

Table 1. Physical and chemical attributes of the soil in a *Eucalyptus benthamii* stand, at 7-year-old, in the Pampa Biome.

Depth	Density	⁽¹⁾ CS	⁽²⁾ FS	Silt	Clay	OM	pH	CTC _{efet.}	Ca
cm	g cm ³			%			(H ₂ O)	cmol _c dm ⁻³	
0-20	1.62a	75a	6 ^a	3 ^a	16a	0.7a	4.2a	2.3a	0.4b
20-40	1.57a	74a	4 ^a	5 ^a	17a	0.7a	4.3a	2.3a	0.5b
40-100	1.48a	69a	6 ^a	5 ^a	20a	0.6a	4.4a	2.7a	0.9a
Depth	Mg	P	K	S	B	Cu	Zn	V	m
cm	cmol _c dm ⁻³			mg dm ⁻³				%	
0-20	0.1b	8.3a	26.9 ^a	5.6 ^a	0.6a	3.5a	0.3a	5.7b	75.2a
20-40	0.1b	4.0b	18.4ab	6.2 ^a	0.6a	3.9a	0.1a	7.2b	70.8a
40-100	0.3a	3.9b	15.9b	4.5 ^a	0.5a	3.5a	0.1a	13.1a	54.7b

⁽¹⁾CS = Coarse Sand; ⁽²⁾FS = Fine Sand; OM = Organic Matter; Equally vertical letters do not differ statistically between the attributes in the 0-20, 20-40 and 40-100 cm layers, respectively, at the 0.05 level of significance, by the Tukey test.

Despite the low fertility observed in the soil of the study area, there is a good growth behavior of this genotype, with an annual average increment with bark of 49.87 m³ ha⁻¹ and a total production of commercial wood with bark of 349.09 m³ ha⁻¹. Among the models tested, for the estimation of height and volume, the best adjustments were provided by the Hohenald-Kreene and Schumacher-Hall models, respectively (Table 2).

Table 2. Dendrometric characteristics of *Eucalyptus benthamii* stands at 7-years-old in the Pampa Biome.

N	DBH			H			G			AAI _{wb}	V _{wb}	AAI _b	V _b
	Mean	σ	CV	Mean	σ	CV	Mean	σ	CV				
922	19.38	4.92	25.38	25.67	3.94	15.35	24.60	0.01	46.17	44.46	311.12	49.87	349.09
Model										Regression adjustment statistics			
										Prob. > F	R ² adj.	Syx(%)	
H = -22.868938 + 3.305093.DBH - 0.0577850.DBH ²										0.0001	0.91	10.51	
Log V _b = -3.634929 + 1.5414769.logDBH + 0.894537.LogH										0.0001	0.99	7.21	
Log V _{wb} = -3.688916 + 1.4279525.logDBH + 0.995249.LogH										0.0001	0.99	8.14	

N = Number of trees per hectare, DBH = diameter at breast height in cm, H = Total height in m, B = basal area in m² ha⁻¹, CV = coefficient of variation; AAI = Annual average increment m³ ha⁻¹; V = Volume m³; wb = without bark; b = with bark.

The values of DBH, H, B, AAI and V verified are similar to those found by Benin [18], studying *E. benthamii*, at 6 years of age, planted in different spacing, in Guarapuava-Paraná. However, it differed from Mendoza [19] which observed an average annual increment of 34 m³ ha⁻¹ in a settlement of *E. benthamii* with 85% survival in northern Argentina at 7 years of age.

The modeling of above-ground biomass in a clone of *Eucalyptus saligna* stand at 10 years of age was performed by Momolli *et al* [20]. The authors found 89; 5.9; 3.2 and 1.8% of the biomass in the wood, bark, branch and leaf respectively. The average annual increment of bark was 54.6 m³ ha⁻¹ and the total biomass above-ground was 269 Mg ha⁻¹. The highest percentages of wood occur due to the maturity of the stand, in addition to being a clone with genetic improvement.

Barros and Comerford [21] explain that the great variation in productivity of plantations with eucalyptus in the different regions is mainly associated to the different types of soils that have available contents and total nutrients in a very wide range. Considering this condition, Guimarães [22] adds that the forester should intervene in the management of the site and, consequently, increase the gains in production and reduce operating costs.

The models selected for biomass estimation presented good predictive capacity and significance, evidenced by the equation adjustment statistics. The biomass production above-ground was 192.0 Mg ha⁻¹, distributed in wood (81.2%)> branches (11%)> bark (6.5%)> leaves (1.3 %) (Table 3).

Table 3. Quantity of biomass and models by components of *Eucalyptus benthamii*, at 7 years of age, in the Pampa Biome.

	(1)Mg ha ⁻¹	Model	R ² adj.	Syx(%)
L	2.5	$y = -0.192074^{ns} + 0.000273411^{**}.d^2h$	0.93	20.26
Br	21.2	$y = 13.87236^{**} + 10.23822^{*}.hv - 0.00049041^{**}.d^2h^2 + 0.00691^{**}.d^3$	0.94	11.52
Ba	12.6	$y = -1.80420^{ns} + 74.59571^{**}.v - 1.162225^{*}.hv$	0.96	6.61
W	156.0	$y = -22.33094^{*} + 869.02702^{**}.v - 0.00034689.d^2h^2$	0.99	4.76
T	192.0	$y = -44.85056^{**} + 1348.85096^{**}.v - 0.00071892^{**}.d^2h^2 + 122064^{ns}.1/h^3$	0.99	3.58
C	23.7	$y = 13.96618^{**} + 10.78749^{**}.hv - 0.00050939^{**}.d^2h^2 + 0.00727^{**}.d^3$	0.97	9.13
WBa	18.3	$y = -23.88367^{*} + 939.37265^{**}.v - 0.00037597^{ns}.d^2h^2$	0.99	4.48

Were: L = leaf; Br = branch; Ba = bark; W = wood; T = total; C = canopy; WBa = wood + bark; d = diameter at breast high; h = high; v = volum (m³); R² adj. = adjusted coefficient of determination; Syx (%) = standard error of estimate; ns = not significant; * Significant at the 5% probability level of error; ** Significant at 1% probability of error.

Hernández *et al.* [23], evaluating *E. dunnii*, aged 9 years in Uruguay, verified that the biomass production of wood was 144 Mg ha⁻¹ and in the branches of 22 Mg ha⁻¹, similar to this study, but for biomass of the bark (29 Mg ha⁻¹) and leaves (13 Mg ha⁻¹) the values observed were higher. Viera *et al.* [14] studying the hybrid of *Eucalyptus urophylla* x *E. globulus*, at 10 years of age in Rio Grande do Sul, also verified a similar biomass production (167.10 Mg ha⁻¹). Guimarães *et al.* [24] studying *E. dunnii*, at the age of 4 years in the same region of this study, found an above-ground production of 104.5 Mg ha⁻¹, with 76.7 Mg ha⁻¹ for wood biomass differing from that observed in this study.

In relation to the relative partition, Schumacher *et al.* [25] evaluating *Eucalyptus* spp. Stands, with different ages, observed that at 2 years of age, 47% of the biomass was allocated to the

wood, and that at 8 years of age, the proportion of biomass in the increased to 74.4%, with the reduction of the relative biomass of the other components, corroborating with the trend found in this work. For Schumacher [26], the difference in the biomass allocation in the tree components is very dynamic due to the carbohydrate distribution, resulting from the photosynthesis, besides the edaphoclimatic factors, species and the density of planting with the age of the stand.

The highest nutrient contents of the biomass components were observed in the leaf, with the exception of Ca and Mg, with higher values in the bark ($p < 0.05$). The largest stocks of nutrients in above-ground biomass were observed in the wood, except for Ca and Mg, which were more accumulated in the bark. The highest nutrient utilization efficiency was verified in wood, which presented the highest values for the biological utilization coefficient, except for the Zn that was in the leaves component (Table 4).

Table 4. Concentrations, nutrient amounts and biological utilization coefficient (BUC) of the above-ground biomass components of *Eucalyptus benthamii*, at seven years of age in the Pampa Biome.

Biomass	N	P	K	Ca	Mg	S	B	Cu	Mn	Zn
	g kg ⁻¹			Concentration			mg kg ⁻¹			
L	31.0a	2.1a	12.7a	6.2b	2.3a	1.5a	32.0a	11.4a	1058.5a	16.8a
Br	6.5b	0.8b	4.8b	9.7b	1.9a	0.5b	10.9b	8.4ab	479.7b	22.5a
Ba	8.0b	1.3b	6.4b	19.7a	3.6a	0.6b	15.5b	5.1b	825.6ab	22.8a
W	1.9c	0.4c	1.5c	0.8c	0.1b	0.3b	4.5c	2.2c	77.2c	29.8a
	kg ha ⁻¹ Amount					g ha ⁻¹				
L	77.5 (12.3)	5.0 (5.0)	32.1 (6.7)	12.3 (2.2)	5.0 (4.6)	3.6 (5.1)	80.5 (6.7)	27.1 (4.5)	2256.0 (6.4)	39.5 (0.6)
Br	137.0 (21.8)	16.7 (16.6)	106.9 (22.4)	172.5 (31.0)	37.0 (33.9)	9.2 (13.2)	237.5 (19.8)	175.4 (29.2)	9385.3 (26.6)	440.0 (7.1)
Ba	82.3 (13.1)	15.1 (15.0)	80.0 (16.8)	248.1 (44.5)	41.4 (37.9)	6.0 (8.5)	164.0 (13.7)	54.6 (9.1)	10825.1 (30.7)	222.7 (3.6)
W	332.7 (52.8)	63.6 (63.4)	257.6 (54.1)	124.1 (22.3)	25.9 (23.6)	51.4 (73.2)	692.8 (59.8)	374.6 (57.2)	12798.3 (36.3)	5462.0 (88.7)
	BUC									
L	32	498	78	204	507	706	31181	92765	1113	63510
Br	155	1265	198	123	573	2297	89163	120765	2257	48131
Ba	152	831	157	51	303	2088	76543	229874	1159	56359
W	468	2450	605	1256	6014	3030	224817	415795	12170	28517

Were: L = leaf; Br = branch; Ba = bark; W = wood. Vertical letters do not differ statistically between the nutrient contents in the biomass components, at the 0.05 level of significance, by the Tukey test. Values in parentheses refer to the relative partition (%) of each nutrient per component in relation to the total quantity.

The magnitude of nutrient concentration, in descending order, was as follows: leaves > bark > branches > wood. The higher levels of nutrients observed in the leaf, as well as the difference in concentration between the components of biomass, can be explained by the tendency of most nutrients to concentrate in the new structures of the plant, where the main

metabolic processes occur. The interaction of these processes is intrinsically related to biochemical cycling, where with age, nutrients from senescent tissues tend to move to regions with higher metabolic activity, and biochemical cycling is more important for the maintenance of nutrients with high mobility (N, P, K and Mg), and lower for low mobility nutrients (Ca, S) and micronutrients [27,28,29].

Considering the nutrient utilization efficiency of wood, is verified that Mg is the macronutrient least required for the production of biomass, followed by S, P, Ca, K and N. Several authors have observed a higher conversion of nutrients to wood than those observed in this study. Silva et al. [30] evaluating 5 eucalyptus species, at 10 years of age, in Itirapina-SP, verified that P provided the highest biological utilization coefficient (BUC) of nutrients, especially *E. grandis* with 43441 for P in wood, but for Mg efficiency was lower among all genotypes. This behavior was also verified by Viera et al. [14] studying the hybrid *Eucalyptus urophylla* x *E. globulus*, with a BUC of 13285 for P in the wood, followed by S, Mg, Ca, K and N; and by Guimarães[22] studying the clonal hybrid *Eucalyptus urophylla* x *E. grandis*, *E. grandis* and *E. dunnii*, at the age of four in Alegrete-RS, who verified in the *E. urograndis* a BUC of 17060 of P in the wood, followed by Mg, Ca, S, K and N. In another study carried out with 8 clonal *E. urograndis* hybrids, at 9 years of age in Aracruz-ES, Neves [31] verified that S was the nutrient that presented the highest conversion in wood (average BUC of 8500), followed by P, Mg, K, N and Ca; which was also observed by Beulch [32], studying clone *E. saligna*, at the age of four in São Francisco de Assis-RS, who verified a BUC of 11688 for S in wood, followed by P, Mg, Ca, S, N and K.

Silva et al. [30] argue that the high efficiency presented by a species in the use of nutrients conditions a lower nutritional requirement, which can be used as an indicator for the selection of species that can be cultivated mainly in soils with low natural fertility. In addition, Santana et al. [33] complement that the use of genetic material that has efficiency compatible with soil fertility, can maintain the productive capacity of the site and, consequently, use smaller amounts of fertilizers.

In this study, there is a great potential for growth and production of the *Eucalyptus benthamii* genotype. The species showed to be less efficient in nutrient utilization, when compared to several seminal and clonal materials, cultivated on a large scale in Brazil. However, considering its good adaptability to cold climate regions such as the Pampa Biome in the state of Rio Grande do Sul, studies on genetic improvement through hybridization with species that present a better efficiency of nutrient utilization in this region and later cloning of the best individuals for commercial cultivation.

4. CONCLUSION

Despite the low natural fertility of the soil in the experimental area, *Eucalyptus benthamii* presented productivity similar to the other eucalyptus species cultivated on a commercial scale in Brazil. The above-ground biomass is predominantly allocated to the wood (81.2%), followed by branches (11%), bark (6.5%) and leaves (1.3%).

The leaves present the highest levels of nutrients, except Ca and Mg (bark) and Zn (wood). The largest amounts of nutrients are allocated to the wood, except for Ca and Mg (bark). The wood presents the highest efficiency of nutrient utilization, with the exception of Zn.

285 **COMPETING INTERESTS**

286

287 Authors have declared that no competing interests exist.

288

289

290 **REFERENCES**

291

292 1. Hall N, Brooker I. Camden White Gum: *Eucalyptus benthamii* Maiden et Cambage.
293 Camberra: Department of National Development Forestry and Timber Bureau, 1973;4.
294 (Forest Tree Series, 57).

295 2. Butcher PA, Skinner AK, Gardiner CA. Increased inbreeding and inter-species gene flow
296 in remnant populations of the rare *Eucalyptus benthamii*. Conservation Genetics.
297 2005;6(2):213-226.

298 3. Graça MEC, Shimizu JY, Tavares FR. Capacidade de rebrota e de enraizamento de
299 *Eucalyptus benthamii*. Boletim de Pesquisa Florestal. 1999;39:135-138. Portuguese.

300 4. Higa RCV, Pereira JCD. Usos potenciais do *Eucalyptus benthamii* Maiden et Cambage.
301 Colombo: Embrapa Florestas. 2003;4. (Comunicado Técnico, 100). Portuguese.

302 5. Viera M, Schumacher MV, Caldeira MVW. Biomassa e exportação de nutrientes pela
303 colheita do eucalipto. In: Viera M, Schumacher MV. Silvicultura do eucalipto no Brasil. Santa
304 Maria: UFSM, 2015a: 245-272. Portuguese.

305 6. Achat DL, Deleuze C, Landmann G, Pousse N, Ranger J, Augusto L. Quantifying
306 consequences of removing harvesting residues on forest soils and tree growth – A meta-
307 analysis. Forest Ecology and Management. 2015;348:124–141.
308 doi:10.1016/j.foreco.2015.03.042

309 7. Alvares CA, Stape JL, Sentelhas PC, Gonçalves JLM, Sparovek G. Köppen's climate
310 classification map for Brazil. Meteorologische Zeitschrift. 2014;22(6):711-728.

311 8. EMBRAPA. Sistema Brasileiro de Classificação de Solos – SBCS. 3 ed. Brasília, DF.
312 2013. 353 p. Portuguese.

313 9. Finger CAG. Fundamentos de Biometria Florestal. Santa Maria: SM/FATEC/CEPEF,
314 1992. 269p. Portuguese.

315 10. Empresa Brasileira De Pesquisa Agropecuária. Centro Nacional de Pesquisa de Solos.
316 Manual de Métodos de Análise de Solos. 2. ed. Rio de Janeiro, 1997. 212p. Portuguese.

317 11. Tedesco MJ, Gianello C, Bissani CA, Bohnen H, Volkweiss SJ. Análise de solo, plantas
318 e outros materiais. 2.ed. Porto Alegre: Departamento de Solos, UFRGS, 1995. 174p.
319 (Boletim Técnico, 5). Portuguese.

320 12. Miyazawa M, Pavan MA, Muraoka T. Análises químicas de tecido vegetal. In: SILVA,
321 F.C. (Org.). Manual de análises químicas de solos, plantas e fertilizantes. Brasília: Embrapa
322 Comunicação para Transferência de Tecnologia, 1999. cap. 4, p.171-224. Portuguese.

- 323 13. SAS. Statistical analysis system: Programa de computador, ambiente VM. Cary, 2003.
324 Versão 6.08.
- 325 14. Viera M, Schumacher MV, Trüby P, Araújo FE. Implicações nutricionais com base em
326 diferentes intensidades de colheita da biomassa de *Eucalyptus urophylla* x
327 *Eucalyptus globulus*. Ciência Rural. 2015b;45(3):432-439. Portuguese.
- 328 15. SBCS-CQFS – Sociedade Brasileira de Ciência do Solo-Comissão de Química e
329 Fertilidade do Solo – RS/SC. Manual de calagem e adubação para os Estados do Rio
330 Grande do Sul e de Santa Catarina. 11ª ed. Solo – Núcleo Regional Sul. Porto Alegre. 2016,
331 376 p. Portuguese.
- 332 16. Lemos RC, Santos RD. Manual de descrição e coleta de solo no campo. Sociedade
333 Brasileira de Ciência do Solo – Centro Nacional de Pesquisa de Solos. 3ª Ed., p. 83,
334 Campinas-SP, 1996. Portuguese.
- 335 17. Reinert DJ, Reichert JM. Propriedades físicas de solos em sistema plantio direto
336 irrigado. In: Carlesso R, Petry M, Rosa G, Ceretta, CA. Irrigação por Aspersão no Rio
337 Grande do Sul, Santa Maria. 2001. p. 114-131. Portuguese.
- 338 18. Benin CC. Efeito do espaçamento na produção, variáveis dendrométricas e
339 Propriedades da madeira de *Eucalyptus benthamii*. 2014. 58 p. Dissertação (Mestrado em
340 Manejo Sustentável de Recursos Florestais) – Universidade Estadual do Centro-Oeste, Irati,
341 2014. Portuguese.
- 342 19. Mendoza L. Notes on *Eucalyptus benthamii* in Argentina. In: COLLOQUES
343 INTERNATIONAL SUR LES EUCALYPTUS RESISTANTS AU FROID, 1983, Bordeaux.
344 Annales... Bordeaux: IUFRO, 1983. p.480.
- 345 20. Momolli DR. et al. Modeling and biomass quantification in *Eucalyptus saligna* Smith
346 stand at the end rotation in the south of Brazil. Journal of Experimental Agriculture
347 International. 2019; 33(3), 1-10. <https://doi.org/10.9734/jeai/2019/v33i330146>.
- 348 21. Barros NF, Comerford NB. Sustentabilidade da produção de florestas plantadas na
349 região tropical. In: Alvarez VVH et al. eds. Tópicos em ciência do solo. Viçosa, Sociedade
350 Brasileira de Ciência do Solo, Viçosa, Folha de Viçosa, 2002. v.2. p.487-592. Portuguese.
- 351 22. Guimarães, C. C. Biomassa e Nutrientes em plantios de Eucaliptos no Bioma Pampa.
352 2014. 63f. Dissertação (Mestrado em Ciências Florestais). Universidade Federal de Santa
353 Maria, Santa Maria, 2014. Portuguese.
- 354 23. Hernández J, Pino A, Salvo L, Arrarte G. Nutrient export and harvest residue
355 decomposition patterns of a *Eucalyptus dunnii* Maiden plantation in temperate climate of
356 Uruguay. Forest Ecology and Management. 2009;258(2):92-99.
- 357 24. Guimarães CC, Schumacher MV, Witshoreck R, Souza HP, Santos JC, Vieira FCB.
358 Biomassa e nutrientes em povoamento de *Eucalyptus dunnii* Maiden no Pampa Gaúcho.
359 Revista Árvore. 2015;39(5):873-882. Portuguese.
- 360 25. Schumacher MV, Witschoreck R, Calil FN. Biomassa em povoamentos de
361 *Eucalyptus* spp. de pequenas propriedades rurais em Vera Cruz - RS. Ciência Florestal.
362 2011;21(1):17-22. Portuguese.

- 363 26. Schumacher MV. Ciclagem de nutrientes como base da produção sustentada em
364 ecossistemas florestais. In: SIMPÓSIO SOBRE ECOSSISTEMAS NATURAIS DO
365 MERCOSUL O AMBIENTE DA FLORESTA, 1., 1996, Santa Maria. Anais... Santa Maria:
366 UFSM/CEPEF, 1996, p.65-77.Portuguese.
- 367 27. Poggiani F, Schumacher MV. Nutrient cycling in native forest. In: Gonçalves JLM,
368 Benedetti V. Forest nutrition and fertilization. Piracicaba: IPEF, 2004. p. 285-
369 305.Portuguese.
- 370 28. Pallardy S. Physiology of woody plants. San Diego: Academic Press, 2008. 454p.
- 371 29. Viera M. Dinâmica nutricional em um povoamento híbrido de *Eucalyptus urophylla* x
372 *Eucalyptus globulus* em Eldorado do Sul-RS, Brasil. 2012. 119 p. Tese (Doutorado em
373 Engenharia Florestal) – Universidade Federal de Santa Maria, Santa Maria, 2012.
374 Portuguese.
- 375 30. Silva HD, Poggiani F, Coelho LBC. Eficiência de utilização de nutrientes em cinco
376 espécies de *Eucalyptus*. Colombo: Embrapa Florestas, 1983. 8 p. (Embrapa Florestas,
377 Boletim de Pesquisa Florestal, n. 6/7).Portuguese.
- 378 31. Neves JCL. Produção e partição de biomassa, aspectos nutricionais e hídricos em
379 plantios clonais de eucalipto na região litorânea do Espírito Santo. 2000. 191f. Tese
380 (Doutorado em Produção Vegetal) – Universidade Estadual do Norte Fluminense, Campos
381 dos Goytacazes, RJ, 2000.Portuguese.
- 382 32. Beulch LS. Biomassa e nutrientes em um povoamento de *Eucalyptus saligna* Smith
383 submetido ao primeiro desbaste.2013. 58 p. Dissertação (Mestrado em Engenharia
384 Florestal) – Universidade Federal de Santa Maria, Santa Maria, 2013.Portuguese.
- 385 33. Santana RC. et al. Biomassa e conteúdo de nutrientes de procedências de
386 *Eucalyptus grandis* e *Eucalyptus saligna* em alguns sítios florestais do Estado de São Paulo.
387 Scientia Forestalis. 1999;56:155-169.Portuguese.
- 388