

1 DIFFERENT LIGHT RADIATION INTENSITIES ON COTTON: A 2 PHYSIOLOGICAL APPROACH

3
4 **Abstract:** The luminosity and the temperature are factors that act directly in the
5 photosynthetic process, where the elevation of the luminous intensity can cause
6 a reduction in the assimilation of carbon, which consequently lowers the
7 development of the cotton. The objective of this work was to know the response
8 of physiological parameters of cotton when subjected to different artificial light
9 intensities. A randomized complete block design was used in a 2x5 factorial
10 scheme, with two varieties of cotton: IMA5801B2RF e IACRDN, interacting with
11 five artificial light intensities, being interacting with 5 densities of light: 0
12 (control); 500; 1000; 1500 and 2000 $\mu\text{mol m}^{-2} \text{s}^{-1}$ of photosynthetically active
13 radiation (PAR) provided by LED bulbs. The following variables were set: rate of
14 CO_2 assimilation (A); transpiration (E); stomatal conductance (GS); inner CO_2
15 concentration in the substomatic chamber (Ci) and efficient use of water (EUW)
16 in which a portable device of gas exchange was used (Infra-Red Gas Analyzer -
17 IRGA, marca ADC BioScientific Ltd, modelo LC-Pro). The cotton varieties
18 responded positively under different luminous intensities until reaching the point
19 of maximum saturation between 1400 a 1600 $\mu\text{mol m}^{-1} \text{s}^{-1}$ of light, which
20 provides a better rate of CO_2 assimilation (A); concentration of CO_2 in the
21 substomatic chamber (Ci) and efficient use of water (EUW). Leaf transpiration
22 (E) and stomatal conductance of the cotton showed a positive linear response
23 with increasing light intensity. The ideal luminous intensity for the use of Infra-
24 Red Gas Analyzer - IRGA was 1500 $\mu\text{mol m}^{-1} \text{s}^{-1}$ in the tested cotton crop.

25
26 **Keywords:** *Gossypium* L .; brightness; CO_2 assimilation; photosynthesis rate

27 28 Introduction

29 Belonging to the Malvaceae family, cotton (*Gossypium* L.) is cultivated as
30 a fiber source for the production of fabrics, as well as crushing of its seeds for
31 the production of linoleic and linolenic oils that are used in the cosmetics or
32 animal feed industry. Because it is an important crop for the Brazilian
33 agricultural scenario, since it makes an alternative in crop rotation in the
34 production of large crops such as corn and soybeans, cotton can suffer

interference during its developmental stages due to climatic factors such as water stresses, pests and diseases and light intensity, especially in the establishment and reproduction phases [1].

Light is the primary source of energy related to photosynthesis and morphogenetic phenomena, and is one of the main factors that influence plant growth and development [2; 3; 4]. Increase in light intensity can reduce the photosynthetic activity through photoinhibition, and this response can be variable between plant species and varieties [5; 6]. The luminous intensity and the temperature are factors that can cause the limitation of the photosynthetic process and also contribute to the reduction of the carbon acquisition, consequently causes a reduction in rate of plant growth [7].

The plants when subjected to medium intensity light show less transpiration when compared to plants that are exposed to more intense light intensity, that is, less light is a limiting factor for leaf transpiration [8]. The importance of light intensity in the physiological process of the plant, is evidenced in its direct link in the activation of enzymes related to carbon fixation and in the control in the opening and closing in the stomatal cleft [9; 10; 11].

It is important to emphasize that the understanding in the balance of intensity levels and the duration of exposure to light that plants can be subjected to makes it an important factor to know the responses of plants to varying light stress. When exposed to direct low-intensity radiation, the plants become more efficient in carrying out their photosynthesis, since the process is started in a gradual way, which does not compromise the pathways of the electrons by the photosystems. But with the increase of this intensity of photons that affect the leaves, the plants present an elevation in the photolysis of the water, which results in a saturation of electrons, causing a reduction in the rate of assimilation of CO₂ and in the efficient use of water [12; 13].

In view of the above, this work had as objective to know the response of selected physiological parameters of different intensities of light radiation on cotton crop.

Material e Methods

The experiment was carried out in December 2018, at the Paulista Agribusiness Technology Agency (APTA), located in the city of Adamantina, State of São Paulo, with geographic coordinates 21°40'24.024"S and 51°8'31.088"W, with an altitude of approximately 420 m. The climate of the region is characterized as Aw according to Köppen, with rainy summers and dry winters; with an annual average temperature of 22.1°C and 1204 mm of rain accumulated in the year.

The experiment was carried out in randomized blocks, in a factorial scheme of 2x5, being 2 varieties of cotton, IMA5801B2RF and IAC-RDN, interacting with 5 densities of light: 0 (control); 500; 1000; 1500 e 2000 $\mu\text{mol m}^{-2} \text{s}^{-1}$ of photosynthetically active radiation (PAR) provided by LED bulbs.

The area soil was classified as Red-yellow Latosol [14] and presented the following chemical attributes (Table 1).

Table 1: Chemical attributes of the soil of the experiment area at the time of sowing of cotton.

pH	OM	P	K	Ca	Mg	H+Al	Al	SB	CEC	V%	m%
CaCl ₂	g dm ⁻³	mg dm ⁻³						mmol _c dm ⁻³			
4.6	12.0	26.0	2.9	8.0	4.0	20.0	1.0	14.9	34.9	43.0	6.0

SB: Sum of bases; V%: Saturation per bases; m%: Saturation per aluminum; CEC: Cation exchange capacity.

Each block consisted of five rows of five meters in length, spaced 0.9 m between rows, where the cotton was sown with a population intensity of 45 thousand plants per hectare. Soil was fertilized as the needs of cotton culture [15]. During the experiment, the cotton was watered until the soil reached field capacity, and the phytosanitary treatments of the crop were done.

Thirty days after the sowing, five plants were randomly selected within each replicate, where four readings were performed on the leaves fully expanded from the apex of the plant, totaling 20 readings for each light intensity for the different cotton varieties. The following parameter were measured: rate of CO₂ assimilation ($A - \mu\text{mol CO}_2 \text{ m}^{-2} \text{s}^{-1}$); transpiration ($E - \text{mmol H}_2\text{O m}^{-2} \text{s}^{-1}$); stomata conductance ($GS - \text{mol H}_2\text{O m}^{-2} \text{s}^{-1}$); inner CO₂ concentration in the substomatic chamber ($C_i - \mu\text{mol mol}^{-1}$), with 380 ppm of CO₂, under 28°C temperature of chamber, a portable device of gas exchange was used (Infra-

96 Red Gas Analyzer - IRGA, ADC BioScientific Ltd, modelo LC-Pro); and efficient
 97 use of water (EUW) by applying the following mathematical formula:
 98

$$EUW = \frac{A}{E}$$

99

100 All variables were **subjected** to the F test ($p < 0.05$) and analyses of
 101 regression were applied to the intensities of artificial light, in which their
 102 standards were tested: linear, quadratic and cubic. Cotton varieties were
 103 **subjected** to Scott & Knott Test, at 5% probability [16]. Statistic program R was
 104 used [17].

105

106 Results

107 IMA5801B2RF showed higher mean values for CO_2 assimilation (A) and
 108 water efficiency (EUW), with a difference of 4.68% and 5.79%, respectively, in
 109 relation to the IAC-RDN variety (Table 2).

110

Table 2: Mean values of rate of CO_2 assimilation ($A - \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$); transpiration ($E - \text{mmol H}_2\text{O m}^{-2} \text{ s}^{-1}$); stomata conductance ($GS - \text{mol H}_2\text{O m}^{-2} \text{ s}^{-1}$); inner CO_2 concentration in the substomatic chamber ($C_i - \mu\text{mol mol}^{-1}$) and the efficient use of water (EUW - $\text{mol CO}_2 \text{ mol H}_2\text{O}^{-1}$) and analysis of variance of the cotton regressions when exposed to different intensities of light radiation, where the models were tested: linear, quadratic and cubic.

Variety (V)	A	E	GS	Ci	EUW		
IMA5801B2RF	16.66a	3.48	0.53	275.63b	4.49a		
IAC-RDN	15.88b	3.45	0.53	282.24a	4.23b		
CV%	12.07	11.54	26.20	6.21	14.38		
OM	16.27	3.46	0.53	278.93	4.36		
Variety (V) of F	8.04**	0.28Ns	0.06Ns	7.28**	8.47**		
Radiation (R) of F	1320.22**	84.09**	42.27**	639.20**	954.15**		
VxR of F	5.36**	0.76Ns	1.73Ns	4.06**	4.36**		
	VF	DF	Regressions middle square				
IMA5801B2RF	Radiation	4	2394.8730	26.7034	1.7038	137064.375	264.5401
	Residue	96	2.8655	0.1641	0.0201	264.7738	0.3051
	Regression	1	Q**	L**	L**	Q**	Q**
IAC-RDN	Radiation	4	1628.0518	24.4440	1.5699	88253.0035	188.8543
	Residue	96	5.7066	0.1602	0.0192	378.7123	0.5611
	Regression	1	Q**	L**	L**	Q**	Q**

CV: Coefficient of variation. OM: Overall mean. F: value of F calculated in the analysis of variance; Ns $p = 0.05$; * $0.01 \leq p < 0.05$; ** $p < 0.01$. The averages in the column followed by the same letter do not differ statistically from each other.

The Scott&Knott test was applied at a 5% probability level. Ns- $p \geq 0.05$;

*0.01=<p<0.05; ** p <0.01. VF: Variation factor; DF: Degrees of freedom. L: polynomial of 1st degree. Q: polynomial of 2nd degree.

111

112 There was no difference between the varieties in the transpiration (E)
113 and stomatal conductance (GS) parameters when the cotton was exposed to
114 different light intensities (Table 2). However, the IAC-RDN variety showed a
115 greater mean in the internal CO₂ concentration in the substamatic chamber (Ci)
116 with a difference of 2.34% more in relation to IMA5801B2RF.

117 When the light intensities are taken into account, the varieties responded
118 in a similar way in all parameters evaluated (Table 2). The varieties presented a
119 positive quadratic response to the CO₂ assimilation rate, (Figure 1), where the
120 IMA5801B2RF variety presented a maximum point up to 1521 $\mu\text{mol m}^{-2} \text{s}^{-1}$
121 while the IAC-RDN variety had a maximum point of 1673 $\mu\text{mol m}^{-2} \text{s}^{-1}$.

122




Fig 1. CO₂ assimilation rate (A) of the cotton when exposed to different intensities of light radiation.

(1) IMA5801B2RF e (2) IAC-RDN.

123

124 While there was an increase in light intensity, the cotton varieties
125 presented a positive linear response to the transpiration parameter of the leaf
126 (E) as shown in Figure 2, already in Figure 1.

127




Fig. 2. Transpiration of cotton leaf (E) when exposed to different intensities of light radiation.

(1) IMA5801B2RF e (2) IAC-RDN.

128

129 Similarly, the varieties exhibited a positive response to the increase in
 130 light intensity on leaf perspiration (E) (Figure 3). Again, the understanding of
 131 these responses regarding leaf water loss with increase in the luminous
 132 intensity is important in the determination of the point of maximum response of
 133 this variable. This becomes an important tool in the decision making in the
 134 cotton cultivation, since it can guarantee a better understanding of the water
 135 availability requirements.

136




Fig. 3. Stomatal conductance (GS) of cotton when exposed to different intensities of light radiation.

(1) IMA5801B2RF e (2) IAC-RDN.

137

138 In contrast, the internal CO₂ concentration in the sub-static chamber (Ci)
 139 of the cotton cultivars presented negative quadratic responses when there was
 140 an increase in light intensity, where a minimum point of 1385 $\mu\text{mol m}^{-2} \text{s}^{-1}$ was
 141 observed in the variety IMA5801B2RF and 1528 $\mu\text{mol m}^{-2} \text{s}^{-1}$ for the IAC-RDN
 142 variety, as shown in Figure 4.
 143




Fig. 4. Internal CO₂ concentration in the substomatic chamber (Ci) of the cotton when exposed to different intensities of light radiation.

(1) IMA5801B2RF e (2) IAC-RDN.

144
 145 With the increase in the intensity of the light radiation on the leaves, the
 146 cotton varieties presented a quadratic positive response in the parameter EW
 147 (water efficient use) (Figure 5), where the maximum points of 1375 $\mu\text{mol m}^{-2} \text{s}^{-1}$
 148 in the variety IMA5801B2RF and 1489 $\mu\text{mol m}^{-2} \text{s}^{-1}$ in the IAC-RDN. This shows
 149 that the light intensity influences positively only until its saturation as pointed out
 150 earlier. This saturation of light causes an increase in the photolysis of the water
 151 which may have led to the saturation of electrons in the photosystem.
 152




Figure 5: Uso eficiente da água (EUW) do algodoeiro quando exposto em diferentes intensidades de radiação luminosa.

(1) IMA5801B2RF e (2) IAC-RDN.

153

154 Negative correlations were observed between the internal CO₂
 155 concentration variable in the **substomatic** chamber (Ci) interacting with leaf
 156 transpiration (E); stomatal conductance (GS); rate of assimilation of CO₂ (A)
 157 and water efficiency (EUW) as shown in Table 3.

158

Table 3: Pearson correlation coefficient *r* values among the analyzed variables of cotton when submitted to different light intensity.

	Ci	E	GS	A
E	-0.5733** < 0.0001	---		
GS	-0.3943** < 0.0001	0.94156** < 0.0001	---	
A	-0.9432** < 0.0001	0.79094** < 0.0001	0.64496** < 0.0001	---
EUW	-0.9955** < 0.0001	0.61386** < 0.0001	0.44222** < 0.0001	0.96071** < 0.0001

*Ns-p>=0.05; *0.01=<p<0.05; **p < 0.01. rate of CO₂ assimilation (A – μmol CO₂ m⁻² s⁻¹); transpiration (E – mmol H₂O m⁻² s⁻¹); stomata conductance (GS – mol H₂O m⁻² s⁻¹); inner CO₂ concentration in the substomatic chamber (Ci – μmol mol⁻¹) and the efficient use of water (EUW - mol CO₂ mol H₂O⁻¹).*

159

160 However, positive correlations were observed between the variable leaf
 161 transpiration (E) interacting with the stomatal conductance (GS); rate of
 162 assimilation of CO₂ (A) and efficient use of water (EUW). In the same way,
 163 stomatal conductance (GS) presented a positive correlation with CO₂
 164 assimilation rate (A) and water efficiency (EUW) and, finally, the rate of

assimilation of CO₂ (A) with the efficient use of water (EUW) showed a positive correlation as shown in Table 3.

DISCUSSION

The plant can respond in different ways with the change of the environment in which it was inserted, where the luminosity is restrictive to the development of the plant, since the quality and the luminous intensities that affect the leaves alter the responses in the PSII and PSI complexes of the photosystem. This can cause changes in the photolysis of the water, which consequently in the release of electrons during photosynthesis due to the increase or restriction of the photons that are affecting the plant [13], in this way, the ideal intensity observed is approximately 1500 $\mu\text{mol m}^{-1} \text{s}^{-1}$ light falling on the leaves of the cotton plant.

It is worth noting that, even at different periods of the day, a variation occurs in the incidence of light energy, which influences the CO₂ assimilation rate of the leaves [18] demonstrate this phenomenon that occurs naturally during the day.

The significant negative correlation between the internal concentration of CO₂ in the sub-static chamber (Ci) and the other variables as shown in Table 4 was already expected, since the internal concentration of CO₂ is reduced while the carbon fixation in the dry matter of the cotton occurs via Rubisco molecule, which results in the elevation of the CO₂ assimilation rate (A). In this way, this interaction can be verified when one observes Figure 1 and Table 2, where the absence of light on the leaves caused a negative assimilation rate (A), while the internal CO₂ concentration was high (Figure 4). And with the increase in light radiation, the stomata were opened, consequently causing an increase in the transpiration rate (E) (Figure 2) and the stomatal conductance (GS) (Figure 3) and thus led to a reduction in concentration (Ci) due to a possible dilution effect, where CO₂ at high internal concentrations is released to the environment due to the stomatal opening and its fixation in the dry mass [13].

It is worth mentioning that the understanding of the mechanism of opening and closing the stomatal cleft can be compromised or enhanced with nutritional stress factors (Table 1), and the availability of H₂O in the soil-plant-atmosphere system [10; 11] and even internal morphology of the leaves of each

species and varieties [3; 4; 5; 6]. As previously mentioned, stomatal conductance presents a positive correlation with the other variables (Table 3).

The positive correlation between the CO₂ assimilation rate (A) interacting with the use of leaf transpiration (E) was already expected, since the relationship between these two variables yields the efficient use of water (EUW), which was elevated with the increase of light radiation between 1300 and 1500 $\mu\text{mol m}^{-2} \text{s}^{-1}$ (Figure 5). When gas exchange occurs through the stomatal cleft, the plant needs a hydrostatic pressure (Kleaf) to efficiently use water (EUW) in the photosynthetic system, where water stress directly influences development in different plant species in the initial phase [5; 11; 7]. Thus, more in-depth studies are needed on the relationship between these variables, since species and varieties present different responses between them.

CONCLUSIONS

The two cotton varieties responded positively under different light intensities up to the maximum saturation point between 1400 and 1600 $\mu\text{mol m}^{-1} \text{s}^{-1}$ of light. This provided a better rate of assimilation of CO₂ (A); concentration of CO₂ in the substamatic chamber (Ci) and efficient use of water (EUW).

Leaf transpiration (E) and stomatal conductance of the cotton showed a positive linear response with increasing light intensity.

The ideal luminous intensity for the use of Infra-Red Gas Analyzer - IRGA was 1500 $\mu\text{mol m}^{-1} \text{s}^{-1}$ in the cotton crop.

COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

233

234 REFERENCES

- 235 1. Echer FR, Zanfolin PRL, Moreira ACM, Santos ACP, Gorni PH. Root
236 growth and carbohydrate partitioning in cotton subjected to shading in
237 the initial phase. *Ciência Rural*. 2016; 49(1):1-8.
238 <http://dx.doi.org/10.1590/0103-8478cr20180749>
- 239 2. Holt JS. Plant response to light: a potencial tool for weed
240 management. *Weed Science*. 1995; 43: 474-482.
- 241 3. Stewart, JJ, Polutchko SK, Adams Iii, WW, Cohu CM, Coleman A, Wenzl
242 CA, Demmig-Adams B. Light, temperature and tocopherol status
243 influence foliar vascular anatomy and leaf function in *Arabidopsis*
244 *thaliana*. *Physiologia Plantarum*. 2017; 160(1):98-110.
245 <http://dx.doi.org/10.1111/ppl.12543>
- 246 4. Feldman AB, Leung H, Baraoidan M, Elmido-Mabilangan A, Canicosa I,
247 Quick WP, Sheehy J, Murchie EH. Increasing leaf vein density via
248 mutagenesis in rice results in an enhanced rate of photosynthesis,
249 smaller cell sizes and can reduce interveinal mesophyll cell
250 number. *Frontiers In Plant Science*. 2017; 8:1-10.
251 <http://dx.doi.org/10.3389/fpls.2017.01883>
- 252 5. Xiong D, Douthe C, Flexas J. Differential coordination of stomatal
253 conductance, mesophyll conductance, and leaf hydraulic conductance in
254 response to changing light across species. *Plant, Cell & Environment*.
255 2018; 41(2):436-450. <http://dx.doi.org/10.1111/pce.13111>
- 256 6. Rockwell FE, Holbrook NM. Leaf Hydraulic Architecture and Stomatal
257 Conductance: A Functional Perspective. *Plant Physiology*. 2017;
258 174(4):1996-2007. <http://dx.doi.org/10.1104/pp.17.00303>
- 259 7. Araújo SAC, Deminiciis BB. Photoinhibition of the
260 Photosynthesis. *Brazilian Journal of Biosciences*. 2006; 7(4): 463-472. In
261 Portuguese
- 262 8. Vieira TO, Degli-Esposti MSO, Souza GM, Rabelo GR, Vitória AP.
263 Photoacclimation capacity in seedling and sapling of *Siparuna*
264 *guianensis* (Siparunaceae): Response to irradiance gradient in tropical
265 forest. *Photosynthetica*. 2015; 53(1):11-22.
266 <http://dx.doi.org/10.1007/s11099-015-0073-x>

- 267 9. Teixeira MC, Vieira TO, Almeida TCM, Vitória AP. Photoinhibition in
268 Atlantic Forest native species: short-term acclimative responses to high
269 irradiance. Theoretical And Experimental Plant Physiology. 2015; 27(3-
270 4):183-189. <http://dx.doi.org/10.1007/s40626-015-0043-5>
- 271 10. Bellasio C, Quirk J, Buckley TN, Beerling DJ. A dynamic hydro-
272 mechanical and biochemical model of stomatal conductance for C4
273 photosynthesis. Plant Physiology. 2017; 175(1):104-119.
274 <http://dx.doi.org/10.1104/pp.17.00666>
- 275 11. Li Y, Li H, Li Y, Zhang S. Improving water-use efficiency by decreasing
276 stomatal conductance and transpiration rate to maintain higher ear
277 photosynthetic rate in drought-resistant wheat. The Crop Journal. 2017;
278 5(3):231-239. <http://dx.doi.org/10.1016/j.cj.2017.01.001>
- 279 12. Atroch EMAC, Soares AM, Alvarenga AA, Castro EM. Growth,
280 chlorophyll content, biomass distribution and anatomical characteristics
281 of young plants of *Bauhinia forficata* link submitted to shading. Ciência e
282 Agrotecnologia. 2001; 25(4):853-862. In Portuguese
- 283 13. Taiz L, E Zeiger. Fisiologia vegetal. 5. ed. Porto Alegre: Artmed. 2013;
284 918p.
- 285 14. Embrapa – Empresa Brasileira de Pesquisa Agropecuária. Sistema
286 brasileiro de classificação de solos. 3.ed. Brasília. 2013; 353p.
- 287 15. Raij B, Cantarella H, Quaggio JÁ, Furlani AMC. Recomendações de
288 adubação e calagem para o Estado de São Paulo. 2.ed. Campinas: IAC.
289 1996; 285p.
- 290 16. Banzatto DA, Kronka SN. Experimentação Agrícola. 4.ed. Funep. 2013;
291 237p.
- 292 17. R Development Core Team. R: A language and environment for
293 statistical computing. R Foundation for Statistical Computing, Vienna,
294 Austria. 2009. ISBN 3-900051-07-0, URL <http://www.R-project.org>
- 295 18. Kim S, Nusinow DA, Sorkin ML, Pruneda-Paz J, Wang X. Interaction and
296 regulation between lipid mediator phosphatidic acid and circadian clock
297 regulators in Arabidopsis. The Plant Cell. 2019; 1-58.
298 <http://dx.doi.org/10.1105/tpc.18.00675>