

1

2 **Original Research Article**

3 **Biodiversity in Forest Fragments Under**

4 **Different Forms of Environmental**

5 **Conservation, Jaboticabal, São Paulo, Brazil**

6

7

8

9

10 **ABSTRACT**

11

The objective of this work was to evaluate the biodiversity of forest fragments at different levels of anthropization in the municipality of Jaboticabal, State of São Paulo, Brazil. The related research was carried out in three forest fragments, an area under reforested conditions, a natural remnant area and an area under natural regeneration. Each fragment was divided into three sampling units of 240 m², constituting the replicates. The total frequency of insects, amphibians, birds, mammals, reptiles and trees was evaluated. Cluster analysis indicated that there were differences between the three fragment types. The analyzed variables were summarized in two main components, which explain 87.1% of the accumulated total variance. It was concluded that the areas of natural remnants and reforestation presented higher frequencies of species and, consequently, greater biological diversity detriment to the natural regeneration area.

12

13 *Keywords: Environment; ecology; fauna; flora; frequency of species.*

14

15 **1. INTRODUCTION**

16

17 In the current situation the environment makes necessary the sum of efforts to better understand the dynamic interplay between man and the environment. The growing environmental awareness and the expansion of knowledge between the different knowledge areas have mobilized the scientific community and the population in favor of getting to know these interaction man/environment [1] and design strategies for the sustainable use of the environment by man.

23

24 Human exploratory actions have promoted significant changes in the dynamic equilibrium of ecosystems. In fact, the interference of man can cause disorders to environmental factors [2], notably because of the disturbances in the natural habitat of several species, such as insect eating birds [3, 4, 5], insect bioindicators [6], and to plant species [7, 8, 9]. This has led to fragmented areas that are in various states of regeneration.

29

30 To quantify the quality of habitat for wildlife is a task that is extremely challenging, this being essential to the development of quantitative techniques with robustness sufficient to express the real ability of the natural shelters [10].

33

34 On the basis of the above, it is emphasized that the stratification of areas to study their quality, is a preponderant step to understand the peculiarities of each environment [11]. Summers et al. [12] point out that areas in reforestation can be divided into three main categories: assisted natural regeneration, direct sowing, and planting of seedlings; these areas being, according to Cunningham et al. [13], essential for the maintenance of

38

biodiversity. Already Botello et al. [14], dealing with areas of natural remnants, reported the contribution of these environments to the richness of fauna and flora. Fiorentin et al. [9] added that processes in natural regeneration areas are highly complex and dynamic due to the interaction of various processes that converge to drive ecological succession.

Owing to the importance of the ecosystems mentioned above, [10] point out that, their quantitative analysis consists of an important strategy for the generation of local environmental quality indicators in order to subsidize the decision making on areas to be destined for legal reserve. However, it is known that the evaluation of forest fragments is usually based on a single taxonomic group, evidencing the need for rapid assessments based on multitax on indicators [15] that can be excellent tools to help conservationists and managers in the definition of environmental conservation strategies [14]. The objective of this work was to analyze the biodiversity of three forest fragments under different forms of environmental conservation.

2. MATERIAL AND METHODS

2.1 Research Coverage Area

The research was carried out in April 2015 in three forest areas in the municipality of Jaboticabal, State of São Paulo, Brazil. The areas covered by the study, consisting of 720 m² each, were characterized according to the level of anthropization, namely: Area under reforestation conditions, denominated fragment 1 (FRA-1), located at 21°14'54.7"S and 48°17'48.5"W; area of natural remnant, fragment 2 (FRA-2), located at 21°14'47.1"S and 48°17'29.4"W; and an area under natural regeneration, fragment 3 (FRA-3), located at 21°15'02.5"S and 48°17'42.3"W. During the month of conduction of the research, climate variables were monitored: relative air humidity (RH%), accumulated rainfall (R mm), mean atmospheric temperature (AT °C), and mean solar radiation (SR MJ m²), as illustrated in Figure 1A and B.




Fig. 1. Climate variables. (A) relative humidity (RH) and rainfall (R), (B) atmospheric temperature (AT) and solar radiation (SR), * days the evaluations were performed. Jaboticabal, SP, 2015.

2.2 Experimental Design

The research was conducted in a completely randomized design (DIC), and the treatments represented by three forest fragments [16] with three replicates. To define the sample unit,

the fragments were divided into three parts of $\approx 240 \text{ m}^2$, where each one represented a repetition. In each fragment, three visits were carried out at different times, the first being at 08:00, the second at 12:00, and the third at 16:00 hours, in a randomized way between the sample units.

2.3 Survey of Data

In the analyzed fragments, variables inherent in the frequency of occurrence of fauna and flora were included in the taxonomic groups: insects (F-INS), amphibians (F-AMP), birds (F-BIR), mammals (F-MAM), reptiles (F-REP), trees (F-TRE), and these are condensed into the variable total species frequency (F-TES). The data obtained from these variables consisted of *in loco* observation. Therefore, these were considered as "clues", in order to facilitate the visualization of copies of the groups; indicators of their existence, such as the diversity of leaves, flowers and fruits that can serve as food and water; trees and soil for shelter; besides aptitude for hunting and coexistence of populations. The research was classified as exploratory [17], of the qualitative type [18].

The collection of vegetation information was carried out based on specialized literature [19], and two species are commonly found in the transition areas of the Atlantic Forest and Cerrado, mainly because they represent the vegetation of the State of São Paulo. The species chosen were araticum-de-terra-fria (*Annona emarginata* (Schltdl.) H. Rainer) and dairy (*Tabernaemontana catharinensis* A. DC.). This categorization was performed for greater precision in the visualization and obtaining of the data [20].

2.4 Statistical Analysis

The original data of the dependent variables were transformed into a sine arc of $\sqrt{(x/100)}$, to normalize the distribution of the deviations [21], data was then subjected to analysis of variance using the 5% probability F-test. For the significant variables, the Tukey test was applied for multiple comparisons of averages [22], in order to detect differences between the fragments. Subsequently, the original data were standardized and subjected to multivariate exploratory analysis, using cluster analysis (Ward's method) and Principal Component Analysis (PCA) [23].

3. RESULTS AND DISCUSSION

It was verified that there were significant differences ($P < .001$) between the analyzed fragments for the variables, total frequency of species (F-SPE), F-INS, F-BIR, F-MAM, and ($P = .05$), F-AMP, while for frequencies of occurrence of F-REP and F-TRE no significant differences were found ($P = .05$) as summarized in Table 1.


Table 1. Summary of variance analyses for the total frequency of species (F-TSP), insects (F-INS), amphibians (F-AMF), birds (F-BIR), mammals (F-MAM), reptiles (F-REP) and trees (F-TRE). Jaboticabal, SP, 2015.

F.V.	GL	Medium Squares						
		F-TSP	F-INS	F-AMF	F-BIR	F-MAM	F-REP	F-TRE
Fragments	2	144.81**	99.74**	33.88	57.16**	43.03**	3.66 ^{ns}	2.54 ^{ns}
Residue	6	8.80	9.10	6.45	5.16	1.99	7.32	0.63
CV (%)		17.19	32.25	68.30	26.68	32.86	212.13	10.49

*, ** and ^{ns} - significant at 1 and 5% probability of error and not significant by the Fischer test, F.V. - sources of variation, GL - degrees of freedom and CV - coefficient of variation.

125
126
127
128
129
130
131
132
133
134

By analyzing the total frequency of verified species, it was possible to verify that fragment two (FRA-2), expressed superiority of 26.7 e 80.0% in relation to the fragments (FRA-1) it's three (FRA-3), though FRA-2 and FRA-3 have not differed statistically from one another, with averages of 11 and 15. The FRA-1 was superior in 72.7% when compared with FRA-3, with an average frequency of 3 (Figure 2A). The same behavior was observed when the F-INS was analyzed, having recorded averages of 4.3, 5.3 and 0.6 for FRA-1, FRA-2 and FRA-3, respectively, with percent differences of 86.0 and 88.7% when comparing FRA-1 and FRA-2 with FRA-3 (Figure 2B).



135
136
137
138
139
140

Fig. 2. Total species frequency (A), with minimum significant difference (MSD) of 7.43 and frequency of insects (B), with MSD of 7.53, verified in three forest fragments. Mean data of three untransformed sine-arc replicates of the $\sqrt{(x/100)}$. Jaboticabal, SP, 2015.

141
142
143
144
145
146
147
148
149
150
151
152
153
154

Forest fragments, given the nature of their classification, express structural differences perceptible to the animals, in order to interfere with their behavior [24]. These authors attribute these behavioral changes, above all, to changes in the natural habitat of the species, while [25] contribute to changes in predation patterns. In a natural remnant fragment, the biological richness is undoubtedly superior to that of anthropized areas [14], reducing the incidence of solar radiation and temperature, increasing the relative air humidity and, thus, favoring the development of several species of fauna and flora [26]. Even in man-altered areas, some species can adapt and take advantage of this situation [27], justifying the frequency of species verified in natural regeneration areas in this work. Among the various species indicative of the quality of ecosystems, insects are often mentioned. For example, beetles [28, 29], caterpillars [30], bee species [31], and ants [32, 33, 6], are often quantified to express the level of environmental disorder based on how often they occur in environments.

155
156
157
158
159
160
161

Regarding the F-AMP, it was verified that FRA-2 had a higher mean (1.3), although it did not differ significantly from FRA-1 that had an average of 1, whereas FRA-3 expressed average 0, differing from FRA-2 being calculated a percentage difference of 100% between these two fragments (Figure 3A). For the variable F-BIR, this was also found to be superior to FRA-2, with an average 4.6, although this did not differ from FRA-1, with an average 2.6, with FRA-2 higher in 86.9% to FRA-3, where the mean was 0.6 (Figure 3B).




Fig. 3. Frequency of amphibians (A), with MSD of 6.37 and birds (B), with MSD of 5.69, verified in three forest fragments. Mean data of three replicates untransformed in sine-arc $\sqrt{(x/100)}$. Jaboticabal, SP, 2015.

Rapid assessments and biotic integrity indexes, although generally based on a single taxonomic group, are effective methods for assessing biodiversity conservation [15]. These researchers confirm the longing for this research by suggesting that multiparameter-based assessments, for example, the frequency of occurrence of amphibians and birds, provide a more robust assessment of environmentally disturbed forest fragments. Studying the distribution of amphibians, reptiles, birds and mammals, [14] mentions that anthropic habitats are unsuitable for these species, while natural remnants and reforested fragments are potentially habitable, explaining the high frequency of amphibians and birds in FRA-2, followed by intermediate frequencies in FRA-1 and critical values evidenced in FRA-3 in this research. It should be noted that amphibians are one of the most endangered animal classes, mainly because of their sensitivity to environmental changes (for example, habitat destruction, climate change, as well as the reduction of air humidity, or the emergence of new pathogens, such as the quitridio fungus, *Batrachochytrium dendrobatidis*) due to its dependence on water and its permeability of the skin [34]. The distribution results of birds verified by [15] corroborate the findings of this research, certainly due to the characteristics of a particular reduction of the presence of ornithopters in places of intense antropic activity. [24], studying bird species, reported that, although the vast majority of bird species are classified as highly and moderately sensitive to environmental disturbances, there are, although in a smaller number, less sensitive species, justifying the occurrence of birds in the FRA-3 of this study.

No significant difference was recorded between the means 1.0 and 1.6 of the FRA-1 and FRA-2, respectively, when analyzed against the variable F-MAM, however, these two fragments differed significantly from the FRA-3, where there was no presence of mammals, characterizing the superiority of 100% of FRA-1 and FRA-2 in relation to FRA-3 (Figure 4A). For the variables F-REP and F-TRE, no significant differences were found (Figure 4B), which can be justified by the high variation of the original data, reflecting a high coefficient of variation of 212.13% for F-REP absence of normal distribution of data F-REP and F-TRE.




Fig. 4. Frequency of mammals (A), with MSD of 3.54, reptiles and trees (B), with MSDs of 6.78 and 1.99, respectively, verified in three forest fragments. Mean data of three untransformed sine-arc replicates of the $\sqrt{(x/100)}$. Jaboticabal, SP, 2015.

Economic interests, to meet the demands of the growing population, have motivated predatory hunting, animal trafficking, forest deforestation, and expansion of arable land [35]. These researchers add that, fragmented forests tend to harbor fewer mammals compared with intact areas. It should be noted that the distribution dynamics of mammals in fragmented areas is also associated with their size. In fact, [36] report that the population of small rodent mammals can be increased in areas where the frequency of large mammals is reduced. In a complementary sense, [37] explain that changes in the distribution of mammals can be influenced by increasing land occupation for agriculture and livestock, as well as suppression of part of vegetation, alteration of hydrological cycles, burning regime and nutrient cycling in ecosystems. The nonoccurrence of differences between the fragments for the frequency of trees and reptiles can be justified by the fact that local climatic conditions favor the propagation and development of the trees, providing an adequate ecosystem for the occurrence of reptiles in the area of the three fragments studied [38].

On the basis of the Euclidean Distance used to summarize the homogeneity between the experimental units within the groups and heterogeneities between the groups, there were two main groups, the first group being represented by fragment three (FRA-3) and the second group by fragments one (FRA-1) and two (FRA-2), denoting the dissimilarities between the groups based on hierarchical grouping (Figure 5).




Fig. 5. Dendrogram of dissimilarities between three forest fragments (F). Jaboticabal, SP, 2015.

In a research to compare two multivariate methodologies in the study of similarities between fragments of Atlantic forest, [39] point out that there is dissimilarity between groups of forest fragments, emphasizing that fragment groupings are due to the similarities of their variables, justifying these similarities due to their geographical proximity. [40] also observed that floristic similarity decreased with increasing distance between areas, in agreement with the ideas of [41] and [42], according to which geographical proximity would be the only reliable factor to predict the similarity between areas. These evidences allow us to infer that, due to the high geographic proximity of the fragments investigated in this research, the differences observed are due to the particular characteristics of these fragments, mainly anthropization in fragments of natural regeneration and reforestation.

It is observed in Table 2, that the set of seven categories (variables) analyzed was summarized in two latent variables (constructs), called Principal Components 1 (PC_1) and 2 (PC_2), which were selected based on eigenvalues, 4.83 and 1.26 because they were ≥ 1 , satisfying the criterion of Kaiser-Meyer-Olkin (KMO).


Table 2. Eigenvalues (AV), relative variance (S^2_r) and absolute (S^2_a) and variable loads. Jaboticabal, SP, 2015.

PCs	AV	S^2_r (%)	S^2_a (%)	Variable Loads						
				F-TES	F-INS	F-ANF	F-AVE	F-MAM	F-REP	F-ARV
PC_1	4.83	69.07	69.07	-0.98	-0.93	-0.91	-0.96	-0.85	0.09	-0.73
PC_2	1.26	18.03	87.11	0.17	0.25	-0.06	0.19	-0.18	0.97	-0.38

PCs: Principal Components.

The PC_1 and PC_2 account for 87.1% of the total cumulative variance, with PC_1 accounting for 69.07% of this total, with PC_2 accounting for 18.04%. It is observed that the FRA-3

247 presented greater dissimilarities compared with the FRA-2, with the FRA-1 occupying
248 intermediate position (Figure 6).
249



250
251 **Fig. 6. Two-dimensional projection (Biplot) of three forest fragments (F) and**
252 **representatives of fauna and flora in two Principal Components (PC₁ and PC₂).**
253 **Jaboticabal, SP, 2015.**
254

255 Analyzing forest fragments through Principal Component Analysis, [35] report that the first
256 two components account for 56% of the total variation in mammalian distribution among the
257 sampled sites. These researchers point out that the lower incidence of mammalian species
258 in altered areas can be explained by the hunter's pursuit of animals, especially game
259 animals and those that cause damage to agricultural crops, while trees and nontarget
260 species of hunting tend to be seen more frequently in anthropized areas.

261
262 On the basis of the results, the use of Principal Component Analysis is justified, since it
263 provides a structural simplification of the original data. In fact, in the research carried out by
264 [39], 462 dimensions were reduced in 10 Principal Components resulting from linear
265 combinations between original variables. Therefore, these authors report that using the first
266 10 PCs is as efficient as the use of the 462 initial variables with regard to the explanation of
267 the variance. Thus, the use of two PCs in this research was sufficient enough to explain the
268 variance under study.

269
270 On the basis of this information, it is believed that the divergences evidenced between the
271 fragments of this research can be a reflection of the anthropic actions, especially of the
272 illegal hunting that can occur in these places. For [39], each fragment exhibits a species
273 composition that appears to result from a series of factors that varied differently over time
274 and space. Perhaps that is why it is so difficult to establish these areas. This difficulty,
275 however, indicates that each fragment presents a set of its own characteristics, which
276 emphasizes its importance in terms of conservation.

277

4. CONCLUSION

There are differences in the biodiversity of the forest fragments analyzed, highlighting the areas of natural remnants and reforestation with greater biological diversity, to the detriment of the natural regeneration areas with insufficient biological indicators, denoting adequacy of the first two fragments and inadequacy of the latter with respect to the potential of use as a legal reserve.

Two groups of environments were evidenced according to the potential hierarchy for use as legal reserve, the first group being characterized as inadequate and the second as adequate. Of the seven analyzed variables, six were considered essential to the correct evaluation of the environments.

COMPETING INTERESTS

The authors have not declared any conflict of interests.

AUTHORS' CONTRIBUTIONS

This work was carried out with the collaboration of all authors. All authors read and approved the final manuscript.

REFERENCES

1. Ashby MF. What is a "Sustainable Development"? In: Ashby MF, editor. Materials and Sustainable Development. Amsterdã: Butterworth-Heinemann; 2016.
2. Powell LL, Stouffer PC, Wolfe JD, Johnson EI, Hines JE, Nichols JD. Heterogeneous movement of insectivorous Amazonian birds through primary and secondary forest: A case study using multistate models with radiotelemetry data. *Biological Conservation*.2015;188(1):100-108. a
3. Arcilla N, Holbehc LH, O'Donnell S. Severe declines of understory birds follow illegal logging in Upper Guinea forests of Ghana, West Africa. *Biological Conservation*. 2015;188(1):41-49.
4. Cordeiro N, Dampf CJ, Borghesio L, Joho MP, Monoski TJ, Mkongewa VJ. Forest fragmentation in an African biodiversity hotspot impacts mixed-species bird flocks. *Biological Conservation*.2015;188(1):61-71.
5. PavlackyJr DC, Possingham HP, Goldizen AW. Integrating life history traits and forest structure to evaluate the vulnerability of rainforest birds along gradients of deforestation and fragmentation in eastern Australia. *Biological Conservation*.2015;188(1):89-99.
6. Rocha WO, Dorval A, Peres Filho O, Vaez C dos, Ribeiro ES. Formigas (Hymenoptera: Formicidae) bioindicadores de degradaçãoambientalemPoxoréu, MatoGrosso, Brasil. *Floresta e Ambiente*.2015;22(1):88-98. Portuguese.
7. Velazco SJE, Galvão F, Keller HA, Bedrij NA. Florística e fitossociologia de uma floresta estacional semidecidual, reservaprivada de Osununú-Misiones, Argentina. *Floresta e Ambiente*.2015;22(1):1-12. Portuguese.

- 331 8. SilvérioNeto R, Bento M de C, Menezes SJM da C de, Almeida FS. Caracterização da
332 coberturaflorestal de unidades de conservação da Mata Atlântica. Floresta e
333 Ambiente.2015;22(1):32-40. Portuguese.
334
- 335 9. Fiorentin LD, Téó SJ, Schneider CR, Costa RH, Batista S. Análise florística e
336 padrão espacial da regeneração natural em área de floresta ombrófila mista na região do
337 Caçador SC. Floresta e ambiente.2015;22(1):60-70. Portuguese.
338
- 339 10. Powell LL, Cordeiro NJ, Stratford J. Ecology and conservation of avian insectivores of
340 the rainforest understory: A pantropical perspective. Biological
341 Conservation.2015;188(1):1-10. b
342
- 343 11. Vogel HF, Cardoso O, Watzlawick LF, Campos JB. Pesquisas em unidades de
344 conservação urbanas no Paraná: conhecimentos raramente divulgados ou aplicados.
345 Ambiência - Revista do Setor de Ciências Agrárias e Ambientais.2015;11(1):75-
346 93. Portuguese.
347
- 348 12. Summers DM, Bryan BA, Nolan M, Hobbs TJ. The costs of reforestation: A spatial
349 model of the costs of establishing environmental and carbon plantings. Land Use
350 Policy.2015;44(1):110-121.
351
- 352 13. Cunningham SC, Mac Nally R, Baker PJ, Cavagnaro TR, Beringer J, Thomsonb JR,
353 Thompsonb RM. Balancing the environmental benefits of reforestation in agricultural
354 regions. Perspectives in Plant Ecology, Evolution and Systematics.2015;17(4):301-317.
355
- 356 14. Botello F, Sarka S, Sánchez-Cordero V. Impact of habitat loss on distributions of
357 terrestrial vertebrates in a high-biodiversity region in Mexico. Biological
358 Conservation.2015;184(1):59-65.
359
- 360 15. Medeiros HR, Bochio GM, Ribeiro MC, Torezan JM, Anjos L. Combining plant and bird
361 data increases the accuracy of an Index of Biotic Integrity to assess conservation levels
362 of tropical forest fragments. Journal for Nature Conservation.2015;25(1):1-7.
363
- 364 16. Mussury RM, Scalón S de PQ, Gomes AA, Batista MR, Scalón Filho H.
365 Flutuação populacional da mesofauna em fragmentos de mata na região de Dourados MS.
366 Ciência e Agrotecnologia, Lavras.2008;32(2):645-650. Portuguese.
367
- 368 17. Laake P, Fagerland MW. Statistical Inference. In: Laake P, Benestad H, Olsen BR,
369 editors. Research in Medical and Biological Sciences. 1st ed. Amsterdã: Academic
370 Press; 2015.
371
- 372 18. Moen K, Middelthon AL. Qualitative Research Methods. In: Laake P, Benestad H, Olsen
373 BR, editors. Research in Medical and Biological Sciences. 1st ed. Amsterdã: Academic
374 Press; 2015.
375
- 376 19. Lorenzi H. Árvores Brasileiras: Manual de identificação e cultivo de
377 plantas arbóreas nativas do Brasil. Nova Odessa: Instituto Plantarum; 2009. Portuguese.
378
- 379 20. Depoy E, Gitlin LN. Collecting data through measurement in experimental-type research.
380 In: Depoy E, Gitlin LN Introduction to Research. 5th ed. Amsterdã: Mosby; 2016.
381
- 382 21. Pimentel-Gomes F. Curso de estatística experimental. 14th ed. Piracicaba: Degaspari;
383 2000. Portuguese.

384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436

22. Santos JW, Almeida FAC, Beltrão NEM, Cavalcante FB. Estatística Experimental Aplicada. 2nd ed. Campina Grande: EmbrapaAlgodão/UFCG; 2008.Portuguese.
23. Hair JF, Black WC, Babin BJ, Anderson RE, Tatham RL. Análisemultivariada de dados. Porto Alegre: Bookman; 2009.Portuguese.
24. Stratford JA, Stouffer PC. Forest fragmentation alters microhabitat availability for Neotropical terrestrial insectivorous birds. *Biological Conservation*.2015;188(1):109-115.
25. Visco DM, Sherry TW. Increased abundance, but reduced nest predation in the chestnut-backed antbird in costa rican rainforest fragments: surprising impacts of a pervasive snake species. *Biological Conservation*.2015;188(1):22-31.
26. Pollock HS, Cheviron ZA, Agin TJ, Brawn JDAC. Absence of microclimate selectivity in insectivorous birds of the neotropical forest understory. *Biological Conservation*.2015;188(1):116-125.
27. Boyle WA, Sigel BJ. Ongoing changes in the avifauna of La Selva Biological Station, Costa Rica: Twenty-three years of Christmas Bird Counts. *Biological Conservation*.2015;188(1):11-21.
28. Filgueiras BKC, Tabarelli M, Leal IR, Vaz-de-Mello F, Iannuzzi L. Dung beetle persistence in human-modified landscapes: Combining indicator species with anthropogenic land use and fragmentation-related effects. *Ecological Indicators*.2015;55(1):65-73.
29. Salomão RP, Iannuzzi L. Dung beetle (Coleoptera, Scarabaeidae) assemblage of a highly fragmented landscape of Atlantic forest: from small to the largest fragments of northeastern Brazilian region. *RevistaBrasileira de Entomologia*.2015;59(1):126-131.
30. Maguire DY, Bennett EM, James PMA, Buddle CM. Landscape connectivity and insect herbivory: A framework for understanding tradeoffs among ecosystem services. *Global Ecology and Conservation*.2015;4(1):73-84.
31. Pereira SAN, Sousa CS. Levantamento da fauna de abelhas no município de montecarmelo-MG. *Getec*.2015;4(7):11-24.Portuguese.
32. De La Mora A, García-Ballinas JA, Philpott SM. Local, landscape, and diversity drivers of predation services provided by ants in a coffee landscape in Chiapas, Mexico. *Agriculture, Ecosystems and Environment*.2015;201(1):83-91.
33. Komonen A, Övermark E, Hytönen J, Halme P. Tree species influences diversity of ground-dwelling insects in afforested fields. *Forest Ecology and Management*.2015;349(1):12-19.
34. Grosjean S, Cruaud C, Ohler A, Hassanin A, Chuaynkern Y. Improving biodiversity assessment of anuran amphibians using DNA barcoding of tadpoles. Case studies from Southeast Asia. *ComptesRendusBiologies*.2015;338(1):351-361.
35. Meyer NFV, Helen JE, Ricardo M, Frank VL, Yorick L, David RO, Chantal BFV, Andrew DC, Clayton KN, Patrick AJ. "An assessment of the terrestrial mammal communities in

- 437 forests of Central Panama, using camera-trap surveys". *Journal for Nature*
438 *Conservation*.2015;(6):28-35.
- 439
- 440 36. Galetti M, Bovendorp SR, Guevara R. Defaunation of large mammals leads to an
441 increase in seed predation in the Atlantic forests. *Global Ecology and*
442 *Conservation*.2015;3(1):824-830.
- 443
- 444 37. Ripple WJ, Newsome TM, Wolf C, Dirzo R, Everatt KT, Galetti M, Hayward MW, Kerley
445 GIH, Levi T, Lindsey PA, Macdonald DW, Malhi Y, Painter LE, Sandom CJ, Terborgh J,
446 Valkenburgh BV. Collapse of the world 's largest herbivores Collapse of the world 's
447 largest herbivore. *Science Advances*.2015;(5):1-12.
- 448
- 449 38. Paula DM de, Groeneveld J, Huth A. Tropical forest degradation and recovery in
450 fragmented landscapes — Simulating changes in tree community, forest hydrology and
451 carbon balance TL - 3. *Global Ecology and Conservation*.2015;(3):664-77.
- 452
- 453 39. Ferreira RLC, Mota AC, Silva JAA, Marangon LC, Santos ES. Comparação de
454 duas metodologias multivariadas no estudo de similaridade entre fragmentos de
455 floresta atlântica. *Revista Árvore*.2008;32(3):511-521. Portuguese.
- 456
- 457 40. Jacquemyn H, Butaye J, Dumortier M, Hermy M, Lust N. Effects of age and distance on
458 the composition of mixed deciduous forest fragments in an agricultural landscape.
459 *Journal of Vegetation Science*.2011;(12):635-642.
- 460
- 461 41. Condit R. Defining and mapping vegetation types in mega-diverse tropical forests.
462 *Trends in Ecology and Evolution*.1996;11(1):4-5.
- 463
- 464 42. Cook SA. Diversity of approaches to the study of species richness. *Trends in Ecology*
465 *and Evolution*.1998;13(9):340-341.