

Antioxidant Potential and anti-sickling activity of different organs of *Curcuma longa*: correlation of the antioxidant capacity on anti-sickling activity

ABSTRACT

Aims: The aim of this research was to evaluate comparatively the anti-sickling activity of aqueous and methanol extracts of different organs of *C. longa*, to determine the polyphenolic compounds and to evaluate the antioxidant potential of different parts of *C. longa* organs. Also, to establish a correlation between the antioxidant potential of the extracts on the anti-sickling and to look for a possible correlation between the antioxidant potential and the anti-sickling activity of the tested extracts in order to predict the roles of the antioxidant potential of extracts on the anti-sickling activity.

Study design: The study used an experimental design. The antioxidant and anti-sickling activities of the crude aqueous and methanol extracts of different organs of *Curcuma longa* were assessed along with their correlation.

Place and Duration of Study: The study was carried out at the Laboratory of Natural Products, Department of Chemistry, Laboratory of Analyses and Research on Food and Nutrition (LARAN), Department of Biology, both at the Faculty of Science and at Centre for the Study of Natural Products from Plant Origin (CESNOV), Faculty of Pharmaceutical Sciences, both at University of Kinshasa in 2018.

Methodology: *C. longa* was cultivated in an experimental setting. The plant was planted in an experimental monoculture garden thanks to its rhizomes and various organs (rhizomes, roots, leaves, petals and sepals) were then harvested, packaged and macerated in methanol and water. The macerates were then filtered and the filtrates dried in the oven to obtain the dry aqueous and methanol extracts used for biological analyses (antioxidant and anti-sickling activity) and for the determination of polyphenolic compounds. The determination of polyphenolic compounds was performed using spectrophotometric methods as reported by standard protocols while the evaluation of antioxidant and anti-sickling activities was performed via the ABTS, DPPH and Emmel tests respectively. The blood used for the Emmel test was obtained at the Centre de Médecine Mixte et d'Anémie SS (CMMSS) from sickle cell patients. The GraphPad Prism 6.0 software was used for data analysis for all statistical analyzes and the determination of IC₅₀.

Results: The findings showed that the polyphenolic compounds are not distributed in the same way in different organs of *C. longa*. The extracts of different parts of *C. longa* do not all have a great antioxidant potential; Rhizome and root extracts showed better antiradical activity using ABTS method compared to other organs, while rhizome and leaf extracts showed better antiradical activity using the DPPH test. The analyses carried out revealed: a positive correlation between (i) the total polyphenol contents and the anti-sickling activity of the studied extracts ($r = 0.08328$ and (ii) between the antioxidant activity by the test ABTS and DPPH along with the anti-sickling activity of different extracts ($r = 0.1281$ and 0.5232 respectively).

Conclusion: The tested extracts displayed good antioxidant and anti-sickling activities. Thus, different organs of *C. longa* are a source of natural antioxidants and/or active ingredients that can be used to manage sickle cell disease. The effect of the antioxidant potential of the extracts tested on their anti-sickling activity has been demonstrated. Nevertheless, in-depth studies are required in order to confirm this correlation.

Keywords : *Curcuma longa*, antioxidant activity, anti-sickling activity, Polyphenol content, Correlation.

1. INTRODUCTION

Sickle cell disease is a genetic disease with autosomal recessive inheritance linked to an abnormality of hemoglobin. On the molecular level, it is a hemoglobinopathy resulting from a point mutation on the

sixth codon of the β -globin gene carried by chromosome 11. It is a substitution of Adenine by Thymine at the 6th level codon of the β -globin gene. At the protein level, this leads to the replacement (in the β -globin chain) of glutamic acid (polar amino acid) in position six by a valine (apolar amino acid) ($\beta 6 \text{ Glu} \rightarrow \text{Val}$) and the synthesis of an abnormal hemoglobin, hemoglobin S (Hb S), in English "sickle" [1-4]. Sickle cell disease is responsible for the loss of life worldwide, the most vulnerable layer of which is children under 5 years old [5].

From its regions of origin (Central and West Africa), it has moved and continues to move with migratory flows (Mediterranean region, Asia, USA, the Caribbean). However, populations of African origin are particularly affected. In some West African countries, 30-40% of the population would carry the gene [3]. Several treatment modalities have been considered to relieve patients, including bone marrow transplantation, gene therapy, blood transfusions and hydroxy-urea intake. The truth is that these treatments are not only ineffective and very costly for poor people in Africa, but can be the cause of several adverse effects (toxicity, erythrocyte incompatibility, graft versus host reaction, HIV/AIDS contamination) [5-7]. In the Democratic Republic of the Congo (DRC), the phytotherapy is currently used as an alternative which can provide relief to sickle cell disease. The review of the literature indicated that several plants have been screened having the anti-sickling potential activity of which "*Curcuma longa*" [2, 5-9, 61-69].

Belonging to Zingiberaceae family, *C. longa* L. is a perennial herbaceous plant measuring up to 1m high. It is mostly cultivated in the tropics but a large part of the production is carried out in India and Southeast Asia [9-10].

Several studies reported that the intense cultivation of *C. longa* is due to the use of rhizomes. For centuries, *C. longa* rhizomes have been efficiently used in various sectors including the food, cosmetology, agri-food, pharmaceutical, textile and traditional medicine industries. These uses have been reported by several authors [9-38]. Furthermore, its leaves are used in industry for the extraction of essential oils, in traditional kitchens for the seasoning of dishes or as a colouring agent. In traditional medicine, it is used as tonic, stimulant and blood purifier in the valley districts of Manipur, India [23, 28, 34-36]. Mbadike *et al.* [9], reported that extracts of rhizomes of *C. longa* have broad pharmacological potential such as anti-cancer, anti-inflammatory, healing, cholesterol-lowering, hypoglycemic, anti-Alzheimer's, antiparasitic, anti-inflammatory, antioxidant, antibacterial, antifungal, anti-venom, antipyretic, analgesic, inhibits the action of the integrase of HIV-1 replication and integrase protein of HIV-1, protects against diabetic retinopathy and numerous other pathologies. However, several studies have shown that the pharmacological actions reported on rhizome extracts of *C. longa* are linked to the presence of polyphenolic compounds, particularly curcuminoids including curcumin, but also to its essential oils [8-10, 12, 23-24, 39-40, 41-53]. Furthermore, the antibacterial, antioxidant, hypoglycemic, analgesic potential of leaf extracts have also been reported [20, 26, 31, 34, 54-57]. Ajay *et al.* [24] and Camatari *et al.* [17], showed in their study that *C. longa* flowers possess the antioxidant potential. In addition, Lassa [58], reported that *C. longa* rhizomes is used for the treatment of SS anemia in traditional medicine of the Democratic Republic of the Congo (DRC). This use was scientifically justified by the research carried out by Mbadike *et al.* [9] whom showed the anti-sickling potential of total methanol extracts from different organs (rhizomes, roots, leaves, sepals and petals) of *C. longa*. Henceforth, the interest to carry out a comparative study on the anti-sickling activity of total methanol and aqueous extracts of different organs of *C. longa*.

However, numerous data from the literature have shown the involvement of oxidative stress in sickle cell anemia. Hierso [1] reported that sickle red blood cells produce more free radicals than normal red blood cells and that the increase in oxidative biomarkers would actively participate in the physiopathology mechanisms of the disease and play a role in the hemolysis, the onset of vaso-occlusive crises and the attacks of various organs. Due to oxidative damage associated with sickle cell disease, the development of therapeutic antioxidant strategies is proving to be essential. Hence, the main goal of this study was to (i) evaluate the antioxidant potential of different parts of *C. longa* (rhizomes, roots, leaves, petals and sepals) and (ii) look a possible correlation between the content of polyphenolic compounds, the antioxidant potential and the anti-sickling activity of the extracts.

2. MATERIALS AND METHODS

2.1 Biological material

Different organs of *C. longa* (rhizome, root, leaf, petal and sepals) were used in this study as biological material. The plant was first cultivated as a monoculture in an experimental garden and the different

parts were later harvested. After harvesting, the organs of *C. longa* were brought to the laboratory where they were washed, dried in the oven (Melag Nur Für Wechselstrom brand) at 30 °C and crushed using a Moulinex electric mill. The powders resulting from these steps were used for extract preparation. Later, they were useful for the phytochemical and biological screening.

2.2. Determination of Total Phenolic content

The dosage of the polyphenolic compounds was carried out according to the method described by Kapepula *et al.* and Adepado *et al.* [59-60]. In fact, 10 mg/mL of each extract were diluted in methanol 80% in order to obtain a solution of 1mg/mL for each extract. Then, a reaction mixture consisted of 0.5 mL extract; 5.0 mL distilled water and 0.5mL Folin-Ciocalteu reagent was prepared for each extract. After three minutes, 1.0 mL of a saturated solution of Na₂CO₃ 20% was added. The mixtures prepared were stirred and incubated at laboratory temperature away from light for one hour. The absorbances of the reaction solutions were read using a spectrophotometer at 725 nm. Each dose was repeated three times. The quantity of total polyphenols was expressed in mg of gallic acid equivalents (GAE)/g using the following equation from the calibration line:

$$y = 1.7097 \ln(x) + 5.2062 \quad [1]$$

and R²=0,965, where x is the absorbance and y is the equivalent of gallic acid (mg/g).

2.3. Dosage of total flavonoids

The estimation of the total flavonoid content of our extracts was performed using a spectrophotometer. Aluminium trichloride forms a yellow complex with flavonoids that absorbs at 415 nm [60]. First, a reaction mixture containing 1mL of the methanol solution (80%) of each of the extracts (concentration 10mg/mL) and 1mL of AlCl₃ (2%) (dissolved in methanol) was prepared. The mixtures were then stirred and incubated for one hour at the laboratory temperature under shade and the absorbances were finally read at 415 nm. The mixtures were prepared in triplicate for each analysis and the average value was retained. For the preparation of the blank (control), it was performed in the same way except that instead of the extract, 1 mL of methanol was added. The flavonoid content of the extracts was expressed in mg quercetin equivalent (QE)/g using the following equation from the calibration line:

$$y = 0.5001 \ln(x) + 3.442 \quad [2]$$

R² =0,944, where x is the absorbance and y is the equivalent of quercetin (mg/g).

2.4. Determination of total anthocyanins

The determination of the anthocyanin content of extracts was carried out following the protocol as per Adepado *et al.* [60]. We took 0.5 mL of the 1mg/mL solution of extract prepared with MeOH 80%, which was mixed with 3 mL of vanillin methanol solution (4%) and 1.5 mL hydrochloric acid. The mixture was incubated for one hour, then the absorbance was measured at 500 nm. Mixtures were prepared in triplicate, and the average value was taken. The same procedure was repeated for the reference substance (D-catechin) in the concentration range (from 4 mg/mL to 0.125 mg/mL) and allowed the calibration line to be established. The anthocyanin content of the extracts was expressed in mg equivalent D-catechin per g of dry vegetable matter (EC/g) using the following equation from the calibration line:

$$y = 0.0728 x + 0,0171 \quad [3]$$

and R²= 0,994, where x is the absorbance and y is the equivalent of catechin (mg/g).

2.5. Assessment of the anti-sickling activity

a. Blood samples

The blood samples used in our experiments were collected at the Centre de Médecine Mixte et d'anémies SS (CMMSS) from the sickle cell patients. Once at the laboratory, they were kept in the refrigerator and the Emmel's test was performed 24 hours later to ensure that these samples were actually taken from sickle cell patients.

b. Preparation of dry methanol extracts of various organs of *C. longa*

Using a precision scale (Kern 440-35N), it was weighed respectively 25g of roots and rhizomes, 20g of leaves, 3.69g of sepals and 5g of petals which were macerated in 100 mL of methanol during 24 hours. The macerates were then filtered with a Whatman n°1 filter paper and oven dried at 37 °C (Melag Nurfur Wechselstrom) for five days.

c. Preparation of dry aqueous extracts of various organs of *C. longa*

With a precision scale (Kern 440-35N), it was weighed respectively 20g of rhizomes and roots, 15g of leaves and 10g of sepals and petals of *C. longa* powders that were diluted in 150 mL of distilled water. After 24 hours of maceration, the macerates were filtered using a Whatman n°1 filter paper, poured into petri dishes and dried in an oven at 40 ° C for a week. The dry extracts obtained were used for the Emmel's test.

d. Emmel's test

The anti-sickling activity of the total aqueous and methanol extracts of different parts of *C. longa* was evaluated using the Emmel's test, as described previously [8-9, 61-69]. Another microscopic preparation not containing extracts was made under the same conditions and served as a negative control. All the analyses were carried out in triplicate in order to confirm the obtained results.

The principle of this test is the following: Hemoglobin S is less soluble when the oxygen content decreases (hypoxia, $pO_2 < 45 \text{ mmHg}$). In sickle cell disease, the sickling of erythrocytes is observed due to the intracellular crystallization of hemoglobin S in the form of tactoids. Therefore, unlike the normal red blood cell, which has a biconcave or circular shape, the red blood cells of sickle cell patients have a sickle or banana shape under hypoxic conditions. The effect of crude plant extracts, which prevents cell sickling under hypoxic conditions (created by the addition of 2% sodium metabisulfite, m/v), is used as an indication of the pharmacological activity of these plants.

First, it was question of preparing different concentrations of extracts of different parts of *C. longa* (1mg/mL ; 500 µg/mL, 250 µg/mL, 125 µg/mL, 62.5 µg/mL, 31.25 µg/mL and 15.62 µg/mL) using saline solution (NaCl 0.9%, m/v) as solvent. From that solution, it allowed to make the microscopic preparation by placing one drop of SS blood in the slide to which was added a drop of sodium metabisulfite ($\text{Na}_2\text{S}_2\text{O}_5$ 2%) in order to create a hypoxic condition. To that preparation was added a drop of the test drug (see different concentrations prepared above) which was mixed thoroughly and was carefully covered in order to avoid the formation of air bubbles., (d) mix thoroughly and cover with the object cover plate avoiding the formation of air bubbles. The edges of the slide was supercooled with paraffin to reinforce the hypoxic solution. The results was read after 3-24 hours using a microscope (Bresser Biolux NV brand) at 40 of magnitude.

2.6. Antioxidant activity

The evaluation of the antioxidant activity was performed using the ABTS and DPPH as described by Kapepula *et al.* [59].

a. Preparation of samples

10mg of dry extract from each sample was dissolved in 1mL of methanol for polar extracts and dichloromethane-methanol mixture (1:1) for apolar extracts (solution A : 10 mg/mL). We then diluted to the following concentration levels: 8mg/mL, 6 mg/mL, 4mg/mL and 2 mg/mL.

b. Preparation of the ABTS radical

In 500 µL of distilled water was dissolved a quantity of the ABTS reagent corresponding to 20 millimoles: solution A. In another 500µL of distilled water was dissolved a quantity of potassium persulphate ($\text{K}_2\text{S}_2\text{O}_8$) corresponding to 10 millimoles: solution B. Both solutions were mixed at the same volume and the mixture was kept under shade for 12 or 16 hours. This mixture constituted the ABTS•⁺ radical stock solution. Then the stock solution of the radical was diluted several times with methanol to obtain an analytical solution having an absorbance ranging from 0.800 to 1.000

c. Preparation of the DPPH radical

A quantity of 3.2 mg of DPPH reagent was dissolved in 100 mL of methanol 80% and this solution was kept under shade for at least one hour. The absorbance of this solution was adjusted to 0.7 ± 0.05 using methanol 80%.

d. ABTS test

When reacted with potassium or sodium persulfate ($\text{K}_2\text{S}_2\text{O}_8$), ABTS (2,2'-azino-bis-3ethylBenz-Thiazolin-6-Sulfonic acid) forms the cationic radical ABTS-⁺ from blue to green. The addition of antioxidant reduces this radical and causes the mixture to discolour. The radical discoloration measured by spectrophotometer at 734 nm is proportional to the antioxidant concentration [59]. For each sample and concentration level (8mg/mL, 6 mg/mL, 4mg/mL and 2 mg/mL), 20µL of the solution was collected

and 1980µL of the ABTS-+ radical analysis solution was added later. The reaction mixtures were incubated for 30 minutes and the solutions were successively read with a spectrophotometer at 734 nm. The same procedure was repeated for the preparation of the white consisting of methanol (20µL) and ABTS (1980µL). The mixtures were made in triplicate for each analysis.

The percentage of inhibition of the ABTS-+ radical by the sample is determined using the following formula:

$$\% \text{ of inhibition} = [1 - (A_x/A_c)] \times 100 \text{ [d]}$$

A_x : the absorbance of the ABTS-+ radical in the presence of the extract

A_c: the absorbance of the ABTS-+ radical (control solution)

e. DPPH test

The DPPH test is based on the degradation of the DPPH- (2,2 DiPhenyl-1- PicrylHydrazyl) radical. The DPPH- radical is a violet coloured radical, the addition of antioxidant reduces this radical and causes the discoloration of the mixture; this discoloration of the radical measured by spectrophotometer at 517 nm is proportional to the concentration of antioxidants [59]. For each sample and dilution level, 20 µL of the solution was taken and 1980µL of the DPPH- radical analysis solution was added. The reaction mixtures were incubated in a dark place for 30 minutes and the solutions were read successively with a spectrophotometer at 517 nm. We proceeded in the same way for the preparation of the blank, except that instead of the extract, 20 µL of methanol was added. As with the ABTS test, the mixtures were made in triplicate.

The percentage of inhibition of the ABTS-+ radical by the sample is determined using the following formula:

$$\% \text{ of inhibition} = [1 - (A_x/A_c)] \times 100 \text{ [d]}$$

A_x : the absorbance of the ABTS-+ radical in the presence of the extract

A_c: the absorbance of the ABTS-+ radical (control solution)

2.7. Data analysis

The GraphPad Prism 6.0 software was used for data analysis for all statistical analyzes and the determination of IC₅₀. For data presented as mean ± standard deviation, we performed the ANOVA one Way test for comparison of sample averages, followed by Tukey's pairwise multiple comparison test. We considered the threshold of significance at α = 0.05. In addition, Pearson's test was used in order to assess the correlation between some variables.

3. RESULTS AND DISCUSSION

3.1. Total Phenolic content

Table 1 reports the contents of polyphenolic phytoconstituents in the various organs of *C. longa*.

Table 1: Content of polyphenolic phytoconstituents in different organs of *C. longa*

Samples	Tota polyphenol content (mg/EAG) (n=3)	Flavonoids content (mg/EQ) (n=3)	Anthocyanin content (mg/EC) (n=3)
Rhizomes	12.82 ± 0.63 ^a	0.020 ± 0.001 ^a	0.29 ± 0.05 ^a
Roots	5.91 ± 0.87 ^b	0.010 ± 0.001 ^b	0.17 ± 0.04 ^{a,b}
Leaves	4.31 ± 0.68 ^c	0.020 ± 0.002 ^c	0.30 ± 0.19 ^a
Petals	1.53 ± 0.28 ^d	0.007 ± 0.000 ^d	0.05 ± 0.01 ^{b,c}
Sepals	0.59 ± 0.30 ^d	0.006 ± 0.001 ^d	0.01 ± 0.00 ^{b,c}

Note: Mean values followed by the same letters in each column do not differ significantly at $p \leq 0.05$ by the Tukey test. n= number of repetition of the experiment

The Analysis of variance showed that extracts from different organs of *C. longa* showed very highly significant differences (P value $<0, 0001$) in total polyphenol content. However, the Tukey pair multiple comparison test ($HSD = 0.4938$) does not reveal a significant difference between roots and leaves and between sepals and petals. This indicates that the total polyphenol content is low in the sepals and petals. It follows the following descending order: Rhizomes> roots, leaves> petals and sepals. This shows that rhizomes are an important source of *C. longa* total polyphenols. Maizura *et al.* [14] also showed that *C. longa* rhizomes are rich in polyphenolic compounds. Although the rhizome samples used in their study had a high content of total polyphenols (67.9 ± 1.0 mg GAE/100g extracts) compared to those used in this study. This may be explained by the fact that these two samples were not collected in the same geographical environment. Other abiotic and biotic factors may also explain this difference. Gayatri *et al.* [70], reported that the rhizomes of *C. longa* have a high content of total polyphenols compared to other species of *Curcuma* genus.

In contrast to the present study, of which floral extracts showed low levels of total polyphenols, the studies by Yan *et al.* [26], Ajay *et al.* [24] and Camatari *et al.* [17] revealed the presence of large amounts of total polyphenols in flower extracts (2013.09 ± 5.13 mg GAE/100g, 210.45 ± 1.32 mg GAE/100 g and 119 mg ± 2 GAE/g respectively). Chang *et al.* [71] also reported that *C. longa* flowers were rich in polyphenolic compounds (including curcuminoids). In fact, the content of secondary metabolites and in this case the polyphenolic compounds may vary with respect to the geographical origin and the period of harvesting of the plant, to the physicochemical and biochemical characteristics of the soil, to the method and solvents used for the treatment. extraction, at the storage and post-harvest treatment conditions, of the genotype of the plant, the part of the plant used [12-13, 56, 72-75].

Regarding our study, the post-harvest treatment of floral pieces may have influenced polyphenolic content. Indeed, while drying floral pieces in the shade at the laboratory temperature; these organs tend to soften and lose color; which implies a deterioration of pigmentary principles including polyphenolic compounds. The degradation of plant polyphenolic compounds under cellular conditions i.e. fresh samples has also been reported by Yan *et al.* [26], and Chang *et al.* [71]. Moreover, Yan *et al.* [26], reported that dried and powdered *C. longa* leaves have a high total polyphenol content as fresh samples. The author justified this observation by the fact that the degradation of phytoconstituents in cellular conditions probably due to the activation of hydrolases and/or cellular oxidases. Secondary metabolites are synthesized mainly in response to environmental constraints (biotic or abiotic stresses) and intervene in plant-environment interactions. Some secondary metabolites are considered as signaling or messenger molecules and are involved in the initiation of certain biochemical and / or physiological processes of the plant. These processes explain the presence or absence of phytoconstituents during a period of the plant cycle or according to the environment [76]. Shiyu *et al.* [12], reported that the concentration of certain polyphenolic compounds, especially curcuminoids in the rhizomes of *C. longa*, becomes important five months after planting and declines between the 9th and 10th months. They also reported that mother or primary rhizomes are rich in phenolic compounds as secondary rhizomes [12].

However, it was observed low levels of flavonoids and total anthocyanins in the various extracts under study. Analysis of variance also showed a very highly significant difference in the distribution of flavonoids in different parts of *C. longa* (P value $<0, 0001$). Pairwise multiple comparison by the Tukey test ($HSD = 0.0005357$) shows that there is no significant difference in flavonoid levels in the petals and sepals. The order of flavonoid concentration in the different parts of *C. longa* is as follows: Leaves> rhizomes> roots> sepals and petals. The flavonoid contents are expressed in mg equivalent of quercetin per g of dry extracts. Meanwhile, the analysis of variance revealed a highly significant difference in anthocyanin levels in the various organs of *C. longa* under study (P value = 0.0073). Although the Tukey multiple comparison test ($HSD = 0.07237$) reveals a significant difference only between rhizomes and sepals, sepals and leaves and petals and leaves. This shows that the petals and sepals are poor in anthocyanins. The findings of this study show that the polyphenolic phytoconstituents studied are not distributed in the same way in the different organs of *C. longa*. It should be noted that studies on the determination of flavonoid and anthocyanin content of different organs of *C. longa* are still poor.

3.2. Anti-sickling activity

The anti-sickling activity of total aqueous and methanol extracts of various organs of *C. longa* are evident from the findings of this study (Figure 1). The anthocyanin-rich leaf extracts showed a great anti-sickling activity (15.62 µg/mL). According to Mpiana *et al.* [61-68], Tula [69] and Ngbolua *et al.* [7], the anti-sickling activity of plants is generally related to the presence of anthocyanins and organic acids. In fact, anthocyanins interact with the hemoglobin S and compete with the polymerization reaction by inhibiting the erythrocyte sickling process [2, 69]. This could justify the fact that leaf extracts showed at the same time high anthocyanin content, and a strong anti-sickling activity. However, the total aqueous extracts of rhizomes, leaves and sepals showed a high anti-sickling activity compared to total methanol extracts. The anti-sickling activity of total methanol extracts of different organs of *C. longa* has also been demonstrated by Mbadiko *et al.* [9].




Figure 1. The normalization of sickle cells treated on the one hand with total methanol leaf extracts of 31.25 µg/mL (B); rhizomes of 62.50 µg/mL (C), petals of 62.50 µg/mL (D), roots of 62.50 µg/mL (E), sepals of 62.50 µg/mL (F), and the other hand, sickle cell treated with total aqueous rhizome extracts of 31.25 µg/mL (G), roots of 62.50 µg/mL (H), petals of 62.50 µg/mL (I), sepals of 31.25 µg/mL (J) and leaves 15.62 µg/mL (K) compared to control, blood SS erythrocytes not treated (A).

The analysis of variance showed a very highly significant difference compared to the anti-sickling activity of the methanol total extracts of different organs (P value <0.0001). This difference is not significant at the 5% probability threshold. The Tukey multiple comparison test did not show a significant difference between rhizome and root, rhizome and sepal, and lastly between rhizome and petal activity. This reflects the following order in relation to the anti-sickling of extracts from different organs: Leaves >

Rhizome, root, petal and sepal. In addition, the ANOVA also showed a very highly significant difference in anti-sickling activity of total aqueous extracts of different organs of *C. longa* (P value <0.0001). However, the Tukey Multiple Comparison Test did not reveal a difference between the rhizome and sepal, and then between root and petal. The anti-sickling activity of aqueous extracts follows the following order: Leaves> rhizome, sepal >Petal root.

3.3. Antioxidant activity

The antioxidant activity of extracts from different organs of *C. longa*, the relationship between polyphenol content and antioxidant activity and the relationship between total polyphenol content and / or antiradical activity and anti-sickling activity is presented in the table below.

Table 2: Evaluation of the antiradical activity of the methanol extracts from different organs of *C. longa* expressed in value of the inhibitory concentration 50 (IC 50) by the ABTS and DPPH techniques

Samples	ABTS (µg/mL) (n=3)	DPPH (µg/mL) (n=3)
Roots	4.66 ± 0.34 ^a	-
Rhizomes	4.43 ± 0.18 ^a	15.63 ± 7.81 ^a
Sepals	46.24 ± 2.51 ^b	1129.8 ± 559.64 ^b
Petals	24.27 ± 1.43 ^c	553.35 ± 250.15 ^c
Leaves	14.93 ± 4.30 ^d	54.83 ± 19.35 ^d
Gallic acid (Control)	0.71 ± 0.08	1.07 ± 0.10

Note: Mean values followed by the same letters in each column do not differ significantly at $p \leq 0.05$ by the Tukey test. n= number of repetition of experiment

The analysis of variance revealed a very highly significant difference with regard to the antioxidant activity (ABTS test) of *C. longa* extracts (P value = 0.0001). The Tukey pair multiple comparison test (HSD = 0.5365) does not reveal a significant difference between rhizome and root extracts. The intensity of antiradical activity by the ABTS test follows the following descending order: Rhizomes, roots > leaves > petals > sepals. However, the ABTS method helps to evaluate the antioxidant activity of the mixtures of substances and also to distinguish their additive and synergistic effects. Because the radical ABTS reacts at the same time with the lipophilic and hydrophilic compounds (in particular the polyphenols) while the DPPH reacts only with the polar or hydrophilic compounds [59, 77]. Thus, the fact that the rhizomes and roots of *C. longa* showed better activity with the ABTS test proves that these organs have a high content of lipophilic (non-polyphenolic compounds) and hydrophilic (polyphenolic compounds) compounds responsible for the antioxidant activity. It could also demonstrate that the antiradical activity of rhizome and root extracts is related to the additive or synergistic effect of polyphenolic and non-polyphenolic compounds. The analysis of variance on the antioxidant activity of extracts of different organs of *C. longa* with DPPH test showed a very highly significant difference in the antioxidant activity of these organs (P value= 0, 0001). However, the Tukey multiple-pair comparison test indicates that there is no significant difference between rhizome and leaf extracts. The antiradical activity of extracts from different parts of *C. longa* follows the following descending order: Rhizomes> leaves > petals > sepals. It seems that the antioxidant activity of *C. longa* leaves is largely related to polyphenolic compounds because the DPPH radical reacts particularly with hydrophilic compounds including polyphenols [59, 77].

In addition, for the same sample, the analysis of the antioxidant activity by the ABTS test showed in most cases lower IC₅₀ values than those of the DPPH test. This asserts the hypothesis on the additive or synergistic effect of polyphenolic and non-polyphenolic compounds in the antioxidant activity of extracts studied and reveals that non-polyphenolic compounds contribute significantly to their antioxidant property. The synergistic action of the polyphenolic and non-polyphenolic fraction on the antioxidant potential of plant extracts has also been reported by Maizura *et al.* [14]. Several studies have shown that rhizome extracts of *C. longa* exhibited an antioxidant activity; and this antioxidant potential is mainly due to the presence of polyphenolic compounds, including curcuminoids. [12-13, 15-16, 18,

27, 32-33, 37-39, 49,51, 73, 78-86]. According to these authors, curcuminoids notably Curcumin is ten times more antioxidant than Vitamin E.

By the way, Jelena *et al.* [28] , reported that the essential oils extracted from *C. longa* rhizomes also have a very strong antioxidant action. There is little information in the literature on the antioxidant activity of root, leaves and floral extracts. The antioxidant activity of *C. longa* leaf, roots and flower extracts was reported as well by Ajay *et al.* [24], Camatari *et al.* [17] and Ritwiz *et al.* [54].

The comparison of total polyphenol content and the antioxidant activity is presented in table 3 below.

Table 3: Comparison of total polyphenol content and antioxidant activity of different organs of *C. longa*.

Extracts from plants	Total Polyphenol Content (mg/EAG)	Antioxidant activity	
		ABTS (µg/mL)	DPPH (µg/mL)
Rhizomes	12.82 ± 0.63	4.43 ± 0.18	15.63 ± 7.81
Roots	5.91 ± 0.87	4.66 ± 0.34	-
Leaves	4.31 ± 0.68	14.93 ± 4.30	54.83 ± 19.35
Petals	1.53 ± 0.28	24.27 ± 1.43	553.35 ± 250.15
Sepals	0.59 ± 0.30	46.24 ± 2.51	1129.80 ± 559.64

The correlation between the total Polyphenol content and the antioxidant action of the ABTS and DPPH test of different extracts was evaluated by the Pearson test. The results obtained highlight the existence of a negative correlation between the concentration of total polyphenols and the antioxidant activity of different extracts by the ABTS test ($r = -0.7807$) but yet this correlation is not significant at the 5% probability level. The analyzes also showed a negative correlation between the concentration of total polyphenols content and the antioxidant activity by the DPPH test ($r = -0.7445$). The non-significant negative correlation between total polyphenol content and the *in vitro* antiradical activity by the ABTS and DPPH test indicates that the antiradical activity of the studied extracts is not related solely to the presence of the Polyphenol compounds. Nevertheless, many studies have reported a positive correlation between the total polyphenol concentration of plant extracts and their antioxidant potential [13-14, 17, 24, 26, 70, 75]. Shan *et al.* [88] and Wu *et al.* [89], have also reported that phenolic compounds contribute significantly to the antioxidant activity of plant extracts. Velioglu *et al.* [90] and Kaur *et al.* [91], have also shown that there is a linear correlation between the total polyphenol content of plants and their antioxidant potential.

The relationship existing between the polyphenol content, antioxidant and anti-sickling activities is given in the following table.

Table 4: Relationship between polyphenol contents, antioxidant activity and anti-sickling activity.

Extracts from plants	Total Polyphenol Content (mg/EAG) (n=3)	Antioxidant activity		Anti-sickling activity /M.E (MNC en µg/mL)	Anti-sickling activity /A.E (MNC en µg/mL)
		ABTS (µg/mL)	DPPH (µg/mL)		
Rhizomes	12.82 ± 0.63	4.43 ± 0.18	15.63 ± 7.81	62.50	31.25

Roots	5.91 ± 0.87	4.66 ± 0.34	-	62.50	62.50
Leaves	4.31 ± 0.68	14.93 ± 4.30	54.83 ± 19.35	31.25	15.62
Petals	1.53 ± 0.28	24.27 ± 1.43	553.35 ± 250.15	62.50	62.50
Sepals	0.59 ± 0.30	46.24 ± 2.51	1129.80 ± 559.64	62.50	31.25

Legend : MNC: Minimum Normalization Concentration, M.E.: Methanol extract, E.A.: Aqueous extract, n= number of repetition of experiment

Correlations between the total polyphenol content, antioxidant activity and the anti-sickling activity of the extracts were demonstrated using the Pearson's test.

The analyses carried out revealed: a positive correlation between (i) the total polyphenol contents and the anti-sickling activity of the studied extracts, in particular methanol extracts ($r = 0.08328$) and (ii) between the antioxidant activity using ABTS and DPPH tests along with the anti-sickling activity of different extracts, in particular methanol extracts ($r = 0.1281$ and 0.4900) respectively). However, these correlations were not significant at 5%. The positive correlation between total polyphenolic contents and anti-sickling activity demonstrates that the polyphenolic compounds would contribute effectively at the same time to the inhibition of hemoglobin S polymerization and to erythrocyte sickling. However, the fact that this correlation (total polyphenol content and antioxidant activity of total methanol extracts) was not significant suggests that polyphenolic compounds act in synergy with other active ingredients. Tshilanda [2] and Ngbolua *et al.* [7], reported that anthocyanins have the ability to inhibit the polymerization of hemoglobin S by engaging in a reaction with this protein.

In addition, the analysis of the correlation using Pearson test did not reveal a correlation between the total polyphenol content or antioxidant activity using ABTS test and the anti-sickling potential of aqueous extracts ($r=-0.1740$, $r=-0.1481$ respectively). Although the correlation between the antioxidant activity of the extracts using DPPH test and the anti-sickling activity of the aqueous extracts was positive ($r=0.3114$). This would predict the involvement of the polyphenolic compounds of *C. longa* and their antioxidant potential on their anti-sickling potential; because the DPPH radical reacts exclusively with the polyphenolic compounds. However, in-depth and especially *in vivo* investigations are necessary to elucidate the implications of plant antioxidants on their anti-sickling potential.

Furthermore, the fact that the results of the present study showed a positive correlation between the antioxidant power of the extracts and their anti-sickling potential would reveal the involvement of the extract antioxidants in the normalization of erythrocytes *in vitro*. Au regard de ces résultats, Considering these findings, Curcuma is a good candidate for the management of sickle cell disease because it has both anti-sickling and antioxidant activities and because of the involvement of oxidative stress in the pathophysiology of sickle cell disease. Following Ngbolua *et al.* [7], anthocyanins are effective scavengers of free radicals and therefore potent inhibitors of lipid peroxidation and can, through their antioxidant properties, prevent the lysis of erythrocytes. A phytomedicine based on these compounds could ensure the protection of the membrane in sickle cell patients by supplementing the enzymatic defense systems of the erythrocyte, which are defective in sickle cell patients [1].

Several studies have reported that some polyphenolic compounds, notably curcumin extracted from *C. longa* rhizomes have the ability to trap free radicals (superoxide anions, hydroxyl radicals, H_2O_2 , nitrite radical, etc.), inhibit or regulate the inducible enzyme for the synthesis of nitrite oxide (NO), iNO-Synthetase (iNOS), prevent lipid peroxidation, allow the chelation of metals, maintain the activity of antioxidant enzymes such as superoxide dismutase, catalase, and glutathione peroxidase [11,18, 23, 84, 87, 92-94]. Hierso [1] reported that administration of N-acetylcysteine, a non-essential amino acid stimulating glutathione production, would significantly decrease the formation of sickle red blood cells and return them to their biconcave form.

Conclusion and perspectives

In the current study was highlighted the distribution of polyphenolic compounds in different organs (rhizome, root, leaf, petal and sepal) of *C. longa*, evaluated the anti-sickling activity of the total aqueous and methanol extracts of *C. longa* and their antioxidant potential. The findings showed that the polyphenolic compounds are not distributed in the same way in different organs (Rhizome, roots, leaves, floral parts) of *C. longa*. Total polyphenol content was higher in the rhizomes followed by roots, leaves, petals and sepals. The flavonoid content is higher in the leaves followed by rhizomes, roots, sepals and petals, while the anthocyanin content is stronger in the leaves then rhizomes and other parts are poor in anthocyanins. In addition, it was found that different organs of *C. longa* displayed an anti-sickling activity although leaf extracts showed a high activity. Aqueous extracts of rhizomes, leaves and sepals were more active than methanol extracts of the same organs. Moreover, it was also found that different organs of this plant do not all have a good antioxidant potential, rhizome and root extracts showed better antiradical activity using ABTS method compared to other organs, while rhizome and leaf extracts showed better antiradical activity using the DPPH test. At last, it was also found that there is a positive correlation

It was also demonstrated that there is a positive correlation between the total polyphenol content and the *in vitro* anti-sickling activity of the methanol extracts, and on the other hand between the *in vitro* antioxidant activity and anti-sickling activity. In view of the involvement of oxidative stress in the pathophysiology of sickle cell disease (sickling and haemolysis, vaso-occlusive attacks and other organ damage). it is necessary to carry out *in vivo* studies not only to prove the anti-sickling activity of the polyphenolic compounds of *C. longa* but also to highlight the involvement of their antioxidant potential in the pathophysiology of sickle cell disease.

Studies on the anti-sickling activity total aqueous extracts from different parts of *C. longa* or those analyzing the correlation between total polyphenolic content and/or the correlation between the antioxidant and anti-sickling activities have not yet been reported in the literature. Further studies are required in order to (i) better understand the roles of plant antioxidants (especially polyphenols) in improving the clinical signs of sickle cell disease and their *in vivo* mechanisms of action and (ii) identify the active ingredients responsible for the anti-sickling activity.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- [1]. Hierso R. Implication du stress oxydant dans la pathologie de la drépanocytose : crises vaso-occlusives, taux d'anticorps anti-bande 3 et oxydation du globule rouge. Thèse, Faculté de Sciences exactes et naturelles. Université des Antilles et de Guyane, 2015, 185 pp.
- [2]. Tshilanda DD. Etude phytochimique et activité antidrépanocytaire de trois espèces du genre *Ocimum* : *Ocimum basilicum* L., *O. canum* Sims, *O. gratissimum* L. (Lamiaceae). Thèse de doctorat, Université de Kinshasa, Faculté des Sciences, Département de chimie, 2016, 274 pp.
- [3]. Janky E, Etienne-Julan M, Kadhel PH, Leborgne-Samuel Y, Melki E. Drépanocytose et grossesse. Extrait des mises à jour en Gynécologie et obstétrique, 2006, Tome XXX.
- [4]. Ravelojaona M. Analyse histologique des répercussions musculaires, structurales, énergétiques et microvasculaires chez des hommes et des femmes drépanocytaires. Thèse de doctorat, Université Jean Monnet Saint Etienne, 2015, 206 pp.
- [5]. Mulungulu DN, Mpiana PT, Mbayo M, Tshisand PP, Badibanga ML. Etude de l'activité antidrépanocytaire et de la thermo-dégradation des extraits bruts aqueux, méthanoliques et éthanoliques d'*Ipomoea batatas*, une plante comestible à vertu thérapeutiques. *Int. J. Innov. & Applied Studies*, 2015; 11(3): 684-690. (<http://www.ijias.issr-journals.org>).
- [6]. Mpiana PT, Mudogo V, Tshibangu DST, Shetonde OM, Ngbolua KN, Mbala BM. Antisickling activity of some Congolese plants, in: Drug discovery from African flora, The 12th Symposium of the Natural Product Research Network for Eastern and Central Africa, July 22-26; University of Makerere, Kampala, Uganda, 2007d, p.45 (PS-6).
- [7]. Ngbolua KN, Mudogo V, Mpiana PT, Malekani MJ, Rafatro H, Urvergratsimamanga S, Takoy L, Rakotoarimana H, Tshibangu DST. Evaluation de l'activité antidrépanocytaire et antipaludique

- de quelques taxons végétaux de la République Démocratique du Congo et de Madagascar. *J. Ethnopharmacologia*, 2013, 5:19 pp.
- [8]. Mbadiko CM, Ngbolua KNN, Mpiana PT, Tshilanda DD, Makengo GK, Pambu AL, Kemfine LL, Bamvingana CK, Bongo GN, Mbemba FT. Assessment of the antisickling activity of total methanol extracts from the rhizomes and roots of *C. longa* and the effect of photodegradation on the antisickling activity. *J. Adv. Med. & Life Sci.*, 2017, 5 (1):1-5.
- [9]. Mbadiko MC, Ngbolua KN, Mpiana PT, Mutambel ' H., Kikakedimau NR, Makengo KG, Pambu LA, Kemfine LL, Bongo NG, Mbemba FT. Phytochemical screening and assessment of anti-sickling activity of total methanol extracts of different organs of *Curcuma longa* L. (Zingiberaceae). *The Pharma. & Chem. J.*, 2017, 4(1):32-40.
- [10]. Hombourger C. Le Curcuma de l'épice au médicament. Thèse de Doctorat, Université Henri Poincaré-Nancy 1, Faculté de Pharmacie, 2010, 222 pp. (<http://www.culture.gouv.fr/culture/infos-pratique/droits/protection.htm>).
- [11]. Tilak JC, Meenal B, Hari M, Devasagayam TPA. Antioxidant Availability of Turmeric in Relation to its Medicinal and Culinary Uses. *Phytother. Res.*, 2004, 18:798–804.
- [12]. Shiyu L, Wei Y, Guangrui D, Ping W, Peiying Y, Bharat AB. Chemical Composition and Product Quality Control of Turmeric (*Curcuma longa* L.). *Pharma. Crops*, 2011, 2:28-35.
- [13]. Tanzeela N, Muneeb I, Ahmad R, Madiha S, Fatima I, Marwa W. Turmeric: A Promising Spice for Phytochemical and Antimicrobial Activities. *American-Eurasian J. Agric. & Environ. Sci.*, 2015 ; 15 (7): 1278-1288.
- [14]. Maizura M, Aminah A, Wan Aida WM. Total phenolic content and antioxidant activity of kesum (*Polygonum minus*), ginger (*Zingiber officinale*) and turmeric (*Curcuma longa*) extract. *Int. Food Res. J.*, 2011, 18: 529-534.
- [15]. Prafulla S, Arjun M, Vidya S. *Curcuma longa* Linn. A Phytochemical and Phytopharmacological Review. *Res. J. Pharmacogn. & Phytochemistry.*, 2013, 5(2):59-68.
- [16]. Roshan PY and Gaur T. Versatility of turmeric: A review the golden spice of life. *J. Pharmacogn. & Phytochemistry*, 2017, 6(1):41-46.
- [17]. Camatari FOS, Lopes KH, Valentim BI, Xavier JA, da Costa JG, Santana AEG, Goulart MOF. Antioxidant Potential of Flours from Cereals, Tubers, Beans and Seeds Chemical Profile of *Curcuma longa* Flour. *J. Nutr. Food Sci.*, 2016, 6(2):1-8..
- [18]. Chainani-Wu N. Safety and anti-inflammatory activity of curcumin: a component of turmeric (*Curcuma longa*). *J. Altern. Complement Med.*, 2003; 9(1):161-168.
- [19]. Harshal P, Mugdha K, Nilesh M, Pravin J, Kavita M. Phytochemical Evaluation and Curcumin Content Determination of Turmeric Rhizomes Collected From Bhandara District of Maharashtra (India). *J. Med. Chem.* 2014, 4(8):588-591.
- [20]. Mohammad NH, Aysha F, Nargis A, Shahnaz R, Shahadat Hossan MD, Mohammed R. Preliminary phytochemical screening, toxicity, antihyperglycemic and analgesic activity studies with *Curcuma longa* leaves. *World J. Pharmacy. & Pharm. Sci.*, 2014, 3 (9) : 81-91.
- [21]. Rohini S, Mehta A, Mehta P, Shukla K. Anthelmintic activity of rhizome extracts of *Curcuma longa* and *Zingiber officinale* (Zingiberaceae). *Int. J. Pharm. & Pharm Sci.*, 2011, 3(2):236-237.
- [22]. Jansen PCM, Grubben GJH, Cardon D. Ressources végétales de l'Afrique tropicale 3. Colorants et tanins. Wageningen. Pays-Bas : PROTA, 2005: 238 pp.
- [23]. Vaquier Anne RL. Intérêt d'un nouveau nutriment a visée anti-inflammatoire dans la gestion de troubles locomoteurs de cheval. Aspects bibliographiques et études cliniques. Ecole vétérinaire d'Alfort, Faculté de Médecine de Cretal, 2010, 199 pp.
- [24]. Ajay K, Monika S, Prem PS, Sandeep KS, Pratima R, Kapil PD. Antioxidant efficacy and curcumin content of turmeric (*Curcuma-longa* L.) Flower. *Int. J. Current Pharm. Res.*, 2016, 8(3):112-114.
- [25]. Kholi K, Ali J, Ansari MJ, Raheman Z. Curcumin: A natural anti-inflammatory agent. *Indian J. Pharmacol.*, 2005, 37 (3):141 – 147.
- [26]. Yan SW and Asmah R. Comparison of total phenolic contents and antioxidant activities of turmeric leaf, pandan leaf and torch ginger flower. *Int. Food Res. J.*, 2010, 17: 417-423
- [27]. Rafie H, Soheila H, Mohsen H, Mahnaz S, Roxanna H. Turmeric (*Curcuma longa*): from a variety of traditional medicinal applications to its novel roles as active antioxidant, anti-inflammatory, anti-cancer, and anti-diabetes. *Int. J. Pharmacol., Phytochemistry and Ethnomed.*, 2015, 1:37-45.
- [28]. Stanojevićv SJ, Stanojević PL, Cvetković JD, Danilović RB. Chemical composition, antioxidant and antimicrobial activity of the turmeric essential oil (*Curcuma longa* L.). *Adv. Techn.*, 2016, 4(2):19-25.

- [29]. Ajaiyeoba EO, Sama W, Essien EE, Olayemi IO, Ekundayo OT, Walker M, Setzer WN. Larvicidal Activity of Turmerone-Rich Essential Oils of *Curcuma longa* Leaf and Rhizome from Nigeria on *Anopheles gambiae*. *Pharma. I Biol.*, 2008 ; 46 (4) : 279–282.
- [30]. Fathima ND. Curcumin -Role in Cancer. *Pharm. Sci. & Res*, 2014; 6 (11):351-355.
- [31]. Essien EE, Schmidt NJ, Tameka M., Walker, William N., Setzer, Olusegun E. Chemotaxonomic Characterization and in-Vitro Antimicrobial and Cytotoxic Activities of the Leaf Essential Oil of *Curcuma longa* Grown in Southern Nigeria. *Med.*, 2015, 2(4), 340-349 (<https://www.ncbi.nlm.nih.gov>).
- [32]. Rajesh H, Rao SN, Rani NM, Prathima. K, Shetty, Rejeesh EP, Chandrashekar R. Phytochemical analysis of methanol extract of *Curcuma longa* Linn rhizome. *International J. Uni. Pharmacy & Bio Sci.*, 2013, 2(2):39-45.
- [33]. Savina P. Le Curcuma, un agent naturel de lutte contre le vieillissement cutané, Synthèse Bibliographique en Biologie et Biotechnologie. Master en Biologie et Gestion, Université de Rennes 1 UFR Sciences de la Vie et de l'Environnement, 2014, 34 pp.
- [34]. Arutselvi R, Balasaravanan T, Ponmurugan P, Saranji MN, Suresh P. Phytochemical screening and comparative study of antimicrobial activity of leaves and rhizomes of turmeric varieties. *Asian J. Plant Sci. & Res.*, 2012, 2(2):212-219.
- [35]. Seema PK. Comparative Study of Pharmacognostical and Preliminary Phytochemical Investigation of *Curcuma longa* Leaves and Rhizomes. *PharmaTutor*, 2016 ; 4(10): 31-36.
- [36]. Chairman K., Jayamala M, Vijila RC, Ranjit SAJA. Phytochemical Screening and Antimicrobial Activity of *Curcuma longa* Natural Dye. *Gen. Med.*, 2015, 3(2):1-4.
- [37]. Wagner DST, Yasmin A, Padoan GLG, Cola ZJ, Murray BI. Turmeric powder and its derivatives from *Curcuma longa* rhizomes: Insecticidal effects on cabbage looper and the role of synergists. *Scientific Reports*, 2016, 6 : 1-11.
- [38]. Virendra VP, Shalini T, Nirmala K, Chetan N, Kalpagam P. *In vitro* evaluation on antioxidant and antimicrobial activity of spice extracts of ginger, turmeric and garlic. *J. Pharmacogn. & Phytochemistry*, 2013, 2(3):143-148.
- [39]. Imma C, Vitulano MEC, D'Onofrio FLA. Extraction And Curcuminoids Activity From The Roots Of *Curcuma longa* By Rslde Using The Naviglio Extractor. *European Scientific J. Special Ed.*, 2016, 1857 – 7881.
- [40]. Anbujeba SJ, Suraj R, Renitha G, Anandrarajagopal K, Gnamakumari AAV, Promwichit P. *In vitro* antimicrobial evaluation of *Zingibera ffinale*, *Curcuma longa* and *Alpinia galangal* Extracts as Natural food preservatives. *Am. J. Food Tech.*, 2002, 4(5):192-200.
- [41]. Araujo CC, Leon LL, Biological activities of *Curcuma longa* L. *Mem Inst Oswaldo Cruz*, 2001, 96(5):723-728.
- [42]. Bruneton J. Pharmacognosie, phytochimie, plantes médicinales, 4er éd., Revue et Augmentée, T 6 et Doc. Editions médicales internationales, Paris. 2009 ; 1288 pp.
- [43]. Cheikh ZA. Etudes chimiques et biologiques d'*Aframomum sceptrum* (Zingibéraceae) et de la curcumine. Université paris-sud, Faculté de pharmacie de Chatnay-Melabry, 2013, 161 pp. (<http://www.tel.archives-ouvertes.fr>).
- [44]. Goel A, Kunnumkkara B, Aggarwal BB. Curcumin as «Curecumin»: From kitchen to clinic. *Biochem. Pharmacol.* 2008, 75: 787 – 809.
- [45]. Nada GAL, Talet YM, Nayyarjehan SA. Studies of the antibacterial effect of different fraction of *Curcuma longa* against urinary tract infections Isolation. *Pakistan J. Biol. Sci.*, 2004, 7(12):2055-2060. (<http://www.Scialert.net>).
- [46]. Niamsa N, Sittwet C. Antimicrobial activity of *Curcuma longa* Aqueous Extract. *J. Pharmacol. & Toxicol.*, 2009, 4 (4):173-177. (<http://www.Scialert.net>).
- [47]. Perry MC. Evaluation de la Curcumine comme agent anticancéreux dans le traitement des tumeurs cérébrales. Mémoire de Master en Chimie, Université du Québec, Montréal, 2008, 97 pp.
- [48]. Sivabalan S, Anuradha CV. A Comparative study on the antioxidant and glucose lowering effects of Curcumin Analogue through *in vitro* assay. *Int. J. Pharmacol.*, 2010, 6 (5):664-669. (<http://www.Scialert.net>).
- [49]. Gupta SK, Lather A, Jaiswall V, Garg S, Kumar JA. Phytochemistry of *Curcuma longa* - An Overview. *J. Pharm. & Biomed. Sci.*, 2010, 4 (1):1-9.
- [50]. Awasthi PK, Dixit SC. Chemical Composition of *Curcuma longa* Leaves and Rhizome Oil from the Plains of Northern India. *J. Young. Pharm.*, 2009, 1(4):312-316.
- [51]. Shawket DS. Screening the antibacterial potency of *Curcuma longa* L. essential oil extract against boils causing *Staphylococcus* species. *Int. J. Adv. Biol. Res.*, 2013, 3(4): 490-500.

- [52]. Supayang VP. Family Zingiberaceae compounds as Functional Antimicrobials, Antioxidants, and Antiradicals. *World J. Med. & Med. Sci.*, 2007, 1(2):227-240.
- [53]. Jurenka JS. Anti-inflammatory properties of curcumin, a major constituent of curcuma longa: a review of preclinical and clinical research. *Altern. Med. Rev.*, 2009, 14 (2):141 – 153.
- [54]. Ritwiz BS, Kusha SB, Ranjeet D, Ganesh N. *Curcuma longa* leaves exhibits a potential antioxidant, antibacterial and immuno-modulating properties. *Int. J. Phytomed.*, 2011, 3(2):270-278.
- [55]. Ancy AR, Salini P, Antony S. Phytochemical screening and comparative study of antimicrobial activity of leaves and rhizomes of turmeric varieties. *Int. Journal of Research in Plant Science*, 2017, 7(1): 7-11.
- [56]. Sawant RS, Godghate AG. Qualitative phytochemical screening of rhizomes Of *Curcuma longa* Linn. *Int. J. Sci., Env. & Tech.*, 2013, 2 (4): 634 – 641.
- [57]. Sindhu S, Chempakam B, Leela NK, Suseela BR. Chemoprevention by essential oil of turmeric leaves (*Curcuma longa* L.) on the growth of *Aspergillus flavus* and aflatoxin production. *Food Chem Toxicol*, 2011, 49(5):88-92.
- [58]. Lassa KL. Valorisation des produits forestiers non ligneux (PFNL) d'origine végétale vendus dans les marchés et leurs disponibilités dans les environs de la ville de Kinshasa : « cas de CADIM ». Mémoire de DEA, Faculté des Sciences, Université de Kinshasa, 2012, 285 pp.
- [59]. Kapepula PM, Mbombo PM, Franck F, Mouithys-Mickalad A, Mumba ND, Dibungi KP, Kabamba NN, Tits M, Frédérick M, Tamfum MJJ. Antioxidant potentiality of three herbal teas consumed in Bandundu rural areas of Congo. *Natural Product Res.*, 2017, 31(16):1-4.
- [60]. Adedapo AA, Jimoh FO, Koduru S, Afolayan AJ, Masika PJ. Antibacterial and antioxidant properties of the methanol extracts of the leaves and stems of *Calpurnia aurea*. *BMC Compl. & Altern. Med.*, 2008, 8(1):1–8.
- [61]. Mpiana PT, Mudogo V, Ngbolua KN, Tshibangu DST, Shetonde OM, Mbala MB. *In vitro* antisickling activity of anthocyanins from *Ocimum basilicum* L. (Lamiaceae). *Int. J. Pharmacol.*, 2007, 3 (4): 371-374.
- [62]. Mpiana PT, Mudogo V, Tshibangu DST, Ekitwa EK, Kanangila AB, Lumbu BS, Ngbolua KN, Atibu EN, Kakule MK. Antisickling activity of anthocyanins from *Bombax pentadrum*, *Ficus capensis* and *Ziziphus mucronata*: Photodegradation effect. *J. Ethnopharmacol.*, 2008, 120 :413-418.
- [63]. Mpiana PT, Kitwa EK, Kanangila AB, Tshibangu DS, Ngbolua KN, Atibu EK, Kalonda EM. Influence de la photodégradation sur l'activité antifalcémiant des anthocyanes extraits de quelques plantes de Lubumbashi (RD Congo). *Ann. Fac. Sci.*, 2009, 45-53.
- [64]. Mpiana PT, Mudogo V, Tshibangu DST, Ngbolua KN, Tshilanda DD, Atibu EK. Antisickling activity of anthocyanins of *Jatropha curcas* L. *Recent Progress Med. Plants*, 2009, 25:101-108.
- [65]. Mpiana PT, Mudogo V, Tshibangu DST, Ngbolua KN, Atibu EN, Kitwa EK, Kanangila AB. Antisickling activity of anthocyanins extracts of *Vigna unguiculata* (L.) Walp. *Recent Progress Med. Plants*, 2009, 25:91-98.
- [66]. Mpiana PT, Tshibangu DST, Shetonde OM, Ngbolua KN. *In vitro* antidrepanocytary activity (anti-sickle cell anemia) of some Congolese plants, *Phytomedicine*, 2007, 14:192-195.
- [67]. Mpiana PT, Mudogo V, Tshibangu DS, Ngbolua KN, Shetonde OM, Mangwala KP, Mavakala BK. *In vitro* antisickling activity of anthocyanins extract of a Congolese plant: *Alchornea cordifolia* M.Arg. *J. Med. Sci.*, 2007, 7(7)1182-1186.
- [68]. Mpiana PT, Makelele LK, Oleko RW, Bokota MT, Tshibangu DST, Ngbolua KN, Mbala MB, Atibu EK, Nsimba SM. Antisickling activity of medicinal plants used in the management of Sickle Cell Disease in Tshopo district, D.R.Congo, *Australian J. Med. Herbalism*, 2010, 22(4), 132-137.
- [69]. Tula YM. Activité anti drépanocytaire et stabilité des anthocyanes extraits des bulbes d'*Hypoxis angustifolia* et de feuilles d'*Ipomoea batatas*, Mémoire de licence, Département de Chimie, Faculté des Sciences, Université de Kinshasa, 2016.
- [70]. Gayatri N. and Rajani KS. Evaluation of antioxidant activity in ethanolic extracts of five *Curcuma* species. *Int. Res. J. Pharm.*, 2011, 2 (12):243-248.
- [71]. Chang CH, Lin HY, Chang CY and Liu YC. Comparisons on the antioxidant properties of fresh, freeze-dried and hot-air-dried tomatoes. *Journal of Food Eng.*, 2006, 77: 478-485.
- [72]. Nilanjana D, Purba M, Ajoy KG. Pharmacognostic and Phytochemical Evaluation of the Rhizomes of *Curcuma longa* Linn. *J. PharmaSciTech.*, 2013; 2(2):81-86.
- [73]. Neha B, Tiwari KL, and Jadhav SK. Comparative phytochemical screening of bioactive compounds in *Curcuma caesia* Roxb. and *Curcuma longa*. *Res. J. Med. Plant*, 2013, 7 (2):113-118.

- [74]. Pawar MA, Patil SS, Nagrik DM. Phytochemical and Physicochemical Investigation of *Curcuma longa* Linn Rhizome. *Int. J. Chem. & Phys. Sci.*, 2015, 4:458-463.
- [75]. Cousins M, Aldelberg J, Chen J, Rick F. Antioxidant capacity of fresh and dried rhizomes from four clones of turmeric (*Curcuma longa* L.) grown *in vitro*. *Industrial Crops & Pro.*, 2007, 2(2):129-135.
- [76]. Walker V. Impact de l'inoculation des microorganismes phytobénéfiques sur le métabolisme secondaire de *Zea mays*, Thèse de doctorat, Université Claude Bernard Lyon 1, 2013
- [77]. Yizhong C, Qiong L, Mei S, Harold C. Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Sci.*, 2003, 74: 2157–2184.
- [78]. Akram M, Shahab U, Afzal A. Khan U., Abdul H., Mohiuddin E., Asif M. *Curcuma longa* and curcumin: A review article. *J. Biol. – Plant Biol.*, 2011; 55 (2). 65–70.
- [79]. Pratap SR and Jain DA. Evaluation of antimicrobial activity of curcuminoids isolated from turmeric. *Int. J. of Pharm. & Life Sci.*, 2012, 3 (1):1368-1376.
- [80]. Lawhavinit OA, Ngampong K. and Boonsong K. Antimicrobial Activity of Curcuminoids from *Curcuma longa* L. on Pathogenic Bacteria of Shrimp and Chicken. *Kasetsart J.*, 2010, 44:364 – 371.
- [81]. Shanmugam MK, Grishma R, Madhu MK, Arfuso F, Arunachalam Ci, Zayed ME, Sulaiman AA, Tan BKH, Kumar AP and Gautam S. The Multifaceted Role of Curcumin in Cancer Prevention and Treatment. *Mol. J.*, 2015, 20:2728-2769.
- [82]. Virendra PV, Shalini T, Nirmala K, Chetan N, Kalpagam P. *In vitro* evaluation on antioxidant and antimicrobial activity of spice extracts of ginger, turmeric and garlic. *J. of Pharmagn. and Phytochem.istry*, 2013, 2(3): 143-148.
- [83]. Sachin K, Narendra NS, Arun S, Neelakshi S, Raman KS. *Use of Curcuma longa* L. extract to stain various tissue samples for histological studies. *Ayu Journal*, 2014, 35(4):447-451.
- [84]. Jasim HN, Ali ATA and Al-Amiery AA H. Study the anticancer activities of ethanolic curcumin extract. *Afr. J. Pure & Applied Chem.*, 2010, 4(5):68-73.
- [85]. Seggani S. et Boukehil H. Corrélation entre le contenu polyphénolique et l'activité antioxydante et antimicrobienne *in vitro* de *Curcuma longa* L., Mémoire de Master, Université des Frères Mentouri Constantine, Faculté des Sciences de la Nature et de la Vie, République Algérienne Démocratique et Populaire, 2017, 108 pp.
- [86]. Boutamani M. Etude de la variation du rendement et de la composition chimique du *Curcuma longa* et *Myristica fragrans* en fonction du temps et de la technique utilisée, Mémoire de Master, Université des Sciences et de la technologie Houari Boumediene Alger, Chimie de Médicament, 2013 (Mémoire Online.net).
- [87]. Omodamiro OD and Ekeleme CM. Comparative study of *in vitro* antioxidant and antimicrobial activities of *Piper guineense*, *Curcuma longa*, *Gongronema latifolium*, *Allium sativum*, *Ocimum gratissimum*. *World J. Med. & Med. Sci.*, 2013, 1 (4) : 51-69.
- [88]. Shan, B., Cai, Y. Z., Sun, M. and Corke, H. Antioxidant capacity of 26 spice extracts and characterization of their phenolic constituents. *J. Agri. Food Chem.*, 2005, 53: 7749–7759.
- [89]. Wu CQ, Chen F, Wang X, Kim HJ, He GQ, Haley-Zitlin V, Huang G. Antioxidant constituents in feverfew (*Tanacetum parthenium*) extract and their chromatographic quantification. *Food Chem.*, 2006, 96:220–227.
- [90]. Velioğlu YS, Mazza G, Gao L. and Oomah BD. Antioxidant Activity and Total Phenolics in Selected Fruits, Vegetables, and Grain Products. *J. Agri. Food Chem.*, 46:4113-4117.
- [91]. Kaur C and Kapoor HC. Antioxidant activity and total phenolic content of some Asian vegetables. *Int. J. Food Sci Tech.*, 2002, 37: 153-161.
- [92]. Jayaprakasha GK, Jagan Mohan RAOL, Sakariah KK. Chemistry and biological activities of *C. longa*. *Trends Food Sci. Tech.*, 2005, 16 (12): 533 – 548.
- [93]. Anand P, Kunnumakkara AB, Newman RA, Aggarwal BB. Bioavailability of curcumin: problems and promises. *Mol. Pharm.*, 2007; 4(6): 807 – 818.
- [94]. Camacho-Barquero L, Villegas I, Sanchez-Calvo JM, Talero E, Sanchez-Fidalgo S, Motivla V, Catalina A. Curcumin, a *Curcuma longa* constituent, acts on MAPK p38 pathway modulating COX-2 and iNOS expression in chronic experimental colitis. *Int. Immunol.*, 2007, 7(3): 333 -342.