

Level of Interferon - gamma, Interlukin – 6, frequency of
Monocyte and Lymphocyte subsets among newly diagnosed
Tuberculosis patients in northwest Ethiopia

Abstracts

Background

Tuberculosis is one of the fundamental health problems in the world. During TB infection immune cells and cytokines play the major role for the pathogenesis of the disease and also for host defence. Cells of immune system play a great role to decrease the progress of the disease with the coordination of cytokines. The aim of our study was determining the level of immune cell and certain cytokine in TB patients and apparently health controls.

Methods

A comparative cross-sectional study design was conducted among newly diagnosed TB patients in Northwest Ethiopia. Blood samples were collected from all study participants. Serum level of interferon-gamma and interleukin-6 were determined by using an Enzyme Linked Immunosorbent Assay (Ready-SET-Go! Kit), Frequency of Monocyte and T cell sub population were measured by using Flow cytometry by using BD Multi TEST™ reagent. Mann-Whitney U test and spearman correlation were applied for statistical test by using graph pad prism.

Result

A total of newly diagnosed smear positive TB patients and healthy blood donors were included in this study. From our study participants 19(23.75%) were females and 61(76.25%) were males with mean age of 30. The median level of IFN- γ and IL-6 in TB patients were higher when compared with controls ($P > 0.001$) and on the other hand the median level of CD4 and BMI were showed lower compared to controls ($P > 0.001$).

Conclusion

Our study showed high level of IFN- γ , IL-6 and low level of monocyte, CD4 and BIM in TB patients than control groups.

Key words; Tuberculosis, CD4T cell, IFN- γ and IL-6, Gondar, Ethiopia

INTRODUCTION

Tuberculosis (TB) continues to be the major health problems and is the main cause of death and morbidity globally. Based on the recent WHO report, there were 9.6 million new TB cases per year and close to 2 million deaths (1). Factors related to the environment, host and pathogen could play roles for TB susceptibility (2). Immune status of the patients play a great role in the progression of TB and known that host protective immune response against the pathogen has a critical role (3). The ability of the host to elicit potent cell mediated immune response the pillar of disease development and outcomes associated TB. Macrophage activation, cytokine production and other immune cells play roles in TB immunity and determine the outcome of the disease (4, 5).

T-helper 1(Th1) and T-helper 2 (Th2) immune responses are the major arms of immune responses for humeral and cell mediated immune system, respectively. The induction of the Th1 type immune response very crucial to combat TB infection and characterized by secretion of interferon-gamma (IFN- γ). Cytokine such as IFN- γ , Tumor necrosis factor – alpha (TNF- α), IL-12 and Interleukin-6 (IL-6) are responsible chemicals to build the proactive immunity against TB pathogens during early stage of the infection. IFN- γ has versatile roles including facilitating the intracellular killing of the TB bacilli by macrophages (6, 7). It has a great role for the function and maturation of immune cells, and it stimulates production of other cytokines and activation factors to augment the function of innate immune system during infections (8, 9). In addition to IFN- γ , other the inflammatory cytokines like IL-6, play pivotal roles in TB immunity. The secretion of IFN- γ for the activation of macrophage by CD4 cells during TB infection is stimulated by the production of IL- 6 (10). IL-6 is a cytokine known by kind of pleiotrophic nature and produced by many cells including T and B-lymphocytes. It plays a great role in the synthesis of C-reactive protein in various stimuli (11, 12) and also has been revealed that it has a relevant role in the pathogenesis of tuberculosis like the secretion of IFN- γ during infection (10, 13).

Therefore, the present study aims to assess the levels of these two essential cytokines which are greatly needed to combat TB infection immune cells which are T cells and monocyte. Thus, the level of these two cytokines and cells among active TB patients will be compared to the level from apparently healthy control.

MATERIAL AND METHODS

Study design and area

A cross-sectional study conducted in newly diagnosed TB patients at the Gondar of University referral hospital from February to May 2014. Gondar University hospital is a tertiary level teaching and referral hospital with 450 beds for inpatients and rendering referral health services for over 5 million inhabitants in North-West Ethiopia. The hospital provides inpatient and outpatient services to the population in the surrounding area of Gondar town and the adjacent regions. The hospital has TB clinic where TB patients are getting their medication and further assessment during follow up period. For comparison purpose, apparently healthy controls were enrolled at the Gondar University Blood Bank.

Data collection

Socio-demographic data were collected using a structured and pretested questionnaire. Whole blood was collected from the study participant using serum separator (14) tubes and EDTA for measurement of cytokines using ELISA technique and the remaining one was used for blood cell count determination, respectively.

Serum level IFN- γ and IL-6 measurement

The levels of IL-6, and IFN- γ were measured using ELISA Ready-SET-Go! Kit. Anti-human cytokines (IL-6 and IFN- γ) capture antibodies (eBioscience) was coated in ELISA (Coat corning costar 9018) plates with 100 μ l/per well in 1x coating buffer and the plates were sealed and incubated overnight at 2-8°C. On the following day, the plates were aspirated and washed three times using 250 μ l wash buffer (1x PBS, 0.05% Tween 20). The standards (human cytokines lyophilized standard) were reconstituted and diluted according to the given instruction. A twofold serial dilution of the top standards with 100 μ l were performed in duplicates to make a standard curve for 8 points and also 100 μ l supernatant of samples were added to incubate at room temperature for 2 hours. After 3-5 washes, 100 μ l/well of detection antibody (anti human cytokine biotin) was added and incubated at room temperature for 1 hour. After 3-5 washes, 100 μ l/well of Avidin-HRP was added and incubated at room temperature for 30 minutes. Finally, 1M H₃PO₄ or 2N H₂SO₄ stop solution was added to stop the reaction, and the plate was read at 450nm length using ELISA reader. The analytical sensitivity and other technical protocols were followed as per the manufacturer guideline.

Cell counting using Flow cytometry

T-lymphocyte sub-populations were characterized using Multi TEST™ fluorescent labelled monoclonal antibodies reagents against surface CD markers (CD₃-fluorescein isothiocyanate (FITC), CD₈- phycoerythrin (PE) and CD₄-allophycocyanin (APC)) after erythrocyte has been lysed. FACS Calibur flow cytometer (Becton Dickinson Biosciences, San Jose, CA, USA) was used for acquisition and analysis of T-lymphocyte sub-population by using Multitest software (Becton Dickinson), and a minimum of 15, 000 events were captured during acquisition. Monocyte cell count done by coulter counter machine.

Statistical analysis

Data were analyzed using graph pad prism. Data were reported as frequency and percentage for categorical variables and mean and standard deviation (SD). Mann-Whitney U test was used to compare the cytokine levels between TB patients and healthy controls. Spearman correlation was used to correlate serum IFN- γ and IL-6 cytokines with peripheral CD4+, CD8+ and Monocyte cells. Statistical significance was considered at 95 % level of confidence and P value less than 0.05.

RESULTS

A total of 40 newly diagnosed TB patients and equivalent number of apparently healthy controls were included in the study. The mean age for the TB patients and the controls were 34 years (range 18 to 57) and 26 Years (range: 20 to 42), respectively. From the total TB patients 27.5% (11/40) were females; 22.5% (9/40) were also females from the control group. From the total cases 70% (28/40) were smear positive pulmonary TB and the remaining were extra pulmonary TB. The BIM of the study participants were, 17.98 for Tb patients and 20.64 for controls. (Table.1)

IFN-gamma, IL-6, and peripheral T-cell subpopulation and Monocyte

The level IFN- γ and IL-6 were significantly higher among active TB patients compared to healthy controls (median of 72.8 Pg/ml versus 44.0 pg/ml, $p < 0.0001$ for IFN-gamma; and 27.4 pg/ml versus 6.0pg/ml, $p < 0.0001$ for IL-6) (table 1). Similarly, significant differences ($p < 0.0001$) and monocyte ($p = 0.018$) was obtained between active TB patients and healthy controls. (Table 1) & (Fig 1).

Correlation of serum IFN- γ and IL-6 cytokines with peripheral CD4⁺, CD8⁺ and Monocyte cells,

By using spearman correlation among cytokine and cells, IL-6 and CD4⁺ showed weak negative correlation ($r = -0.4929$; $p = 0.001$) but monocyte and CD8⁺ cell showed weak positive correlation ($r = 0.3962$; $p = 0.0167$). (Fig 2)

DISCUSSION

In our study, the level of CD4 cell count is significantly higher in the controls when compared with that of newly diagnosed TB-patients. This is in agreement with previously literature. Evidence also shown that a low CD4 cell count associated with TB disease and CD4 cell count is substantially low in more advanced disease among HIV-negative TB patients and more advanced disease was shown CD4 lymphocytopenia (14). Because of protective immune response against this pathogen mediated by cellular immunity mediated by Th1 cells. This leads the patient for the progression of tuberculosis (3). More over most of TB cases show low BMI in our study, which may play a great role for the decreasing of CD4 counts and their functionality by reducing the transportation of different co-factors and elements than controls with normal BMI value. This Low level of CD4 cells play a great for the pathogenesis of TB disease, because it may be due to the low respond in delayed type immune response and due to low level of cell frequency may lead to decrease in granuloma formation, which is fundamental for proper immune response.

IFN- γ is one of the Th1 cytokine where identified as the most important agents of antimycobacterial cytokine (15) by activate macrophage to enhance intracellular killing (16) and enhance the production other cytokine (17). In our study the level of IFN- γ was significantly higher in TB patients compared with the control groups. The finding is consistent with other similar studies on TB (18-20) and the level of this cytokine is decrease through and after treatment (20) The probable reason why INF- γ is higher in TB patients while CD4 cell are low is that, may be due to IFN- γ comes out from both local production and spill over of it from the activated lymphocytes sequestered at the site of MTB infection. This condition can able to contribute to the cytokine not be able to elicit downstream events involving in effective activation of macrophages and intracellular killing of MTB. All these conditions play a great role for the pathogenesis of MTB infections.

The level of IL- 6 in TB patients a higher than that of the controls, other similar studies had the result which is in line with our findings, done in humans (21, 22). On the other hand also study showed that there is increased IL- 6 in active TB disease condition when compared with that of the latent condition of TB infection (23). In addition study showed that the concentration of IL-6 very high in TB patients with pulmonary cavities than without cavities. This great indication that IL-6 plays it own role for the TB disease outcome (24). High level of IL-6 during TB infection can inhibit the type 1 interferon signalling on macrophage and

consequently lead to the disease progression (25). On the other hand of IL-6 level correlated with disease severity, nutritional status has a great contribution for the incereasement of IL-6 concentration (24), our studies support this because of the level of BMI in TB patients were significantly lower than that on health controls.

The level of monocyte in the blood of TB patients is higher than that of the controls and show significant association, one of the probable reason that the increase of the monocyte during TB infection is that during TB infection there is a production of pro-inflammatory cytokines and chemo attractant factors that direct the monocyte from the bone marrow to lymph node and other infected tissue to act like macrophages. At this condition there may be an increasment of monocyte in the vascular. On the other hand cytokines that is produced by infected macrophages may have an impact on monocyte not to change in to macrophages to go to the infected area. This high frequency of monocyet in TB patients may also contribute for the increasement of IL- 6 in TB patients; they have high level than of the controls.

CONCLUSION

In our study patients with tuberculosis showed higher level of IFN- γ , IL-6 and monocyte with lower level of CD4 and BMI than the controls. This may indicate the progression of the disease. Future studies are required for further describing the mechanisms of this value alteration during TB disease and for better clinical application of these parameters.

Abbreviations

BMI: body mass index

CD: cluster of differentiation

EDTA : Ethylenediaminetetraacetic acid

ELISA: Enzyme linked immunosorbant assay

IL: Interlukine

IFN - γ : Interferon gamma

TB: Tuberculosis

TNF- α : Tumor necrosis factor alpha

SD: Standard deviation

WHO: world health organization

Declarations

Aknowledgement

We would like to thank Yohans Sink, Eden W/gerima, Debaka Belet and Samirawit Sisay in the form university of Gondar hospital HIV laboratory. And also Esay Derebe Asnkew Belet Tigist, Koku Sisay, Nebyu Mekonn from University of Gondar hospital TB clinic and laboratory for their indispensable cooperation. Finally we would like to acknowledge University of department of Immunology and molecular biology for great cooperation to finalize this work.

Funding

Not applicable

Availability of data and materials

The datasets supporting the conclusions of this article are included within the article.

Authors' contributions

ML & EA: Conceived and designed the experiments; ML, MW & BM, carried out data collection and laboratory analysis; EA supervised the data collection and laboratory analysis; ML & SY ;analyzed the data and prepared the first manuscript draft EA , MW & ML; reviewed the draft. All authors read and approved the final manuscript.

Competing interests

The authors declare that there is no competing interest with regard to the present study.

Consent for publication

Not applicable.

Ethics approval and consent to participate

The study was approved by the ethical committee of School of Biomedical and Laboratory Sciences, University of Gondar. In addition, written informed consent was obtained from each participant.





REFERENCE

1. Organization WH. Global tuberculosis report 2015: World Health Organization; 2015.
2. McShane H. Susceptibility to tuberculosis—the importance of the pathogen as well as the host. *Clinical & Experimental Immunology*. 2003;133(1):20-1.
3. Kinjo Y, Kawakami K, Uezu K, Yara S, Miyagi K, Koguchi Y, et al. Contribution of IL-18 to Th1 response and host defense against infection by *Mycobacterium tuberculosis*: a comparative study with IL-12p40. *The Journal of Immunology*. 2002;169(1):323-9.
4. Munk M, Emoto M. Functions of T-cell subsets and cytokines in mycobacterial infections. *The European respiratory journal Supplement*. 1995;20:668s-75s.
5. Jo E-K, Park J-K, Dockrell HM. Dynamics of cytokine generation in patients with active pulmonary tuberculosis. *Current opinion in infectious diseases*. 2003;16(3):205-10.
6. Kaufmann SH. Is the development of a new tuberculosis vaccine possible? *Nature medicine*. 2000;6(9):955-60.
7. Mosmann T, Coffman R. TH1 and TH2 cells: different patterns of lymphokine secretion lead to different functional properties. *Annual review of immunology*. 1989;7(1):145-73.
8. Miller CH, Maher SG, Young HA. Clinical Use of Interferon- γ . *Annals of the New York Academy of Sciences*. 2009;1182(1):69-79.
9. Mootoo A, Stylianou E, Arias MA, Reljic R. TNF- α in Tuberculosis: A Cytokine with a Split Personality. *Inflammation & Allergy-Drug Targets (Formerly Current Drug Targets-Inflammation & Allergy)*. 2009;8(1):53-62.
10. Saunders BM, Frank AA, Orme IM, Cooper AM. Interleukin-6 Induces Early Gamma Interferon Production in the Infected Lung but Is Not Required for Generation of Specific Immunity to *Mycobacterium tuberculosis* Infection. *Infection and immunity*. 2000;68(6):3322-6.
11. Papanicolaou DA, Wilder RL, Manolagas SC, Chrousos GP. The pathophysiologic roles of interleukin-6 in human disease. *Annals of internal medicine*. 1998;128(2):127-37.
12. Gabay C, Kushner I. Acute-phase proteins and other systemic responses to inflammation. *New England journal of medicine*. 1999;340(6):448-54.
13. Ladel CH, Blum C, Dreher A, Reifenberg K, Kopf M, Kaufmann S. Lethal tuberculosis in interleukin-6-deficient mutant mice. *Infection and immunity*. 1997;65(11):4843-9.

14. Skogmar S, Schön T, Balcha TT, Jemal ZH, Tibesso G, Björk J, et al. CD4 cell levels during treatment for tuberculosis (TB) in Ethiopian adults and clinical markers associated with CD4 lymphocytopenia. *PLoS One*. 2013;8(12):e83270.
15. Lin PL, Flynn JL. Understanding latent tuberculosis: a moving target. *The Journal of Immunology*. 2010;185(1):15-22.
16. Cooper AM, Khader SA. The role of cytokines in the initiation, expansion, and control of cellular immunity to tuberculosis. *Immunological reviews*. 2008;226(1):191-204.
17. Fulton SA, Johnsen JM, Wolf SF, Sieburth DS, Boom WH. Interleukin-12 production by human monocytes infected with *Mycobacterium tuberculosis*: role of phagocytosis. *Infection and Immunity*. 1996;64(7):2523-31.
18. Liu Y, Janeway CA. Interferon gamma plays a critical role in induced cell death of effector T cell: a possible third mechanism of self-tolerance. *Journal of Experimental Medicine*. 1990;172(6):1735-9.
19. Katti MK. Assessment of serum IL-1, IL-2 and IFN- γ levels in untreated pulmonary tuberculosis patients: role in pathogenesis. *Archives of medical research*. 2011;42(3):199-201.
20. Lee SW, Lee C-T, Yim J-J. Serial interferon-gamma release assays during treatment of active tuberculosis in young adults. *BMC infectious diseases*. 2010;10(1):300.
21. Lopes FHA, Assis LCd, Pires Neto RdJ, Botelho KP, Sá KM, Frota CC, et al. Serum levels of interleukin-6 in contacts of active pulmonary tuberculosis. *Jornal Brasileiro de Patologia e Medicina Laboratorial*. 2013;49(6):410-4.
22. Correia J, Freitas M, Queiroz J, PereiraPerrin M, Cavadas B. Interleukin-6 blood levels in sensitive and multiresistant tuberculosis. *Infection*. 2009;37(2):138-41.
23. Djoba Siawaya J, Beyers N, Van Helden P, Walzl G. Differential cytokine secretion and early treatment response in patients with pulmonary tuberculosis. *Clinical & Experimental Immunology*. 2009;156(1):69-77.
24. Karyadi E, Dolmans WM, West CE, Van Crevel R, Nelwan RH, Amin Z, et al. Cytokines related to nutritional status in patients with untreated pulmonary tuberculosis in Indonesia. *Asia Pacific journal of clinical nutrition*. 2007;16(2):218-26.
25. Martinez AN, Mehra S, Kaushal D. Role of interleukin 6 in innate immunity to *Mycobacterium tuberculosis* infection. *The Journal of infectious diseases*. 2013;207(8):1253-61.

Table 1. Level of IFN- γ , IL- 6, CD4T cell, CD8 T cell and Monocyte cell in TB patients and controls

Parameters	TB patients	Control	p-value
IFN- γ (pg/ μ l) (median \pm SD)	72.8	44.0	< 0.0001
IL-6 (pg/ μ l) (median \pm SD)	13.45	4.05	< 0.0001
CD4 cells($\times 10^3$ / μ l)	565.5	857.0	< 0.0001
CD8 cells($\times 10^3$ / μ l)	539.5	551.5	< 0.4162
Monocyte ($\times 10^3$ / μ l)	400.0	300.0	< 0.0193











Fig. 1 Serum level of IFN-gamma, IL-6, and measurement of peripheral blood CD4+ cells, CD8+ cells and Monocytes among newly diagnosed TB patients (n=40) and apparently healthy controls (n=40)






Fig.2 Correlation of serum level cytokine with peripheral cell in newly diagnosed TB patients (A) IFN- γ with CD4⁺ cells, (B) IFN- γ with IL-6, (C) IFN- γ with monocyte , (D) IFN- γ with CD8⁺ , (E) IL-6 with CD4⁺ cells, (F) IL-6 with CD8⁺ cells (G) IL-6 with monocyte cells and (H) CD8+ cell with monocyte cells.