

PREVALENCE OF INTESTINAL PARASITIC INFECTIONS AMONG PATIENTS ATTENDING USMANU DANFODIYO UNIVERSITY TEACHING HOSPITAL, SOKOTO, NIGERIA

ABSTRACT

Background: Intestinal parasitic infection is one of the major health issue in developing countries particularly in Sub-Saharan Africa. It has been estimated to affect about 3.5 billion people globally and 450 million people are thought to be ill as a result of such infections, the majority being children.

Aims: The study is aimed at determining the prevalence and associated risk factors of intestinal parasitic infections among patients attending Usmanu Danfodiyo University Teaching Hospital, Sokoto, Nigeria.

Study Design: This was a cross-sectional, descriptive study.

Place and Duration of Study: This study was conducted among patients attending Usmanu Danfodiyo University, Teaching Hospital, Sokoto, Sokoto state, between May to November 2017.

Methodology: A total of 245 participants were enrolled in the study. Standard parasitological examination was carried out on stool samples using microscopic examination.

Results: Finding revealed that 29 (11.8%) were positive for intestinal parasitic infections. Males recorded higher prevalence than the females with 19 (7.8%) and 10 (4.1%), respectively. Intestinal parasites continue to remain a serious public health problem in North-western Nigeria.

Conclusion: Low level of education, occupational status, poor water supply were among the significant risk factors for these infections. Creating awareness, level of sanitation, water supply and deworming programme among school children will reduce prevalence and intensity of parasitic infections among the study community.

Keywords: Prevalence study, Intestinal parasitic infection, UDUTH, Sokoto State, Nigeria.

1.0 INTRODUCTION

Intestinal parasitic infection is one of the major public health burdens in developing countries particularly in Sub-Saharan Africa. It has been found to affect about 3.5 billion people globally and 450 million people are thought to be ill as a result of such infections, the majority being children [1]. In

Comment [C1]: was

Formatted: Highlight

Formatted: Font color: Red

Formatted: Highlight

Comment [C2]: Should not be included in the results. If you wish, describe in "conclusion". In your conclusion section, you said that the prevalence was low!!!

Formatted: Font color: Red, Strikethrough

Comment [C3]: Rewrite (or) rephrase!

35 Nigeria, intestinal helminthes infections have continued to prevail because of poor standards of living,
36 poor environmental sanitation and ignorance of simple health promoting behaviours[2,3].Intestinal
37 helminthes infections are most common in school age children and they tend to occur in high intensity
38 in this age group [4,5,6].

39 .

40 These infections have been associated with an increased risk for nutritional anaemias, protein energy
41 malnutrition, growth deficits in children, physical weakness and low educational performance of
42 school children [7,8] and also causing high morbidity and mortality rate [9].

43 Parasitic infections are governed by behavioural factors, biological, environmental, socioeconomic
44 and health systems factors. Local conditions such as quality of domestic and village infrastructure;
45 economic factors such as monthly income, employment and occupation and social factors such as
46 education influence the risk of infection, disease transmission and associated morbidity and mortality
47 [10,11]. These infections are more prevalent among the poor segments of the population. They are
48 closely associated with low household income, poor personal and environmental sanitation, and
49 overcrowding, limited access to clean water, tropical climate and low altitude. Intestinal parasitic
50 infections such as amoebiasis, ascariasis, hookworm infection and trichiuriasis are among the ten
51 most common infections in the world [12].

52 There is dearth of information on the magnitude of intestinal parasitic infections and predictors among
53 patients attending Usmanu Danfodiyo University Teaching Hospital, Sokoto, Nigeria. Information
54 generated could be used for planning public health control programmes which is an important step for
55 initiation of treatment and prevention strategies as well as reducing morbidity and mortality due to
56 parasitic infections in the area.

57 MATERIALS AND METHODS

58 STUDY AREA

59 The study area is Usmanu Danfodiyo University Teaching Hospital, a tertiary health facility located in
60 Sokoto metropolis, the Sokoto State Capital. It serves as a referral centre for more than 10 million
61 people of the Nigerian States of Sokoto, Zamfara and Kebbi; and neighbouring Niger and Benin
62 Republic in the West African sub-region [13].

63 Sokoto State is located at the extreme part of North-Western Nigeria between longitude 3° and 7°
64 east and between latitude 10° and 14° north of the equator. It shares borders with Niger-Republic to
65 the North, Kebbi State to the South-West and Zamfara State to the East [13]. The state covers a total
66 land area of about 32,000 square kilometres and a population of 4,602,298 million based on 2013
67 projection [14]. Sokoto State has semi-arid climate and vegetation is largely Sudan Savannah with an
68 annual rainfall between 500 – 1300mm and temperature ranges between 150°C and over 400°C
69 during warm days [13].

Comment [C4]: was




Figure 1: Map of Sokoto State Showing Study Area [13].

73 **STUDY DESIGN**

74 This is a cross-sectional descriptive study that was carried out on 245 samples collected from UDUTH
75 Service laboratory, from May to November 2017.

76 **SUBJECT AND SELECTION**

77 The subjects were selected or recruited in UDUTH Service laboratory using systematic sampling
78 technique to recruit all patients that meet the inclusion criteria.

79 **Inclusion criteria:**

- 80 1. All patients with uncontaminated stool sample (formed, semi formed and unformed) were
81 recruited for study and
- 82 2. Patients who give their consent to participate in the study.

83 **Exclusion criteria:**

- 84 1. Patients with stool sample contaminated with urine or mixed with soil were excluded from the
85 study and
- 86 2. Patients who refuse to give consent in the study.

87 **SAMPLE SIZE**

88 The sample size was calculated using the formula outlined below [15]. Prevalence was set
89 at 17.5% [16].

90
$$n = \frac{(z-a)^2 (p) (1-p)}{d^2}$$

92
$$n = 221$$

93 Using an attrition rate of 10%, a total of 245 patients were selected.

94 **SAMPLING METHOD**

95 A systematic random sampling method was used to recruit the subjects (or) respondents. all patients
96 that meet the inclusion criteria.

97 The laboratory register had about six hundred patients (600) that submitted their stool for evaluation
98 in the previous year (January 2016 to December 2017). This was used to determine the sampling
99 frame.

100
$$K=N/n: 600/245 = 2.4 \sim 3$$

101 A sampling interval of 3 was achieved.

102 Using simple random sampling; the first patient was chosen between number 1 and 3 for the first
103 week of study.

104 For any randomly chosen numbered patient; thereafter a sampling interval of 3 would be used for the
105 subsequent patients that present themselves at the facility until the sample size was achieved.

106

Comment [C5]: Rewrite! (For example, systematic random sampling was used to select the participants!)

Comment [C6]: You don't need to describe both inclusion and exclusion criteria. Description of only one (either inclusion or exclusion) is enough!

Formatted: Highlight

Formatted: Font color: Red

Formatted: Font color: Red, Strikethrough

Formatted: Strikethrough

Formatted: Font color: Red, Strikethrough

107 **SAMPLE COLLECTION**

108 An approximate amount of 100g faeces was collected into clean, dry and screw cap, leak proof
109 containers.

110 **STUDY TOOL**

111 A structured questionnaire was administered to obtain patient information. It was structured into the
112 following subheadings; demographic information, socio-economic data, clinical history and laboratory
113 investigation. The questionnaire was pretested and validated at a similar site to the study area in the
114 state specialist hospital, Sokoto and corrections were made where necessary.

115 **SAMPLE PROCESSING**

116 The stool specimen was examined macroscopically for the presence of adult worms. The consistency,
117 color and presence of abnormal structures were recorded. It was also examined microscopically using
118 direct saline and wet iodine mount, and also formal-ether concentration method.
119

120 **DIRECT MICROSCOPIC EXAMINATION USING NORMAL SALINE AND IODINE** 121 **PREPARATION**

122 For each sample, normal saline mount and iodine mount was prepared on a slide and examined
123 microscopically at 10X and 40X for the presence of *Intestinal helminths*. Iodine preparation allows the
124 examination of the characteristics features of the protozoa and the identification of the *Entamoeba*
125 *histolytica/dispar* (*Entamoeba histolytica/dispar*) cyst from the commensal *Entamoeba coli*. [17]
126

127 **FORMALIN-ETHER SEDIMENTATION METHOD**

128 0.5g of faecal sample was added to a glass container containing 10mls of 10% formalin and then
129 mixed thoroughly. A Funnel was placed on a gauge and strain into a 15mls centrifuge tube and
130 centrifuged for 2minutes at 1500 rpm. Then the supernatant was discarded and the sediment was re-
131 suspended into 10mls of physiological saline, and centrifuged for 2 minutes at 1500 rpm. The
132 supernatant was discarded and the sediment re-suspended again in 7mls of 10% formaldehyde, 3mls
133 of ether (diethyl) was also added. The tube was closed with a glass stopper and mixed vigorously,
134 and then the stopper was removed and centrifuged for 2minutes at 1500 rpm. The supernatant was
135 poured out and the sediment carefully placed on a clean glass slide and covered with cover slip and
136 this was examined at x10 and x40 magnification on a light microscope [19].

137 **DATA MANAGEMENT**

138 Data were entered independently at two separate occasions using Microsoft Excel 2016. Double data
139 entry analysis was done to ensure data quality. statistical package for social sciences (SPSS) version
140 20 was used for the analysis. Categorical variable was assessed using Chi-square test to determine
141 the association. Simple and multiple logistic regression analysis was used to determine associated
142 risk factors of the infections. Values were considered statistically significant at $p < 0.05$.

143 **ETHICAL COSIDERATION**

144 Ethical clearance was obtained from the ethics and research committee of UDUTH, Sokoto and
145 consent was sought from the participants prior to sample collection.

146

147

Comment [C7]: Statistical (Capital S)

148 RESULT

149 The prevalence of intestinal parasites among the overall population studied was 11.8%. The
 150 highest prevalence of 6.2% was noted for Hookworm and *Ascaris lumbricoides* infection
 151 while the lowest prevalence was seen with *H. nana* (2.1%) as shown in Table 1. Of the total
 152 study subjects 160 (65.3%) were males and 85 (34.7%) were females. The males showed a
 153 higher prevalence of intestinal parasite infections of 11.9% (Table 2) than the females which
 154 showed a prevalence rate of (11.8%). However, this difference was ~~is~~ not statistically
 155 significant ($p>0.05$). The age range of 11-15 had the highest prevalence of parasitic infection
 156 with 2.9% and none was recorded among the age group 31 and above. There was a
 157 statistically significant difference ~~of worm infestation among between~~ age groups ($p=0.004$).

Comment [C8]: 2.0 %

Formatted: Font color: Red, Strikethrough

Formatted: Font color: Red, Strikethrough

158
 159 Prevalence of intestinal parasitic infection among study population based on water source
 160 shows that those that consume river/stream water 13 (5.3%) have highest risk of intestinal
 161 parasitic infection, followed by those that drink other source of water with prevalence of 8
 162 (3.3%) then followed by those that drink well water with 4 (1.6%), Tap water 3 (1.2) and
 163 lastly those that consumed sachet water have the lowest prevalence of 1 (0.4%). Comparing
 164 the different prevalence rates in relation to intestinal parasites by water source is statistically
 165 significant ($p=0.001$).

166 The distribution of intestinal parasitic infection among study population based on frequency
 167 of eating vegetables, walking bare footed, type of toilet facility, occupation, educational level
 168 etc. are shown in Table 4.

169

170 **Table 1: Prevalence of intestinal parasitic Infection**

171

172

Parasites	Frequency (n=245)	Prevalence (%)
-----------	-------------------	----------------

173

174 *Hook worm and* 15 6.1

175 *Ascaris Lumbricoides*

176

177 *G. lamblia and* 9 3.7

178 *Entamoeba. histolytica*

179 *H. nana* 5 2.0

180 Total 29 11.8

181

182

183

184

185

186 TABLE 2: Showing Prevalence in relation to gender.

Gender	No examined (%)	Intestinal Parasites		p-value
		Infected (%)	Not Infected (%)	
Male	160 (65.3)	19 (7.8)	141(57.6)	0.413
Female	85 (34.7)	10(4.1)	75 (30.6)	
Total	245	29	216	

Comment [C9]: Format of the Tables 2, 3 and 4 should be the same (uniform/consistent)!!!

Comment [C10]: Calculate row percent. E.g., 19 out of 160 (11.9%) among males whereas 10 out of 85 (11.8%) among females.

193
194 TABLE 3: Showing age distribution

Ages	No examined (%)	Intestinal parasites		p-value
		Infected (%)	Not Infected (%)	
0-5	33(13.5)	5(2.0)	28(11.4)	0.004*
6-10	19(7.8)	4(1.6)	15(6.1)	
11-15	19(7.8)	7(2.9)	12(4.9)	
16-20	57(23.3)	6(2.4)	51(20.8)	
21-25	83(33.9)	5(2.0)	78(31.8)	
26-30	13(5.3)	2(0.8)	11(4.5)	
≥ 31	21(8.6)	0	21(8.6)	
Total	245	29	216	

Comment [C11]: Showing Prevalence in relation to age-group

Comment [C12]: Must be row percent! Calculate row percent (prevalence) in each age-group!

Table 4: Distribution of intestinal parasitic infection among study population with respect to some sociodemographic characteristics.

Variables	Intestinal parasites					
	Infected		Not infected		Total	
<i>p</i> -value ^a						
	n	%	n	%	N	
Educational status						
None	7	(2.9)	77	(31.4)	84	0.021 *
Informal	2	(0.8)	14	(5.7)	16	
Primary	3	(1.2)	27	(11.0)	30	
Secondary	11	(4.5)	29	(11.8)	40	
Tertiary	6	(2.4)	69	(28.1)	75	
Occupation						
Business	3	(1.2)	23	(9.4)	26	0.905
Farming	1	(0.4)	14	(5.7)	15	
Civil servant	7	(2.9)	40	(16.3)	47	
Unemployed	4	(1.6)	37	(15.1)	41	
Student	14	(5.7)	102	(41.6)	116	
Water source						
Tap water	3	(1.2)	59	(24.1)	62	0.001 *
Well water	4	(1.6)	33	(13.5)	37	
River/stream	13	(5.3)	36	(14.7)	49	
Sachet water	1	(0.4)	73	(29.8)	74	
Others	8	(3.3)	15	(6.1)	23	
Frequency of eating vegetables						

Frequent	11	(4.5)	57	(23.3)	68	0.201
Not frequent	18	(7.3)	145	(59.2)	163	
Not at all	0	(0.00)	14	(5.7)	14	
Do you walk bare foot						
Yes	25	(10.2)	155	(63.3)	180	0.098
No	4	(1.6)	61	(24.9)	65	
Do you wash your hand						
Yes	8	(3.3)	42	(17.1)	50	0.307
NO	21	(8.6)	174	(71.0)	195	
Type of toilet facility						
Pit latrine	9	(3.7)	48	(19.6)	57	0.379
Bucket latrine	6	(2.4)	39	(15.9)	45	
Open space	2	(0.8)	41	(16.7)	43	
Water System	12	(4.9)	88	(35.9)	100	

205 Key: a = Pearson chi-square test, n = Number of parasites, * = statistically significant

206 5.1 DISCUSSION

207 This study reveals a parasitic prevalence rate of 11.8% among 245 patients attending the Usmanu
 208 Danfodiyo University Sokoto Teaching Hospital, which were selected at random from May to
 209 November, 2017.

210 The low prevalence of this study is in line with the study observed in North western Nigeria of 15.67%
 211 by [19] and 12% in South India by [20]. However, the results are considerably lower than studies
 212 reported in North western Nigeria by [21], North central Nigeria by [22], western Tajikistan by [23] and
 213 North eastern Ethiopia by [24]. The lower prevalence might be due to improved environmental
 214 sanitation, better knowledge of personnel health and hygiene, and educational status of the subjects
 215 found in the study area.

216 The present study revealed that males were a little more susceptible to infection (7.8%) than the
 217 females (4.1%). This finding was found to be similar with that reported by [26]. This might be due to
 218 the common feeding pattern in which a great number of men eat outside their homes while on daily
 219 activities to earn a living. And also due to the contamination of soil by human faeces, use of raw
 220 sewage for agricultural purposes; use of waste water irrigated vegetables and contaminated imported
 221 vegetables [27].

Comment [C13]: Must be row percent!
 Calculate row percent (prevalence) in each
 category!

Comment [C14]: Check your findings! 11.9%
 among males and 11.8% among females!

Prevalence is not dependent on sex among the sampled population which disagrees with the work of [26] who observed a higher prevalence of intestinal parasite in females than in males. And the work is in contrast with that of [28], who reported that male was found to have higher prevalence rate in a study carried out in North western Ethiopia.

However, 11-15 years aged group and 16-20 years had a highest prevalence of 2.9% and 2.4% respectively. This finding was found to be similar with that reported by [21, 29]. This study is also similar to the work of [30], who reported highest prevalence in the ages 9-10 years among children [30]. Even though WHO confirmed that intestinal protozoan parasite (IPP) are dependent on age and greater severity of the infection is found in the younger children [31]. This could be attributed to the different host responses and other related factors such as nutritional status [32].

The most common intestinal parasitic infection identified in the community include amongst others *H. nana*, *Ascaris lumbricoides*, *G. lamblia*, *E. histolytica* and Hookworm specie. However Hookworm and *A. lumbricoides* recorded the highest prevalence of 15 (6.1%) followed by *G. lamblia* and *E. histolytica* 9 (3.7%) and *H. nana* recorded the least prevalence of 5 (2.0%). This finding was similar to those reported in Ethiopia [28], and in contrast with the study in Nigeria [33].

In this study, occupation, type of toilet facility and frequency eating of vegetables were not significantly associated with intestinal parasitic infections. However, according to the study conducted by [34] and [35], they were strongly associated with infections. This is more likely due to high level of education, better sanitation condition, better knowledge about the faeco-oral transmission of intestinal parasite through their unwashed hands and the contamination of vegetables with faecal materials in the farm. Season could be another important predictor of intestinal parasitic infections especially during rainy season where agricultural activities is said to be highest. This finding is in agreement with the findings of other researchers that indicated seasonal variations contributed to the higher prevalence of the disease [36, 37].

5.2 CONCLUSION

This present study revealed that there is low prevalence of intestinal parasitic infection among patients attending Usmanu Danfodiyo University Teaching Hospital, Sokoto.

The different potential risk factors assessed in the study include occupation, educational status, water source, and type of toilet facility. Water source and educational status were strongly associated with intestinal parasitic infection. However, the low prevalence of infection might be attributed to proper management of organic refuse, public health enlightenment about the risk of intestinal parasitic infections, adequate supply of clean water and proper drainage among the study participants.

Therefore, all stakeholders should give attention to raising awareness about control of intestinal parasitic infection, personal and environmental hygiene, and improving the quality of drinking water source.

REFERENCES

- [1] WHO (2000). Conquering Suffering Enriching humanity. World Health Organisation: Geneva.
- [2] Nwosu ABC. The community of soil transmitted helminth infection of humans in a hyper-endemic area of southern Nigeria. *Annals Trop Med Parasite*, 1981; **75**:197-203.
- [3] Udonso, J. K. Necator americanus: a cross-sectional study of rural community in relay to some clinical signs. *Annal. Trop. Med. Parasitol.* 1984; **78**:443-445.

263 [4] Albonico, M. Wright, V. Jape, K. Haji, H. J. Taylor, M. Savioli, L. Bickle, O. Soil transmitted
 264 nematode infections and mebendazole treatment in Mafia Island School children. *Annal Trop Med*
 265 *Parasitol*, 2002 **96**:717-726.

266 [5] Alvañiz AFA, de Castro EC, Tablizo BJB, Flores MJC, Maghirang ESV, Esmeli LAR, Pacificador
 267 AY, Sumalapao DEP. Prevalence, physiologic effects, and risk factors of soil-transmitted helminth
 268 infections among grade school children. *National Journal of Physiology, Pharmacy and*
 269 *Pharmacology*. 2017; 7(9):907-913.

270 [6] Flores MJC, Bautista YM, Carandang LGE, Go KM, Olalia MBIT, Maghirang ESV, De Peralta AS,
 271 Esmeli LAR, Sumalapao DEP. Risk factors associated with intestinal parasitic infections among
 272 school children. *Annual Research & Review in Biology*. 2018; 24(4):1-6.

273 [7] Nokes, C. S. M. Grantham-Mcgregor, A.W. Sawyer, E.S. Looper. D.A.P Bundy. Parasitic
 274 helminthic infection and cognitive function in school children. 1992

275 [8] Nokes, C. Bundy, DAP. Does helminth infection affect mental processing and educational
 276 achievement? *Parasitol Today*. 1994; **10**:(14-18).

277 [9] Sackey, M. E. Weigel, M. M. Armijos, R. X. (2003). Predictors and nutritional consequences of
 278 intestinal parasitic infections in rural Ecuadorian children. *J. Trop. Pediatr*. **49**: 17-23.

279 [10] Yakubu, N. Musa, G. Yakubu, S. E. Seasonal changes in the distribution and infection rate of
 280 *Schistosoma* intermediate hosts in River Kubanni and its tributaries. *Bio Res Com* 2003; **15**:207-214.

281 [11] Wang, K. Tang, D. Wang, M. Lu, J. Yu, H. Liu, J. *Journal of cell science* 2009; 122(Pt12):20556

282 [12] WHO. Report Series 749: Prevention and control of intestinal parasitic infections. Geneva: WHO;
 283 1987.

284 [13] Sokoto State Business Directory (SSBD). A Publication of the Commerce Department, Ministry of
 285 Commerce, Industry and Tourism, Sokoto. 2007; Pp 14-18.

286 [14] UNFPA Annual report (2013). Publisher UNFPA ISBN 978-0-89714-994-5.

287 [15] Taofeek, I. Sample Size Determination. Research Methodology and Dissertation Writing for
 288 Health and Allied Health Professionals 1st edition, Cress Global Link Limited, Abuja, 2010; **Pp** 74-75.

289 [16] Muhammad, I. M., Umoru, A. M., Isyaka, T. M., Intestinal parasitic infections among patients
 290 attending a Tertiary Health Institution in North-eastern Nigeria. *American Journal of Research*
 291 *Communication*. 2014; **2**(6): 88-96.

292 [17] Cheesbrough, M. District Laboratory Practice in tropical countries part 1, Cambridge University
 293 Press, and International U.K. **Pp** 184-205. Children in southeast Nigeria. *Journal 7, Environmental*
 294 *health*. 2005; **1**(2); 8-64.

295 [18] Ochei, J. Kolhatkar, A. Medical Laboratory Science. Theory and Practice. Tata Mcgraw-Hill
 296 Publishing Company Limited: New Delhi. 2nd Edition. 2000; **Pp** 331-349.

297 [19] Garba, D. D. Thomas, H. Z. Jatau, E. D. Inabo, H. I. Prevalence of intestinal helminths among
 298 primary school children in chikun and kaduna south local Government Area of Kaduna state, Nigeria.
 299 *J. Med. Res*. 2014; Vol 2(2), Pp 6-1.

300 [20] Baragundi, M. C. Sonth, S. B. Solahannwar, S. Patil, C. S. The prevalence of parasitic infections
 301 in patients attending tertiary care hospital, *Nat J Bas Med Sci*. 2011; **2**(1):314.

302 [21] Kabiru, M., Mohammed, R.A., and Julia O. Intestinal Parasitic Infection and Assessments of Risk
 303 Factors in North-Western, Nigeria: A community based study. *International Journal of Pharma*
 304 *Medicine and Biological Sciences*. 2015; **4:2**

305 [22] Ikeh, E., M. Obadofin, B. Brindeiro. Intestinal parasitism in rural and urban areas of North
 306 Central Nigeria, an update, *Internet J. of Microbiology*. 2006; vol. 2, no. 1.

307 [23] Matthys, B. M. Bobieva, G. Karimova, Z. Mengliboeva, V. Jean, Richard,. Prevalence and risk
 308 factors of helminths and intestinal protozoa infections among children from primary schools in
 309 Western Tajikistan, *Parasites and Vectors*. 2011; vol. 4, no. 195, **Pp**. 1-13.

310 [24] Missaye, A. D. Mulat, A. Alemu, and Agersew, A. Prevalence of intestinal parasite and
 311 associated risk factors among HIV/AIDS Patients with pre-ART and on-ART Clinic, North-east
 312 Ethiopia, *AIDS Research Therapy J*. 2013; vol. 10, no. 7, **Pp**. 1-9.

313 [25] Okon, K.O. Moses, A. E. Zailani, S. B. Tettey, E. N. "Prevelence of human intestinal parasites in
 314 a semi-arid area of Nigeria: A five year review". *Nig. J experimental applied biology*. 2003; **17**: 503-
 315 24.

316 [26] Brooker, S, Hotez P. J. Bundy, D. A. Hookworm-Related Anaemia. Among Pregnant Women: A
 317 *Systematic Review of Trop*. 2008; **Pp** 2-4.

318 [27] Atu, B.O. Galadima, M. and Alice, F. Prevalence of intestinal parasites in Etulo, Benue State
 319 Nigeria. *Nig. J. of Parasitology*. 2006; **7**: **Pp** 1-16.

320 [28] Gelaw, A. A. Belay, Nigssie, B. Prevalence of intestinal parasitic infections and risk factors
 321 among School children at the university of Gondar Community School, North-west Ethiopia, a cross
 322 sectional study, *BMC Public Health*, 2013; vol. 13, no. 1, pp. 257.

323 [29] Abou-EL, F. R. Salama, Taha, N. "Predictors of intestinal parasitic infection among pre-school
 324 children in Rural Lower Egypt, *The Egyptian Journal of Community Medicine*. 2009; vol. 27, no. 1, **Pp**.
 325 1-9.

326 [30] Oguoma, V. M. Anyasodor A. E. Mbata, T. I. Valence of Intestinal Protozoan parasites among
 327 children aged 5-12 years in Owerri Metropolis South Eastern Nigerian. *Trop. J. Health Sci*. 2008;
 328 15(2): **Pp** 43-48.

329 [31] WHO. De-worming for health and development. Report of the third global meeting of the partners
 330 for parasite control. 2005.

331 [32] Adekunle, L. Intestinal parasites and nutritional status of Nigerian children. *Afr. J. Biomed.Res*.
 332 2002; **5**: **Pp** 115-119.

333 [33] Akinbo, F. O. Omerigie R. Eromwan, R. Prevalence of intestinal parasites among patients in
 334 tertiary hospital in benin city, Nigeria. *North American journal of medical science*. 2011; 3(10):462-
 335 464.

336 [34] Erko, B. Medhin, G. Human helmenhesis in wodogenet, southern Ethiopia, with emphasis on
 337 geohelminthiasis, *Ethiop MED J*. 2003; **41**: **Pp** 333-44.

338 [35] Asrat, A. Tewodros, D. Alemayehu, W. Prevalence and risk factors of intestinal paraties among
 339 Delgi school children, North Gondar, Ethiopia. *Journal of parasitology and vector Biology*. 2011; Vol.
 340 3(5) **Pp** 75-81.

341 [36] Miller SA, Rosario CL, Rojas E, Scorza JV. Intestinal parasitic infection and associated symptoms
 342 in children attending day care centres in Trujillo, Venezuela. *Trop Med Int Health*. 2003; **8**:342-7.

343 [37] Nasiri V, Esmailnia K, Karim G, Nasir M, Akhavan O. Intestinal parasitic infections among
344 inhabitants of Karaj City, Tehran province, Iran in 2006-2008. Korean J Parasitol. 2009; 47:265-8..

UNDER PEER REVIEW