

Original Research Article

Pattern of condom usage among male and female youths in Nigeria from 1999 to 2008

ABSTRACT

Condom use is one of the most effective strategies for combating the spread of Human Immunodeficiency Virus (HIV). Despite increased sexual knowledge, adolescents in Nigeria are poor condom users. They are less likely than adults to consistently use condoms or other methods of protection that could reduce their chances of infection. The objective of the study is was to describe the trend in the condom use-usage and to determine the significant association between demographic variables and condom use-usage among youths aged 15-24 from 1999-2008.

A comparative cross-sectional population based study was carried out in which a secondary data review analyses of Nigeria Demographic and Health Survey (NDHS) 1999, 2003 and 2008 data were done. Statistical package of social science version 17.0 was used for determining frequencies, chi square and the p-values.

There was a significant increase in the usage of condom from 8.2% in 1999 to 9.3% in 2008 to 10.5% in 2008 among the female youths, though the percentages are still very low. There was an insignificant increase in condom usage among the male youths from 31.1% in 1999 to 38.7% in 2003 and then a decrease to 36.8% in 2008. There was significant association between the urban dwellers, 20-24 years age group, Christians, literate and single female youths

and condom use over the years while the rural dwellers, single and literate male youths experienced significant increase in condom use over the years at $p < 0.05$.

Condom usage by youths significantly increased over the years. However, the increase is still considered low and not enough to effectively reduce the prevalence of HIV and AIDS. Religion, marital status, age group, education and literacy are variables that influenced the use of condom by young people. Campaign on the usage of condom during sexual intercourse among young people at every level of education, urban or rural residence, age group and religion should be intensified.

Keywords: Condom use, HIV and AIDS, Youths, Nigeria, Secondary data analyses

INTRODUCTION

Acquired immunodeficiency syndrome (AIDS) is one of the world's serious public health concern, and it poses an enormous challenge to most developing countries. The HIV pandemic is on the increase with young people at a greater risk of infection, due to low risk-perception, hindering their commitment to behavioral change in spite of the fact that knowledge of some aspects of the disease is quite high. Nigerian youths between the age of 15 and 24 years have been found to be the most affected because of their engagement in risky sexual behavior, experimentation with alcohol and drugs, and failure to see that they are at the risk of infection due to their poor risk perception [1]. In many regions of the world, young people (15–24 years of age) are heavily prone to new HIV infection and they accounted for 40% of total new HIV infection [2].

According to WHO 2005, the term youth is defined as young people belonging to the age group of 15-24 years. On the other hand, Sexuality involves the entire thoughts and behavior concerning sexual activity. Nearly half of the global population is less than 25 years old [3]. The period of adolescence and youth occupies a unique stage in every person's life. It is a period of transition from childhood to adulthood [4]. Youth has also been described as a stage among human beings where a lot of physiological as well as anatomical changes take place resulting in reproductive maturity in the adolescents [5]. Many adolescents manage this transformation successfully while others experience major stress and find themselves engaging in behaviors (e.g. sexual experimentation, exploration and promiscuity etc.) that place their well-being at risk [4]

Sexual behaviour of adolescents is important not only because of the possible reproductive outcomes but also because risky sexual behaviour, such as unprotected sex and low

and inconsistent use of condom during sexual intercourse have been associated with HIV infection. Early sexual initiation poses health risks for both young men and women. Most young adults who entered into a sexual relationship for the first time did not use any form of contraception and were ignorant of the consequences of their acts leaving them vulnerable to unintended pregnancies and unplanned parenthood [6-8]. There are a lot of socio-demographic factors that had been discovered to have substantial effect on the sexual behavior of the youth and the people at large. In Kenya like other developing countries, rapid urbanization has presented development challenges leading to deteriorating living conditions and growing urban poverty [9]. Young people form a large proportion of those moving from rural to urban areas in search of livelihood opportunities and most find urban slums as the first entry points into the cities. These present enormous challenges faced by most of these urban slums are underserved by health facilities, and challenged by other socio-economic amenities [3]. Because of limited livelihood opportunities and the frustrations of unemployment, many young adults in these settings turn to risky sexual behaviours. For example, they seek comfort in prostitution and drug abuse which expose them to HIV. Other behavioral factors that increase young urban women's risk for HIV infection include having older sexual partners, inconsistent condom use, forced sex, and transactional sex [10].

Condom use is one of the most effective strategies for combating the spread of HIV. However, opinions on educating young people about condom use are sometimes diverse, with some people believing it promotes early sexual initiation [11], while others believe it encourages sexual experimentation. The total number of condoms provided by international donors has been relatively low. Between 2000 and 2005, the average number of condoms distributed in Nigeria by donors was 5.9 per man, per year [12]. Restrictions on condom promotion have hampered

HIV prevention efforts. In 2001, a radio advertisement was suspended by the Advertising Practitioners Council of Nigeria (APCON) for promoting messages suggesting that it is acceptable to engage in premarital sex as long as a condom is used. Premarital sex engagement predisposes an individual to high risk of HIV and other sexually transmitted diseases. Hence, the promotion of condom usage is highly encouraged. In 2006, APCON also started to enforce stricter regulations on condom advertisements that might encourage increased engagement in unprotected sexual relationship [12].

In some African countries, qualitative interviews were carried out about the misconception people have about condom use. Different beliefs appear to influence some men's behavior. For example, some men in the primary survey said that they did not use condoms because condoms were infected with 'AIDS worms.' Qualitative interviews in Namibia revealed the key spiritual and genocidal beliefs that HIV is a punishment from God and that condoms cannot be trusted e.g. manufacturers put holes in condoms; condoms spread disease and government condoms are of low quality [14].

Despite increased sexual knowledge, adolescents in Nigeria are poor contraceptive users [15]. They are less likely than adults to consistently use condoms or other modern methods of contraceptives that could reduce their chances of infection [4]. Neither specific teaching about contraception nor improving the contraceptive service consistently increases effective contraceptive use by the youth [5]. Some other misconceptions harbored by many young people includes, the belief that mosquitoes can transmit HIV infection, false claims that use of contraceptives can cause infertility, and contracting HIV through hugging, holding of hands and sharing of a spoon of an infected person [11]

The Nigerian HIV and AIDS Emergency Action Plan (HEAP) brought to light the lack of support for non-schooling (Never been to school and out of school) youth [16]. There seems to have been an increase in the exposure to risks of sexually transmitted infections including HIV and AIDS, amongst these young people and this is often based on their risky behaviors and sexual interactions with high-risk groups [17-19]. Non-schooling youths are young people that have never been to school or have been out of school and they mostly engage in multiple sexual partners and unprotected sexual activities [20,21]. Very few percentage of unmarried young people use condom at sexual debut, while those that use it do so inconsistently [22]. A study in Republic of Benin have recorded figures of condom use with a recent non-regular partner as low as 20% amongst men with no formal education, 18–19% in those with primary education and 39% in those with secondary education and above [23]. This clearly shows that while young people already own their sexual conduct [24] the capacity to take definite decisions to protect themselves from harmful consequences of unprotected sex is still generally lacking.

Despite series of strategies to improve condom use among the young people, the proportion of youth that make use of condom during sexual intercourse is still low. Furthermore, the pattern of condom usage in Nigeria cannot be ascertained. Therefore, this study is carried out to describe the trend of condom usage among male and female Nigerian youths between 15-24 years age group over the years from 1999 to 2008. The study is carried out to identify some socio-demographic factors that could influence the usage of condom among the youths and to determine the association between socio-demographic variables and condom use among youth aged 15-24 years over the period.

METHODOLOGY

Study Area: The primary survey, a national study (NDHS 1999, 2003 and 2008) for Nigeria was used. Nigeria is made up of 36 states and a Federal Capital Territory (FCT), grouped into six geopolitical zones: North Central, North East, North West, South East, South South, and South West. There are also 774 constitutionally recognized local government areas (LGAs) in the country. There are about 374 identifiable ethnic groups in Nigeria with the Igbo, Hausa, and Yoruba as major groups.

Study Population: The population covered by the 2003 and 2008 NDHS is defined as the universe of all men and women aged 15-49 years in Nigeria.

Unit of Enquiry: For this study, the study unit of enquiry was the young men and women aged 15-24 years from the study population of the 1999, 2003 and 2008 NDHS. According to the United Nations, 'youth', were defined as those persons between the ages of 15 and 24 years, thereby ensuring statistical consistency [13].

Study Design: This is a comparative cross-sectional population-based study. It is a secondary data analysis of data from 1999, 2003 and 2008 NDHS. For this study, condom use by young people aged 15-24 years in Nigeria and the outcomes on the burden of the scourge were compared using the 1999, 2003 and 2008 NDHS.

Sampling Technique: The sampling technique used for 1999, 2003 and 2008 NDHS was a stratified two-stage cluster design [11]. The primary sampling unit (PSU) also known as cluster for the 2003 and 2008 NDHS is defined on the basis of Enumeration Areas, EAs from the 1991 and 2006 EAs census frames.

Sampling Frame: The sample frame for this survey was the list of enumeration areas (EAs) developed for the 1991 Population Census. Administratively, at the time the survey was planned, Nigeria was divided into 36 states and the Federal Capital Territory (FCT) of Abuja. Each state was subdivided into local government area (LGA) units and each LGA was divided into localities. In addition to these administrative units, for implementation of the 1991 Population Census, each locality was subdivided into enumeration areas (EAs). The list of approximately 212,080 EAs, with household and population information (from the 1991 census) for each EA, was evaluated as a potential sampling frame for the 1999, 2003 and 2008 NDHS. The EAs are grouped by states, by LGAs within a state, and by localities within an LGA, stratified separately by urban and rural areas. For NDHS 2008, at the time of survey implementation, the list of EAs of 2006 population census used did not have census information for households and the population because the census frame is under segmentation revision. Therefore, no household or population information was available at the EA level. Therefore, each EA was approximately classified as urban or rural.

Stratification: In the current preliminary census frame, the EAs are grouped by states, by LGAs within a state, and by localities within an LGA. The EAs are stratified separately by urban and rural areas. Any locality with less than 20,000 populations in each LGA constitutes the rural area in the LGA.

Sample Selection: The NDHS sample was selected using a stratified, two-stage cluster design consisting of a total of 365 clusters, 165 in urban and 200 in rural areas for 2003 survey and 888 clusters, 286 in the urban and 602 in the rural areas selected in 2008 survey.

Sample Size: A total of 7,864 households and 36800 households were recruited in 2003 and 2008 NDHS respectively. The 2008 NDHS sample was selected using a stratified two-stage cluster design [11] consisting of 888 clusters, 286 in the urban and 602 in the rural areas¹. A representative sample of 36,800 and 7,864 households was selected for the 2008 and 2003 NDHS survey respectively, with a minimum target of 950 completed interviews per state. In each state, the number of households was distributed proportionately among its urban and rural areas. About 734, 879 and 4910 men of age 15-24 years were interviewed in 1999, 2003 and 2008 NDHS while 2482, 3210 and 12626 women of age 15-24 years were interviewed in 1999, 2003 and 2008 NDHS.

Data Management: For the purpose of this study, the secondary data was analyzed using SPSS statistical package 17.0 and Epi-info package. Different changes such as dataset merging, variables recoding and removal of some missing data were carried out. Dependent variables used in the analysis included last intercourse used condom while independent variables include: region, place of residence, educational attainment, marital status, literacy and religion. Different recording and computation was carried out. Descriptive statistics such as frequencies and percentages were used to summarize socio-demographic characteristics and dependent variables while cross-tabulations and proportions were used to summarize dependent variables with respect to the socio-demographic properties of the respondents. Chi-square and significance obtained using StatCalc of Epi-Info statistical software were used for to investigate trends and

associations between dependent and independent variables. All analyses were done at the 5% level of significance and 95% confidence interval.

Ethical approval was from the ethical review committee for the research work. Additional approval and consent was given by the National Demographic and Health Survey (NDHS) for the release of the data regarding condom use among young people.

RESULTS

Table 1 below shows the distribution of the socio-demographic characteristics of female and male youths aged 15-24 years. About 53.7%, 54.4% and 51.9% of female youths involved in the study are within the age group 15-19 years for 1999, 2003 and 2008. Young women that live in the urban residence ranges between 31% to 40% and higher percentage (60% to 68%) of these young women reside in the rural residences. Higher proportion of female youth had secondary education while about 5% of the respondents were able to attain higher level of education. More proportion of Christian female youths were interviewed in the course of the three surveys compared to the Muslim respondents. Over 50% of the respondents interviewed during the surveys were singles while only about 2% of the respondents were divorced.

A higher percentage (51.7% to 62%) of male youths enrolled in 1999, 2003 and 2008 in the study are within the age group 15-19 years. About 54.4% to 66.4% of the male respondents live in the rural residence. In the aspect of education, higher proportion (>60%) of male respondents had secondary education as their highest level of education attainment at the time of

survey. About 61.2%, 55.1% and 55.2% of the male youths in 1999, 2003 and 2008 studies respectively are Christians. Over 90% of the male respondents are single.

Table 1: Socio-demographic characteristics of female and male youths

	1999	2003	2008	1999	2003	2008
	FEMALE			MALE		
Variables	n (%)	n (%)	n (%)	n (%)	n (%)	n (%)
Age group						
15-19 years	1774 (53.7)	1749 (54.4)	6591 (51.9)	513 (62.0)	453 (50.7)	2571 (51.7)
20-24 years	1528 (46.3)	1464 (45.6)	6103 (48.1)	315 (38.0)	441 (49.3)	2399 (48.3)
Total	3302 (100.0)	3213 (100.0)	12694 (100.0)	828 (100.0)	894 (100.0)	4970 (100.0)
Residence						
Urban	1095 (33.2)	1282 (39.9)	4028 (31.7)	285 (34.4)	408 (10.6)	1671 (33.6)
Rural	2207 (66.8)	1931 (60.1)	8666 (68.3)	543 (65.6)	486 (54.4)	3299 (66.4)
Total	3302 (100.0)	3213 (100.0)	12694 (100.0)	828 (100.0)	894 (100.0)	4970 (100.0)
Education						
No formal Education	956 (29.0)	901 (28.0)	3831 (30.2)	106 (12.8)	95 (10.6)	732 (14.7)
Primary	691 (20.9)	646 (20.1)	1938 (15.3)	184 (22.2)	200 (22.4)	723 (14.5)
Secondary	1523 (46.1)	1520 (47.3)	6260 (49.3)	504 (60.9)	544 (60.9)	3194 (64.3)
Higher	132 (4.0)	146 (4.5)	665 (5.2)	34 (4.1)	55 (6.2)	321 (6.5)
Total	3302 (100.0)	3213 (100.0)	12694 (100.0)	828 (100.0)	894 (100.0)	4970 (100.0)
Region						
North-Central	594 (18.0)	521 (16.2)	2484 (19.6)	164 (19.8)	166 (18.6)	1077 (21.7)
North East	413 (12.5)	547 (17.0)	2361 (18.6)	103 (12.4)	124 (13.9)	821 (16.5)
North West	717 (21.7)	710 (22.1)	2580 (20.3)	112 (13.5)	205 (22.9)	945 (19.0)
South East	401 (12.1)	498 (15.5)	1456 (11.5)	94 (11.4)	114 (12.8)	502 (10.1)
South South	614 (18.6)	433 (13.5)	1832 (14.4)	183 (22.1)	129 (14.4)	856 (17.2)
South West	563 (17.1)	504 (15.7)	1981 (15.6)	172 (20.8)	156 (17.4)	769 (15.5)
Total	3302 (100.0)	3213 (100.0)	12694 (100.0)	828 (100.0)	894 (100.0)	4970 (100.0)
Religion						
Christianity	1848 (56.0)	1754 (54.6)	6853 (53.9)	506 (61.2)	493 (55.1)	2739 (55.1)
Islam	1409 (42.6)	1432 (44.6)	5618 (44.3)	310 (37.4)	394 (44.1)	2165 (43.6)
Others	45 (1.4)	27 (0.8)	223 (1.8)	12 (1.4)	7 (0.8)	66 (1.3)
Total	3302 (100.0)	3213 (100.0)	12694 (100.0)	828 (100.0)	894 (100.0)	4970 (100.0)
Marital Status						
Single	1831 (55.5)	1836 (57.1)	6747 (53.2)	759 (91.7)	827 (92.5)	4537 (91.3)
Married	1418 (42.9)	1310 (40.8)	5773 (45.5)	57 (6.9)	59 (6.6)	414 (8.3)
Divorced	53 (1.6)	67 (2.1)	173 (1.4)	12 (1.4)	8 (0.9)	19 (0.4)

Total	3302 (100.0)	3213 (100.0)	12693 (100.0)	828 (100.0)	894 (100.0)	4970 (100.0)
-------	--------------	--------------	---------------	-------------	-------------	--------------

Table 2 below shows that among the female youths, there was a significant increase in the usage of condom from 8.2% in 1999 to 9.3% in 2003 to 10.5% in 2008, though the percentages are still very low. There was an insignificant increase in condom usage among the male youths from 31.1% in 1999 to 38.7% in 2003 and then a decrease to 36.8% in 2008.

Table 2: Condom use among female and male youths

		1999	2003	2008
Variables		n (%)	n (%)	n (%)
Last intercourse	Yes	146 (8.2)	169 (9.3)	796 (10.5)
Used condom	No	1626 (91.8)	1640 (90.7)	6780 (89.5)
(FEMALE)	Total	1772 (100.0)	1809 (100.0)	7576 (100.0)
Last intercourse	Yes	104(31.1)	126(38.7)	623 (36.8)
Used condom	No	230 (68.9)	200(61.3)	1072 (63.2)
(MALE)	Total	334(100.0)	326(100.0)	1695 (100.0)

Figure 1: Chart of condom use among female and male youths




Table 3 describes the relationship between the socio-demographic characteristics and condom use among female youths. There was a statistically significant increase at $P < 0.05$ ($X^2 = 19.85$, $P = 0.00001$) of condom use during last intercourse among the youths that reside in the urban areas. There was a slight reduction in the proportion of female youth living in the rural environments, that used condom during last intercourse from 1999 to 2003 and increase from 2003 to 2008 though not statistically significant at $P < 0.05$ ($X^2 = 1.36$, $P = 0.244$).

There was significant increase ($X^2 = 16.75$, $P = 0.00004$) in the use of condom among respondent at the secondary level from 1999 to 2008 but there was statistical decline ($X^2 = 3.94$, $P = 0.04715$) from 1999 to 2008 for respondent at the primary level of education. There was no significant increase in the usage of condom among those that had no formal education and higher education.

All geo-political zones in Nigeria showed an insignificant increase in condom use except the youth in the South South region that had significant increase at $P < 0.05$ ($X^2 = 23.08$, $P = 0.00001$) in the proportion of youth that used condom in their last intercourse from 1999 to

2008. There was a statistical and steady increase at $P < 0.05$ ($X^2 = 32.60$, $P = 0.00001$) of condom usage of among the literate female youths from 1999 to 2008.

Table 3: ASSOCIATION BETWEEN SOCIODEMOGRAPHIC FACTORS AND CONDOM USE AMONG FEMALE YOUTHS

	1999	2003	2008		
	Yes (%)	Yes (%)	Yes (%)	Chi square	P value
Age group					
15-19	53(8.3)	69(9.5)	246(8.7)	0.006	.94
20-24	93(8.2)	100(9.2)	550(11.6)	13.25	.00027*
Total	146(8.2)	169(9.3)	796(10.5)	9.15	.0025*
Residence					
Urban	67(12.5)	105(16.1)	410(20.3)	19.85	0.00001*
Rural	79(6.4)	64(5.5)	386(6.9)	1.36	0.24
Total	146(8.2)	169(9.3)	796(10.5)	9.15	0.0025*
Education					
No Formal Education	6(0.9)	2(0.3)	23 (0.7)	0.007	0.96
Primary	24(6.5)	17 (5.0)	47 (4.0)	3.94	0.047*
Secondary	89(13.3)	118(18.4)	560(20.4)	16.75	0.00004*
Higher	27(30.7)	32(43.8)	166(40.6)	1.99	0.16
Total	146(8.2)	169(9.3)	796(10.5)	9.15	0.0025*
Region					
North Central	35(11.3)	29(10.8)	128(10.3)	0.27	0.60
North East	7(2.9)	3(0.8)	45(2.7)	0.58	0.45
North West	4(0.8)	4(0.7)	11(0.5)	0.69	0.40
South East	29(22.1)	30(17.1)	141(26.3)	2.87	0.090
South South	43(12.6)	42(16.7)	207(24.5)	23.08	0.000001*
South West	28(10.4)	61(32.4)	264(21.4)	7.70	0.0055*
Total	146(8.2)	169(9.3)	796(10.5)	9.15	0.0025*
Marital Status					
Single	122(22.1)	143(28.1)	655(32.4)	22.28	0.000001*
Married	21(1.8)	22(1.8)	129(2.4)	2.33	0.13

Divorced	3(6.1)	4(8.0)	12(10.9)	1.02	0.31
Total	146(8.2)	169(9.3)	796(10.5)	9.15	0.0025*
Religion					
Christians	122(14.0)	139(18.0)	675(20.2)	17.47	0.00003*
Islam	23(2.6)	29 (2.8)	110(2.7)	0.000	0.99
Others	1 (5.3)	1 (10.0)	8(7.4)	0.054	0.82
Total	146(8.3)	169(9.3)	793(10.5)	9.15	0.0025*
Literacy					
Illiterate	10(1.3)	12 (1.3)	44(1.1)	52.89	0.000001*
Partly literate	15(7.9)	5 (5.7)	14(4.5)	2.39	0.12
Literate	120(14.8)	152(19.5)	733(22.4)	32.604	0.000001*
Total	145(8.2)	169(9.5)	791(10.5)	9.17	0.0025*

*= Significance at $P < 0.05$

Table 4 describes the relationship between the socio-demographic characteristics and condom use among female youths. There was a steady increase in the usage of condom among male youths both in the urban and rural residences from 1999 to 2008 though there was only significant increase at $P < 0.05$ ($X^2 = 41.65$, $P = 0.00001$) among the respondents in the rural areas from 1999 to 2008. There was no significant increase in the proportion of young male youths that used condom during their last intercourse for the three years.

There was a significant increase ($X^2 = 0.54$, $P = 0.0331$) in the proportion of male youths in the South East regions from 1999 to 2008. There is no significant increase in proportion among those in the other regions. The proportion of young male respondents that used condom during the last intercourse significantly increased ($X^2 = 19.15$, $P = 0.00001$) from 1999 to 2003 but slightly declined in 2008. The proportion of young men that are literate is highest in the three years.

Table 4: ASSOCIATION BETWEEN DEMOGRAPHIC CHARACTERISTICS AND CONDOM USE AMONG MALE YOUTH

	1999	2003	2008		
Variables	Yes (%)	Yes (%)	Yes (%)	Chi square	P value
Age group					
15-19	34(25.4)	28(31.1)	141(31.8)	1.83	0.18
20-24	70(35.0)	98(41.5)	482(38.5)	0.25	0.62
Total	104(31.1)	126(38.7)	623(36.8)	2.36	0.12
Residence					
Urban	50(45.9)	74(51.4)	291(53.8)	2.23	0.14
Rural	54(24.0)	52(28.6)	332(28.8)	41.65	0.000001*
Total	104(31.1)	126(38.7)	623(36.8)	2.36	0.12
Education					
No Formal Education	1(2.4)	0(0.0)	6(2.8)	0.145	0.70
Primary	14(17.1)	10(14.3)	32(16.8)	0.001	0.97
Secondary	74(39.4)	97(47.3)	466(41.8)	0.001	0.97
Higher	15(68.2)	19(79.2)	119(68.8)	0.116	0.73
Total	104(31.1)	126(38.7)	623(36.8)	2.36	0.12
Region					
North Central	17(24.3)	29(42.0)	128(32.5)	0.47	0.50
North East	7(17.5)	8(17.0)	32(12.7)	0.99	0.32
North West	4(13.8)	4(8.5)	12(8.7)	0.50	0.48
South East	16(40.0)	20(51.3)	92(58.6)	0.54	0.033*
South South	22(27.5)	23(41.8)	162(39.2)	2.95	0.086
South West	38(50.7)	42(60.9)	197(57.6)	0.70	0.40
Total	104(31.1)	126(38.7)	623(36.8)	2.36	0.12
Marital Status					
Single	97(36.5)	122(46.7)	592(46.4)	6.85	0.009*
Married	2(3.6)	3(5.2)	27(6.7)	0.93	0.33

Divorced	5(41.7)	1(14.3)	4(28.6)	0.46	0.50
Total	104(31.1)	126(38.7)	623(36.8)	2.36	0.12
Religion					
Christians	80(37.7)	89(45.6)	160(24.8)	21.03	0.000001*
Islam	23(19.7)	37(28.9)	12(8.7)	5.75	0.016*
Others	1(20.0)	0(0.0)	92(58.6)	5.22	0.022*
Total	104(31.1)	126(38.7)	264(28.1)	3.29	0.070
Literacy					
Illiterate	3(3.9)	4(8.7)	15(5.5)	3.01	0.083
Partly literate	6(13.3)	3(14.3)	13(14.9)	0.062	0.80
Literate	95(40.6)	119(46.8)	590(44.5)	19.15	0.00001*
Total	104(31.1)	126(38.7)	623(36.8)	2.36	0.12

*= Significance at $P < 0.05$

DISCUSSION

Over 10% of the world population is affected by the scourge of HIV and AIDS [2]. In 2006, UNAIDS stated that young people of age 15 and above account for about 40% of new HIV infections. The future course of the national response to the HIV and AIDS epidemic in Nigeria had shown to greatly depend on a number of factors including levels of HIV and AIDS-related knowledge among the general population; social stigmatization; risk behavior modification; risk perception; access to quality services for sexually transmitted infections (STI); correct and consistent use of condom; provision and uptake of HIV counseling and testing; and access to care and anti-retroviral therapy (ART)[11].

Adolescence is a transition period between childhood and adulthood and this is a time when many youth experience critical life-defining changes, such as their first sexual experience, marriage, pregnancy, and parenthood [7]. Initiation of sexual intercourse has been described as a milestone in the physical and psychological development of men and women in all societies [28].

This study reported a pattern with about 8.2% to 10.5% of female youths and 31.1% to 38.7% of male youths used condom during their last sexual intercourse. Increase in the usage of condom among the female youths explains that young women are now getting more aware of the importance of condom usage to prevent unwanted pregnancy and sexually transmitted diseases (STIs). This result is supported by the result of a study of adolescent sexuality in Peru which showed that 38% of male youths used condom at the time of first intercourse [6]. This low proportion might be due to the belief that educating young people on condom use may lead to early initiation into sex among the youths. Also in this study, it was observed that there was an overall improvement in condom use among male youths through the years from 1999 to 2003 and a slight decrease in 2008. This could be as a result of poor compliance in the usage of condom during sex by the male respondents. The increase appears to be accounted for by the early adults aged 20-24 years living in the urban residence and those that were Christians, never married and at their secondary level of education. This means that the early adults have adequate knowledge about condom use while young people between 15-19 years of age are more likely to be infected with HIV, thereby increasing the burden of HIV and AIDS in Nigeria.

Conversely among the female youths, the pattern appears different as female youth in the rural area shows significant increase in condom use during their last intercourse while those in the urban shows an increase over the years but was not significant. This shows that most rural dwellers do not have adequate information about the use of condom. They are more likely to have higher burden of HIV compared to the female youths living in the urban areas. Therefore, residence plays an important role in describing predictors of the trend of condom usage. The result above is also supported by a study that indicated that 42% of adolescent girls in a rural community of Rivers state had had induces abortion or STIs, including gonorrhea because of

lower contraception usage [29]. The proportion of young women that use condom was considerably low compared to their male counterpart because the sexual act might be unplanned or preference of sexual enjoyment over protection against HIV and other STIs. Even though there was an increase in the proportion over the years but the increase was marginal. In a related study carried out among out of school youths in a local government area in Nigeria [25], higher proportion of male youths (44.7%) uses condom during their last sexual intercourse compared to female youths (29.7%). Comparatively, in a study carried out in 2012 [26], the proportion of urban youths that used condom during sexual intercourse was higher than those that reside in the rural area. A closely related finding was observed among South African citizens [27], 8% and 11% of men and women respectively living in rural residence and less educated used condom during their last intercourse compared to 29% and 34% of men and women living in urban and more educated. Some of the observed facts show the success of some of the programs and intervention targeted towards enlightening people, especially the young individuals about condom use. Effort is expected to be geared towards increasing condom use among youths (aged 15-24 years) especially the female youths and those living in the rural areas.

Also from this study, about 2.4-2.8% of young men that have no formal education, 14.3-17.1% of those with primary education and about 39.4 % of those with secondary education used condom during their last sexual intercourse. This is in accordance with a study in Benin Republic in 2000 [23], the study showed that 20% of men with formal education, 18-19% of those at primary school, about 39% of those in secondary school and above of non regular young used condom during sexual intercourse. This observed fact was further buttressed by the study carried in Nigeria [25], higher proportion of respondent that had secondary level (53.4%) of education used condom compared to those with no formal education (15.4%), and thereby showing that

education is an important factor that could influence condom use. The percentage of respondents in this study at the secondary and higher level of education increased from 1999-2003 but declined in 2008. This could be because of the programs geared towards educating young people at this level of education seem to be failing in recent times.

This study also shows higher proportion of female and male youths between age 15-24 years that have never married used condom during their last sexual intercourse compared to those that are married or divorced. This means that the singles have high sense of responsibilities of preventing themselves from contracting HIV or other STIs during sex because they are not sure if their partners have multiple sexual partners which might increase the odds of having HIV. However, the married youths have the belief that they don't have to use condom with their partners because they believe that they are the only sexual partners they have. Some of the married respondents also believe it's absurd to be using condom with their partners. The study may not be entirely different from what is known in a study carried out in South Africa [27] in which 15% and 18% of married women consistently and occasionally used condom, and which is considered low compared to the proportion of singles that use condom with their partners during sex. This contributes that marital status is an essential factor that can determine or influence condom use among the youths. .

CONCLUSION AND RECOMMENDATION

Young people between the ages 15 to 24 years have accounted for about 40% of new HIV infection and the positive response to this scourge can only be successful if they adopt behaviours that will protect them from the infection and increase their comprehensive knowledge about the mode of transmission, use of condom, means of contracting the infection. Low

proportion of young men and women used condom in their last sexually intercourse. Condom usage by women significantly increase from 8.2% in 1999 to 10.5% in 2008, this little change over 10 year period is not enough to effectively reduce the prevalence of HIV/AIDS. Likewise, religion, marital status, age group and education are variables that influence the use of condom by young people. Higher percentage of Christians use condom compared to the Muslims, which calls for further investigation on why Muslim respondents don't use condom. Also higher percentage of respondent that are never married and that are within age group 20 to 24 uses condom during sexual intercourse.

Effort should be made by the government and various HIV control organizations towards increasing condom usage among young people which have always been very low. Campaign on the usage of condom during sexual intercourse among young people at every level of education especially at the higher education. Rural dwellers should also be sensitized more on the benefits of condom use during sex. Emphasis should be on the benefit of its usage and implication of its non usage during sexual intercourse. Use of condom among the married youths should not only be positioned in contraception during family planning but also in preventing the contraction of HIV and AIDS which is important in the reduction of the prevalence of the disease.

Conclusively, more work should be done on examining the trend of condom usage beyond 2008 and to monitor the effectiveness of programs and campaign that had been targeted towards increasing the knowledge over time and possibly change their strategies if need be. Also, it will add to knowledge and help our policy making if a work is done on comparing the trend of comprehensive HIV and AIDS knowledge and condom use among young men and women. Therefore, policy makers should integrate education on HIV and AIDS prevention in schools and

religious organizations in order to increase awareness of possible health burden of HIV and AIDS on the socio-economic characteristics of individuals and the country.

Limitation

The only limited observed in the study is the fact that the study is secondary data analysis. We didn't meet the respondents used for the original survey in order to be able to obtain other vital information about their condom use over the years. Furthermore, the study was carried out in the year 2012 and was only able to review three surveys (1999, 2003 and 2008). We couldn't include 2013 survey in the study of the pattern of condom use among the young people over the years. Therefore, researchers should extend the investigation of the trend of condom use beyond 2008.

REFERENCES

1. Makinwa-Adebusoye P. Sexual behavior, reproductive knowledge and contraceptive use among young urban Nigerians. *International Family Planning Perspectives*. 1992;18 (2):66–70.
2. UNAIDS. AIDS epidemic update. Geneva, Switzerland: UNAIDS; November; 2006.
3. UNAIDS, author. End 2004 global epidemic update. Geneva: 2004.
4. Adegoke AA. Adolescents in Africa: Revealing the problems of teenagers in contemporary African society, Ibadan. *Hadassah Publishing*; 2003: 37-41.

5. Kirby D, Brener ND, Brown NL, Peterfreund N, Hillard P and Harrist R. The impact of condom availability (correction of distribution) in seattle schools on sexual behavior and condom use. *Am J Public Health*, 1999; 89(2); 182-187.
6. Magnani RJ, Seiber EE, Gutierrez EZ, Vereau D. Correlates of sexual activity and condom use among secondary-school students in urban Peru. *Stud Fam Plann* 2001; 32:53-66.
7. Oyediran KA, Ishola GP, Adewuyi AA. Knowledge of possible pregnancy at first coitus: a study of in-school adolescents in Ibadan, Nigera. *Biosoc Sci* 2002; 34: 233-48.
8. Mberu BU. Protection before the harm: the case of condom use at the onset of premarital sexual relationship among youths in Nigeria. *Afr Popul Stud* 2008; 23:57-83.
9. Walter Odhiambo, Damiano Kulundu Manda. Urban poverty and labour force participation in Kenya. Nairobi, Kenya: Kenya Institute for Public Policy Research and Analysis (KIPPRA); November; 2003; 12: 89-103
10. Pettifora Audrey E, van der Stratena Ariane, Dunbara Megan S, Shiboskia SC, Padiana Nancy S. Early age of first sex: a risk factor for HIV infection among women in Zimbabwe. Lippincott Williams & Wilkins; 2004.
11. Nigeria Demographic and Health Survey (NDHS) 2008. 2008 Edition. Accessed 27th August, 2018<http://www.census.gov.ph/hhld/ndhs_2008.html>
12. United Nations Population Fund (UNFPA) Condom Programming for HIV Prevention. 2005 UNFPA, World Health Organization, PATH. ISBN: 0-89714-733-2
13. United Nations Population Fund (UNFPA) UNFPA Annual Report 2007. UNFPA, ISBN: 978-89714-883-2

14. Mufune P. Myths about condoms and HIV/AIDS in rural northern Namibia. *International Social Science Journal* 2005; 57(186) 1468-2451
15. Esere MO. HIV/AIDS awareness of in-school adolescents in Nigeria: Implications for adolescence sexuality. *Journal of Psychology in Africa*. 2006; 16(2):255-258
16. Federal Ministry of Health. HIV/AIDS Emergency Action Plan (HEAP) Nigeria: FMOH; 2001.
17. Dare OO, Oladepo O, Cleland JG, Badru OB. Reproductive health needs of young persons in markets and motor parks in South West Nigeria. *Afr J Sci*. 2001; 30:199–205.
18. Onwuamanam DO. Sexual networking among youth in Southwestern Nigeria. *Health transition review supplement*. 1995; 5:57–66.
19. Feyisetan B, Pebley AR. Premarital Sexuality in Urban Nigeria. *Studies in Family Planning*. 1989;20(6.1):343–346.
20. Jinadu MK, Odesanmi WO. Adolescent sexual behavior and condom Use in Ile-Ife, Nigeria. *Clinical Nursing Research*. 1993;(2):111–118.
21. Araoye MO, Adegoke A. AIDS-related knowledge, attitude and behavior among selected adolescents in Nigeria. *Journal of Adolescence*. 1996; 19 (2):179–181.
22. Egger M, Ferrie J, Gorter A, Gonzales S, Gutierrez R, Pauw J, Smith GD. HIV/AIDS-related knowledge, attitudes and practices among Managuan secondary school students. *Bulletin of the Pan American Health Organization*. 1993; 27 (4):360–369
23. UNICEF. Prevention of infection among adolescents and young people. Child info, monitoring the situation of children and women. UNICEF; 2009.

24. Dowsett G, Aggleton P, Mane P. Re-thinking young people's sexuality and HIV risk: New Research findings from UNAIDS. *International AIDS Society Newsletter* 1997; (7):6–8.
25. Adebisi O.A and Azuzu M.C. Condom use amongst out of school youths in a local government area in Nigeria. *African Health Science*, 2009; 9(2): 92–97.
26. Doyle AM, Mavedzenge SN, Plummer ML, Ross DA. The sexual behavior of adolescents in sub-Saharan Africa: patterns and trends from national surveys. *Tropical medicine & International health*, 2012; 17 (7) 796-807.
27. Maharaj, Pranitha. Patterns of condom use: Perspectives of men in KwaZulu-Natal, South Africa. *Development Southern Africa*, 2005 (22) 187-197.
28. Singh S, Wulf D, Samara R, Cuca YP. Gender differences in the timing of first intercourse: data from 14 countries. *Int Fam Plann Perspect* 2000;26:21-8, 43.
29. Brabin L, Kemp J, Obunge OK, Ikimalo J, Dolimore N, Odu NN *et al.* Reproductive tract infections and abortion among adolescent girls in rural Nigeria. *Lancet* 1995;345:270-1.