

Evaluation of the nutrients, antinutrients and metals content of five leafy vegetables in Dengi metropolis

ABSTRACT

Aim: To quantify the levels of amino acids, some metals, and phytochemicals/antinutritional factors in leafy vegetables: *Cucurbita pepo*, *Vitex doniana*, *Hibiscus cannabinus*, *Leptadenia hastata*, *Balanites aegyptiaca*.

Study design: The research work is descriptive.

Place and duration of study: Department of Biochemistry, University of Jos between March 2016 and September 2016.

Methodology: Five samples each of the vegetables were collected at random from different locations in the area of study. Atomic Absorption Spectrophotometry was applied to analyse for metals. Levels of amino acids in samples were determined using the PTH amino acids analyser. Antinutrients were quantified using standard methods. One way ANOVA was used to analyse the data obtained at 95% level of significance.

Results: Mean levels of Pb and Cd were 0.030 ± 0.01 ppm and 0.004 ± 0.001 ppm respectively ($P > 0.05$) relative permissible limits. Mean Mg, Mn and Fe content were 1.084 ± 0.02 ppm, 0.069 ± 0.01 ppm and 1.534 ± 0.10 ppm respectively ($P = 0.05$). Amino acid profile indicated mean values (g/100g proteins) of Glutamate 8.34, Aspartate 8.14, Leucine 8.34, Lysine 4.03, Isoleucine 3.30, Phenylalanine 4.17, Tryptophan 2.25, Valine 4.6, Methionine 1.12, Proline 2.84, Arginine 4.99, Tyrosine 2.75, Histidine 2.23, Cysteine 1.09, Alanine 3.71, Glycine 4.08, Threonine 2.88, Serine 2.99. Mean range of antinutrients were: tannins, 0.51 ± 0.13 % to 0.60 ± 0.12 %, oxalates, 0.14 ± 0.14 % to 0.60 ± 0.20 %, phytates, 1.70 ± 0.01 mg/100g to 4.10 ± 0.01 mg/100g, saponins, 11.85 ± 1.85 % to 15.13 ± 1.50 %, cyanogenic glycosides, 4.82 ± 1.30 % to 7.59 ± 1.20 %, total alkaloids and total flavonoids were 16.22 ± 1.61 % to 19.37 ± 1.23 % and 9.87 ± 1.32 % to 14.71 ± 2.30 % respectively.

Conclusion: Although samples analysed contained significant amounts of antinutrients, they are very good sources of amino acids; especially Lysine, Methionine, Leucine, tryptophan which are essential; and mineral elements. Levels of lead and cadmium in the samples were lower than safe limits. These vegetables are good sources of nutrients. Their consumption will replenish nutrients to the cell thereby improving the well being of consumers.

Key words: amino acids, antinutrients, nutrients, metals, vegetables

41

42

43 INTRODUCTION

44 Leafy vegetables have been shown to be valuable sources of nutrients [1] with some having
45 medicinal properties [2]. ~~These~~ ~~these~~ vegetables serve as valuable sources of nutrients especially in
46 rural areas like Dengi where they exist in the open country. The feeding pattern of people in Dengi
47 metropolis suggests a heavy reliance on many leafy vegetables commonly found in the town. The five
48 leafy vegetables for this study are the most common ones found in ~~the~~ town.

49 Leafy vegetables might contain significant levels of trace elements, heavy metals, amino
50 acids as well as antinutrients. Leafy vegetables can contribute substantially to food security in the
51 rural areas where people's diet is based on mostly carbohydrates and legumes which are high in
52 calories but deficient in essential micronutrients and proteins [3].

53 Antinutritional factors reduce the nutritive values of many plants due to their natural inherence in the
54 plants. They are capable of eliciting deleterious effect in man and animals [4]. Oxalate tends to render
55 calcium unavailable by binding to the calcium ion to form complexes [1, 5, and 6]. Phytic acid acts as a
56 strong chelator forming protein and mineral-phytic acid complexes thereby decreasing protein and
57 mineral bioavailability [7]. Phytate is associated with nutritional diseases such as rickets and
58 osteomalacia in children and adult, respectively. Tannins are water soluble phenolic compounds with
59 a molecular weight greater than 500 and with the ability to precipitate proteins from aqueous solution.
60 They occur in all vascular plants. Tannins bind to proteins making them bio-unavailable [8]. This work
61 seeks to quantify both the nutrients and antinutrients in these plants samples.

62 Heavy metals such as arsenic, cadmium, lead are toxic to cells [9], thus it is important to
63 determine their levels in especially in plant-based foods and diets. In general, information on edibility
64 and therapeutic properties of wild plants is scanty but data on their nutritional composition and mineral
65 content is negligible [10]. Manganese is an essential trace element, which plays an important role as a
66 cofactor for many enzyme systems such as hexokinase and superoxide dismutase. At high level
67 however, it can cause damage to the brain [11]. Magnesium is another nutrient required in the plasma
68 and extra cellular fluid, where it helps in maintaining osmotic equilibrium. It is required in many
69 enzyme-catalysed reactions, especially those in which nucleotides participate where the reactive
70 species is the magnesium salt (eg $MgATP^{2-}$). Lack of Mg is associated with abnormal irritability of
71 muscle and convulsions whereas excess level is implicated in depression [12].

72 Iron is necessary for the formation of haemoglobin and also plays an important role in oxygen
73 and electron transfer in human body [13], also in the functioning of the central nervous system as well
74 as oxidation of carbohydrates, proteins and fats [14]. Cadmium is a heavy metal that causes both
75 acute and chronic poisoning; adverse effect on kidney, liver, vascular and immune system [15].
76 Chronic exposure to chromium may result in liver, kidney and lung damage [16].

77 Lead causes both acute and chronic poisoning with the kidney and liver as primary targets. It
78 inhibits the catalytic action of δ -amino levulinic acid dehydratase (Porphobilinogen Synthase) in the
79 haem biosynthetic pathway therefore exerting a toxic effect on the vascular and immune system [17].
80
81

82 **MATERIALS AND METHODS**

83 **Materials**

84 **Chemicals and reagents**

85 All reagents and chemicals were of analytical grade from British Drug Houses.

86 **Equipment used**

87 OHUAS (Ohaus Harvard Trip Balance) digital balance, applied biosystems PTH (phenylthiohydantoin)
88 amino acid analyzer and Soxhlet assembly set up.
89

90 Raw samples of the leafy vegetables were collected from different farms in and around Dengi
91 metropolis. They were destalked, washed with distilled water and air dried. Samples were pulverised
92 in ceramic mortar and pestle. This was followed by sieving the samples to obtain fine particles 10g
93 each for the analysis using the EnoDecott machine.

94 **Amino acids analysis**

95 The amino acid profile was determined using the method described by [18]. Where each of the dried
96 samples were defatted (by refluxing 100g of the air-dried powdered samples with 250ml petroleum
97 ether for 4 hours, the resulting residue was then dried and subjected to aqueous extraction
98 using Soxhlet assembly. Extracts were thereafter evaporated and loaded into the “applied biosystems
99 PTH amino acid analyzer” which separated and analysed free acidic, neutral and basic amino acids of
100 the hydrolysate.

101 **Determination of mineral elements content**

102 The minerals content of the different samples was evaluated for Mn, Mg, Fe, Cd, Cr and Pb by dry
103 ashing of dried powdered sample (5g) in a muffle furnace set at 775°C. The ash obtained was
104 dissolved in 5 mL of 20% HCl and analysed using the atomic absorption spectrophotometer at their
105 respective wavelengths of maximum absorption (λ_{max}) thus: 385nm, 285.5nm, 405nm, 357.8nm,
106 582nm, 389.6nm for Mn, Mg, Fe, Cd, Cr and Pb in that order.

107 **Determination of antinutrients**

108 Tannins were quantified according to Bainbridge *et al* [19] total oxalate quantified applying Day and
109 Underwood [20]. Phytate content was determined by the Reddy and Love method [21].

110 **Statistical Analysis**

111 The statistical method employed for all the analysis was one way ANOVA and all results are means of
112 three determinations (\pm SD). $P = .05$ was considered significant.

114 **RESULTS AND DISCUSSION**

115 This research sought to assess the amino acids, antinutrients, phytochemicals and metals
116 composition of five commonly consumed leafy vegetables in Dengi metropolis. Green leafy vegetables
117 constitute an indispensable constituent of human diet in Africa generally and West Africa in particular;
118 the varieties of leafy vegetables utilized are diverse, ranging from leaves of annuals and shrubs to
119 leaves of trees. Leafy vegetables are generally good sources of nutrients, important protective foods,

120 highly beneficial for the maintenance of health and prevention of diseases as they contain valuable
121 food ingredients which can be utilized to build up and repair the body. They are valuable in
122 maintaining alkaline reserve in the body and are valued mainly for their high vitamin, dietary fibre and
123 mineral content[22]. The dark green leaves and deep yellow fruits provide a high amount of carotene,
124 ascorbic acid and micro-minerals which play important roles in nutrient metabolism and slowing down
125 of degenerative diseases [22]. (This paragraph should be in the Introduction section rather than in
126 Materials and Methods. Delete redundant sentences or words when insert in Introduction)
127

128 Eighteen amino acids were analysed in different proportions in the vegetables. The amino
129 acid contents are generally high in all samples with the highest, based on dry weight, observed in
130 *Letadenia hastata* (93.20g/100g protein) others range between 71.41g to 74.62g/100g proteins. Level
131 of Glutamic Acid was the highest amongst other amino acids in all samples (with an average 11.19
132 g/100g proteins) followed by Aspartic Acid with an average of 7.92g/100g proteins whereas Cysteine
133 and Methionine were low with average of 1.37g/100g proteins. All the samples analysed contained
134 high levels of Glutamate with *Letadenia hastata* have the highest level at 12.11g/g proteins. Levels of
135 Cysteine and Methionine in *Balanites aegyptiaca* were 0.78g/100g and 0.78g/100g
136 proteins respectively. The samples contain essential amino acids such as Methionine, Lysine,
137 Leucine, Isoleucine, Tryptophan, Phenylalanine, Valine and Histidine albeit the levels were lower than
138 the non essential amino acids. Leucine stimulates muscle strength and growth, regulate blood
139 sugar level by moderating insulin into the body during and after exercise and can even help prevent
140 depression by the way it acts on neurotransmitters in the brain [23]. Lysine is responsible for proper
141 growth and in the production of carnitine (a nutrient responsible for converting fatty acids into fuel to
142 lower cholesterol). It also helps the body absorb calcium for further bone strength and also aids in
143 collagen production. Methionine helps form cartilage in the body through the use of sulphur. Histidine is
144 involved in transport neurotransmitters to the brain and also helps overall muscle health within each
145 muscle cells. Valine is needed for optimal muscle growth and repair[23].

146 Table 2 shows the mineral content of the vegetables. The permissible limit of iron in edible
147 plants is 20 ppm [24]. Iron is necessary for the formation of haemoglobin and also plays an important
148 role in oxygen and electron transfer in human body [24] and normal functioning of the central nervous
149 system and in the oxidation of carbohydrates, proteins and fats [25], the highest iron content of 2.223
150 ppm was found in the leaves of *Leptadenia hastata* while the leaves of *Balanites aegyptiaca* contain the
151 least iron content of 0.772 ppm, the leaves of *Vitex doniana*, *Hibiscus*
152 *cannabinus* and *Cucurbitapepo* have significantly higher iron content of 1.081, 0.916 and 0.831 ppm,
153 respectively. An average culinary preparation contains about 300g of the fresh leaves of vegetables
154 and this would result in an intake of 30g dry weight leaves per serving portion. Therefore 1.081, 0.916
155 and 0.831 ppm in *Vitex doniana*, *Hibiscus cannabinus* and *Cucurbitapepo* respectively, will contribute
156 up to, in mg, 3.2, 2.7 and 2.3/serving portion of Fe respectively, to the recommended dietary
157 allowance of Fe (10-15mg/day) [26].

158 Manganese level, in ppm, was found to be 0.133 in *Leptadenia hastata*, 0.065 in *Hibiscus*
159 *cannabinus*, 0.041 in *Balanitesaegyptiaca*, 0.061 in *Cucurbita pepo* and 0.051 in *Vitex doniana* with
160 respectively.

161
162 In table 2, level of Mg, in ppm, was 1.091 in *Cucurbita pepo*, which is the highest compared to
163 others whose content ranged from 0.915 to 1.080. Mg is required in the plasma and extra cellular fluid,
164 where it helps in maintaining osmotic equilibrium [24]. It is required in many enzyme – catalysed
165 reactions, especially those in which nucleotide participate where the reactive species is the
166 magnesium salt, MgATP²⁻. Deficiency of Mg is associated with abnormal irritability of muscle, and
167 convulsions. Excess Mg predisposes to depression [24].

168 Cadmium concentration, in ppm, of the samples ranged from 0.003 in *Hibiscus cannabinus* to
169 0.004 in *Leptadenia hastata*. These values are below the permissible limit of 0.212 in edible plant [24].
170 In medicinal plants however, the permissible limit by WHO is 0.310. The low level of Cd in all the
171 samples means they are safe for consumption.

172 As for Pb content of the samples, *Leptadenia hastata* had 0.002 ppm
173 whereas *Balanitesaegyptiaca* contained 0.055 ppm. These levels in the samples are below the
174 permissible level, 0.43 ppm [27].

175 Results of the phytochemical analysis are presented in Table 3. Antinutritional factors have
176 been shown to limit the use of many plants due to their ubiquitous occurrence as natural compounds
177 capable of eliciting deleterious effect in man and animals [28]. The major antinutritional factors
178 commonly found in green leafy vegetables are phytic acid, oxalic acid and tannins [29]. High levels
179 of phytates and oxalates have been shown to inhibit the absorption and utilization of minerals such as
180 calcium by animals including man [30]. Tannins decrease protein quality by reducing the digestibility
181 and palatability; they interfere with absorption of iron and a possible carcinogenic effect [31].

182 The oxalate content in these vegetables ranged between 0.14 mg/100g in *Vitex doniana* to
183 0.60 mg/100g in *Cucurbita pepo*. These values are below the established toxic level [32]. The
184 phytate level was between 1.7 mg/100g in *Cucurbita pepo* to 4.1 mg/100g in *Leptadenia hastata*. Results
185 obtained are below the toxic level [24]. According to [33] a phytate diet of 1-6% over a long period of
186 time decreases the bioavailability of mineral elements in mono gastric animals.

187 Cyanogens are glycosides of a sugar, sugars and cyanide containing aglycone. Cyanogens
188 can be hydrolyzed by enzymes to release a volatile HCN gas [34]. Excess cyanide ion inhibits the
189 cytochrome oxidase which stops ATP formation and so tissues suffer energy deprivation and death
190 follows rapidly. High level of HCN has been implicated for cerebral damage and lethargy in man and
191 animal [18]. In table 3, the levels were 7.59 % in *Cucurbita pepo* and 4.82% in *Balanitesaegyptiaca*
192 which are below lethal level as indicated on the table 3. As for saponins, level ranged between 11.85
193 % in *Leptadenia hastata* to 15.13 % in *Vitex doniana*. Saponins are glycosides containing polycyclic
194 aglycone moiety of either C₂₇ steroid or C₃₀ triterpenoids attached to a carbohydrate. High saponin
195 level has been associated with gastroenteritis manifested by diarrhoea and dysentery [35].

196 Tannins are water soluble phenolic compounds with a molecular weight greater than 500 and
197 with the ability to precipitate proteins from aqueous solution. They occur in all vascular plants. Tannin
198 binds to proteins making them bio unavailable [36]. From the results, the level obtained was between

199 0.51% in *Balanitesaegyptiaca* and 0.60 % in *Vitex donina*. There was no significant difference in the
 200 Tannin level among the vegetables ($p>0.05$).

201 Alkaloids have been implicated in the inhibition activities of many bacterial species [37]. The
 202 levels of alkaloids in the various leafy vegetables analysed ranges from 16.22% observed in
 203 *Letadenia hastata* and 19.37% in *Vitex doniana*. Plant phenolics such as flavonoids have been shown
 204 to have antioxidant properties and also contribute to their medicinal significance [38]. In this work, the
 205 levels of flavonoids ranged from 9.87 % in *Balanitesaegyptiaca* to 14.71 % in *Vitex*
 206 *doniana*. Consumption of these vegetables would further enhance the capacity of the cell to mop up
 207 the highly reactive oxygen radicals generated due to oxidative metabolic reactions that occur in cells.
 208

209 **TABLE 1:** Amino acid composition of five leafy vegetables in Dengi metropolis.

AMINO ACIDS g/100g proteins	<i>Cucurbita pepo</i>	<i>Vitex doniana</i>	<i>Hibiscus cannabinus</i>	<i>Letadenia hastata</i>	<i>Balanites aegyptiaca</i>	MEAN	S.D	C.V %
Leucine	8.14	6.48	7.29	8.34	6.48	7.35	0.9	12.03
Lysine	4.03	3.63	3.97	5.54	4.37	4.31	0.7	17.11
Isoleucine	3.3	3.01	2.94	4.78	3.4	3.49	0.8	21.47
Phenylalanine	4.17	4.43	4.34	5.58	4.7	4.64	0.6	12
Tryptophan	2.25	1.26	2.1	2.86	1.12	1.92	0.7	37.78
Valine	4.5	3.59	4	5.61	4.09	4.36	0.8	17.69
Methionine	1.12	1.23	1.71	2	0.8	1.37	0.5	34.95
Proline	2.84	3.14	2.84	4.57	2.23	3.12	0.9	27.96
Arginine	4.99	6.11	5.33	7.83	4.99	5.85	1.2	20.47
Tyrosine	2.75	3.09	3.09	4.13	3.09	3.23	0.5	16.23
Histidine	2.23	2.17	2.39	2.11	2.65	2.31	0.2	9.39
Cysteine	1.09	1.21	1.57	2.18	0.78	1.37	0.5	39.22
Alanine	3.71	4.4	3.94	5.4	4.28	4.35	0.7	14.95
Glutamic acid	9.99	11.28	11.58	12.11	10.97	11.19	0.8	7.06
Glycine	4.08	4.11	3.75	4.46	3.89	4.06	0.3	6.61
Threonine	2.88	2.99	3.27	3.38	2.88	3.08	0.2	7.51
Serine	2.99	3.29	3.13	3.83	3.25	3.3	0.3	9.69
Aspartic Acid	8.31	8	7.38	8.49	7.44	7.92	0.5	6.33
TOTAL	73.37	73.42	74.62	93.2	71.41	77.2	9	11.68

210

211

212

213

214

215

216

217

218

219

220 **TABLE 2:** Levels of metals in five leafy vegetables in Dengi metropolis

SAMPLES	Pb	Cd	Mg	Mn	Fe
concentration(ppm)					
<i>Cucurbitapepo</i>	0.0229±0.004	0.003±0.001	1.0871±0.002	0.0557±0.001	0.7997±0.003
<i>Vitexdoniana</i>	0.0302±0.002	0.0036±0.001	1.1371±0.010	0.0499±0.001	1.0799±0.100
<i>Hibiscus cannabinus</i>	0.0396±0.021	0.029±0.001	1.1255±0.004	0.0653±0.004	0.9159±0.012
<i>Letadenia hastata</i>	0.0021±0.001	0.037±0.010	0.9951±0.010	0.1331±0.000	2.2231±0.900
<i>Balanitesaegyptiaca</i>	0.0552±0.012	0.041±0.001	1.0753±0.002	0.0405±0.001	0.7723±0.123
<i>permissible levels</i>	0.10	0.05		-	-

221 Values represent means of triplicate determination± SD

222 Source for permissible limits: FAO (Mg/L) recommended maximum Concentration of trace elements
223 for vegetables (1985).

224

225 **TABLE 3:** Phytochemical content of five leafy vegetables in Dengi metropolis

SAMPLES	Tannins (%)	Oxalate (mg/100g)	Phytate (mg/100g)	Saponins (%)	Cyanogenic Glycosides (mg/100g)	Alkaloids (%)	Flavonoids (%)
<i>Cucurbitapepo</i>	0.53± .012	0.60±0.20	1.70±0.01	12.49±1.20	7.59±1.20	17.45±2.51	13.72±1.21
<i>Vitexdoniana</i>	0.60±0.12	0.14±0.01	2.10±0.21	15.13±1.50	5.33±1.11	19.37±1.23	14.71±2.30
<i>Hibiscus cannabinus</i>	0.59±0.13	0.32±0.12	3.14±0.12	13.35±2.35	4.93±2.13	17.10±2.50	10.38±1.30
<i>Letadenia hastate</i>	0.55±0.21	0.22±0.03	4.10±0.01	11.85±1.85	6.21±1.20	16.22±1.61	13.22±1.23
<i>Balanitesaegyptiaca</i>	0.51±0.13	0.31±0.01	3.08±0.02	13.48±3.12	4.82±1.30	18.13±2.31	9.87±1.32

226 The values represent the mean of three determinations± SD.

227

228 CONCLUSION AND CONTRIBUTION TO KNOWLEDGE

229 From these results, leaves of the vegetables constitute rich sources of amino acids and mineral
230 elements. The vegetables are therefore rich alternatives (and/or supplements) which can replenish
231 the cellular requirements of the nutrients. Also, samples contain antinutritional factors such as
232 oxalates and phytates but at low levels. The content of flavonoids and alkaloids in appreciable amounts
233 in the samples is critical given the therapeutic/medicinal use of phytochemicals. Hence, leafy
234 vegetables could contribute to the alleviation of protein malnutrition and micronutrient deficiencies if
235 consumed.

236 REFERENCES

237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293

1. Ryan, MF. The role of magnesium in clinical biochemistry: an overview. *Ann Clin Biochemistry*. 1991; 28:19-26.
2. Hilou A, Nacoulma OG, Guiguemde TR. *In vivo* antimalarial activities of extract from *Amaranthus spinosus* L and *Boerhaavia erecta* L. *J. Ethnopharmacol*. 2006; 103: 236-240.
3. Yiridoe EK, Anchirinah VM. Garden production systems and food security in Ghana: Characteristics of traditional knowledge and management systems. *Renew. Agric. Food Syst*. 2005; 20: 168-180.
4. Kubmarawa DI, Andenyang FH, Magomya AM. Proximate composition and amino acid profile of two non-conventional leafy vegetables (*Hibiscus cannabinus* and *Haematostaphis barberi*). *Pakistan Journal of Nutrition*. 2013; 12 (10): 949-956 ISSN 1680-5194.
6. Nkafamiya II, Manji AJ. A study of cyanogenetic glucoside contents of some edible Nuts and Seeds. *J. Chem. Soc*. 2006; Niger 31 (1 and 2) 12-14.
7. Nkafamiya II, Modibbo UU, Manji AJ, Haggai D. Nutrient Content of Seeds of Some Wild Plants. *Afr. J. Biotechnol*. 2007; 6(15): 1665-1669.
8. Fasusi AO. Nutritional potentials of some tropical vegetable meals. Chemical characterization and functional properties. *Afr. J. Biotechnol*. 2006; 5(1): 49-53.
9. Bagepallis S, Narasinga R, Tatinemi P. Tannin contents of foods commonly consumed in India and its influence on ionisable iron. *J. Sci. Food Afric*. 1993; 33:89-96.
10. Schumacher M, Bosque, MA, Domingo JL, Corbella J. Dietary intake of lead and cadmium from food in Tarragona Province, Spain. *Bull. Env. Cont. Toxicol*. 1991; 46:320-328.
11. Aloskar LV, Kakkar KK, Chakra OJ. Second Supplement to Glossary of Indian Medicinal Plants with Active Principles, 1992. Part-I (A-K), NISC, CSIR, New Delhi, pp: A.O.A.C. (Association of Official Agriculture Chemists).
12. Aschner JL, Aschner M. Nutritional aspect of manganese homeostasis. *Molecular aspects of Medicine*. 2005
13. Kaya I, Incekara N. Contents of some wild plants species consumed as food in Aegean region. 2010; *J. Turk. Weed Scie* 3:56-64.
15. Adeyeye EI, Otokiti, MKO. Proximate composition and some nutritionally valuable minerals of two varieties of *Capsicum annum* (Bell and Cherry peppers). *Discovery and Innovation* 11:75-81 1999; Aegean region. *J. Turk. Weed Scie* 3:56-6.
16. Heyes RB. The Carcinogenicity of metals in humans: *Cancer Causes Control* 8:371-385. 1997.
17. Zayed AM, Terry N. Chromium in the environment: factors affecting biological remediation. *Plant. Soil*. 2003; 249:139-156.
18. Bainbridge ZK, Tomlins A, Westby A. Analysis of condensed tannins using acidified vanillin. *J. Food Sci. Agric*. 1996: 29: 77-79.
19. Day RA, Underwood AL. *Quantitative analysis*. 5th ed. Prentice Hall, 701. 1986.
20. Phytate determination Reddy MB, Love M (1999). The impacts of food processing on the nutritional quality of vitamins and minerals. *Adv. Exp. Med. Bio*. 459: 99-106.

- 294 21. Olubunmi AO, Olaofe O, Olunayo RA. Amino Acid Composition of Ten Commonly Eaten
295 Indigenous Leafy Vegetables of South-West Nigeria. *World Journal of Nutrition and Health*. 2017;
296 3(1).
- 297 22. Guoyao, W. Functional Amino Acids in growth, reproduction and health. *Advances in Nutrition*.
298 2010; 1(1): 31-37.
- 299 23. FAO/WHO. Contaminants. In: *Codex Alimentarius* (1st ed, XVII), FAO/WHO, 1984.
300
- 301 24. FAO/WHO. United Nations Food and Agriculture Organization/World Health Organization: Fruit
302 and Vegetables for Health. Report of a Joint FAO/WHO Workshop, Kobe, Japan, 2004.
- 303 25. Allison RG, Margot MP. Identifying the Threshold of Iron Deficiency in the Central Nervous System
304 of the Rat by the Auditory Brainstem Response. *ASN Neuro*. 2015; 7(1): 17-21.
- 305 26. Hurrell R, Egli I. Iron bioavailability and dietary reference values. *Am J Clin Nutr* 2010;91:1461S-
306 7S.
- 307 27. Zurera G, Estrada B, Rincón F, Pozo R. Lead and cadmium contamination levels in edible
308 vegetables. *Bull Environ Contam Toxicol* 1987; 38: 805-812.
- 309 28. Habtumu F, Fecadu G. Antinutritional factors in plant foods: Potential health benefits and adverse
310 effects. *International Journal of Nutrition and Food Sciences* 2014; 3(4): 284-289.
- 311 29. Kaushalya G, Wagle DS. Nutritional and antinutritional factors of green leafy vegetables. *Journal*
312 *of Agricultural and Food Chemistry* 1988; 36(3). DOI: 10.1021/jf00081a016.
- 313 30. Norhaizan ME, Nor F AW. Determination of Phytate, Iron, Zinc, Calcium Contents and Their Molar
314 Ratios in Commonly Consumed Raw and Prepared Food in Malaysia. *Mal J Nutr* 2009; 15(2): 213 –
315 222.
316
- 317 31. Donald ES, Collins WB, Thomas AH, Cassara NE, Carnaha AM. The impact of tannins on protein,
318 dry matter, and energy digestion in moose (*Alces alces*). *Canadian Journal of Zoology* 2010; 88: 977-
319 987.
320
- 321 32. American Conference of Governmental Industrial Hygienists. Documentation of the Threshold
322 Limit Values and Biological Exposure Indices. 5th ed. Cincinnati, OH: American Conference of
323 Governmental Industrial Hygienists, 1986; p 451.
- 324 33. Grases F, Rafaei MP, Antonis C. Dietary Phytate and Interactions with Mineral Nutrients. In book:
325 *Clinical Aspects of Natural and Added Phosphorus in Foods* 10.1007/978-1-4939-6566-3_12. 2017.
- 326
- 327 34. Ewa J, Żaneta P, Sylwia N, Jacek N. Cyanides in the environment—analysis—problems and
328 challenges. *Environ Sci Pollut Res Int*. 2017; 24(19): 15929–15948.
- 329 35. Michael R R, Ka IT, Melody KB. Prevention of rotavirus infections *in vitro* with aqueous extracts of
330 *Quillaja saponaria* Molina. *Future Med Chem*. 2010; 2(7): 1083–1097.
- 331 36. Ann EH. Tannin—Protein Interactions; Phenolic Compounds in Food and Their Effects on Health.
332 Department of Chemistry, Miami University, Oxford, 1992; Copyright © 1992 American Chemical
333 Society. ISBN13: 9780841224759; 236–247.
- 334 37. Tim TPC, Benjamert C, Andrew JB. Alkaloids: An overview of their antibacterial, antibiotic-
335 enhancing and antivirulence activities. *International Journal of Antimicrobial Agents* 2014; 44(5).
- 336 38. Sofna DSB, Nina A. Antioxidant properties of Flavonoids. *Med J. Indones*, 2014; 23 (4): 239-242.