Original Research Article

Soil fertility as influenced by incorporation of K enriched Azolla

Abstract

Use of chemical fertilizer in judiciously without organic manure has created many soil health problems. ~~Therefore, alternate sources of N has to be evolved to supply crop demanded N with less or no environmental pollution.~~Biological Nitrogen Fixation (BNF) is a major source of fixed N for plant life and to sustain production and productivity of agricultural crops. All other biofertilizers simply solubilize or mobilize the nutrients that are already present in soils. Whereas the *Azolla* is unique in the sense that it acts as host to the N-fixing cyanobacteria after which it is used virtually as a green manure. An incubation experiment was conducted by growing *Azollae filiculoides* with 2 agriculturally important potassic fertilizers (Potassium Chloride, Potassium shulphate) as main plot in seven concentrations (0, 5, 10, 20, 30, 40 and 50 ppm of K) as sub-plots laid down in split plot design replicated three times. The collected azolla was incorporated with soil at 10 t/ha and maintained at two moisture condition such as 60 and 100 percent and assessed soil fertility by estimating various available plant nutrients and organic carbon status. ~~Soil fertility was influenced by the /humid substances formed during the decomposition ofAzolla~~.The mean organic carbon content of the soil was 0.657 and 0.525% by K enriched azolla at 60 and 100% moisture contents respectively.. The available N content ranged from 216.2 to 327.3 and 191.1 to 285.3 kg/ha from 0 to 40 ppm of K concentration at 60 and 100 % moisture respectively because the Azolla ~~/~~ had a high N content released into the soil after decomposition. Azolla also contributed to the supply of phosphorus, potassium, sulfur, zinc, iron and molybdenum in sufficient amounts in addition to other micronutrients besides addition of nitrogen. Among the various concentration, 40 and 50 ppm K were significantly maintained higher and equal soil available P status of 75.17 and 77.33 kg / ha

Comment [EA1]: No need of referencing in the abstracts

respectively. The fertilizer, K_2SO_4 with 30, 40 and 50 ppm at 60 % moisture content and 40 and 50 ppm of $K_2SO_4$and 40 ppm of KCl produced statistically higher and equal available K in azolla incorporated soil maintained at 100 % moisture. The soil biological health, mineralization and consequent increase in nutrient status by the application of K enriched Azolla was more under 60 % soil moisture content than fully saturated soil.

Keywords: ~~between 4-8 words~~ **Biofertilizer, Azolla, K-enrichment, Soil fertility, Organic carbon and Available NPK.**

1. Introduction

Intensive crop production is the demand of time to feed the vast growing population in India. This has created a pressure to use more chemical fertilizer. Use of chemical fertilizer injudiciously without organic manure has created many soil health problems like low fertilizer use efficiency, poor soil physical condition, reduced water holding capacity, degraded rhizospharic properties, and low fertility (Awodun, 2008). Rice crops remove around 16-17 kg N for the production of each ton of paddy cultivation (Sahrawat, 2000). Most of the rice soils of the world are deficient in N, so, fertilizer N applications are required to meet its N demand (Datta and Buresh, 1989). Generally, urea is applied as the N source for rice production. But the efficiency of added urea-N is very low, due to denitrification, NH_3 volatilization and leaching (De Datta and Buresh 1989). Therefore, alternate sources of N has to be evolved to supply crop demanded N with less or no environmental pollution. The demand of fertilizers and manures is increasing at the present days scenarios to maximize the crop production. Biological Nitrogen Fixation (BNF) is a major source of fixed N for plant life and to sustain production and productivity of agricultural crops. Estimate of global terrestrial showed that, the BNF ranged from 100 to 290 million tonnes of N/year. Of this, 40–48 million tonnes is estimated to be biologically fixed in agricultural crops and fields. BNF is one of the natural sources of nitrogen for rice and *Azolla*-cyanobacteria biomass has

Comment [EA2]: Same like the abstract introduction

Comment [EA3]: Reference cannot be CROSS check

Comment [EA4]: Important point but with no reference

been identified as potential source of nitrogen. The integrated nutrient management is to maintain or adjust plant nutrient supply to achieve a given level of crop production by optimizing the benefits from all possible sources of plant nutrients (Subba Rao, 2005). Organic manures are considered to play a significant role in nutrient contribution. The use of organic fertilizer is a way to improve soil fertility. Azolla can be used as organic fertilizer (Syamsiyah *et al*., 2015). Subedi and Shrestha (2015) explained that, Azolla does not only increase the productivity of rice but also improve the long-term soil fertility. All other biofertilizers simply solubilize or mobilize the nutrients that are already present in soils. Whereas the *Azolla*is unique in the sense that it acts as host to the N-fixing cyanobacteria after which it is used virtually as a green manure. In the process, it adds not only the biologically fixed N but also the other nutrients absorbed from the soil and present in its biomass. Against the total anticipated biofertilizers demand of 1 million tonne in the country, the current supply position is very low (<10 000 tonnes). The present investigation studied soil fertility improvement by azolla grown under varied K fertilizer commonly used in agriculture.

2. Materials and Methods

2.1. Study site

Comment [EA5]: Basic information about the study/experimental site, meteorological and soil information

An incubation experiment was conducted by growing *Azollae filiculoides* with 2 agriculturally important potassic fertilizers (Potassium Chloride, Potassium shulphate) as main plot in seven concentrations (0, 5, 10, 20, 30, 40 and 50 ppm of K) as sub-plots laid down in split plot design replicated three times. The experimental site was Rice Research Station, Ambasamudram located between 8º 42' N and 77º 28' E with an altitude of 64.8 M above mean sea level. The mean annual rainfall received was 913 mm. The experimental soil was acidic in reaction (pH 5.84) and free from salinity (0.08 dS m^{-1}) with sandy clay in texture. The organic carbon content was 0.56 %. The soil was low in available nitrogen (210 kg ha^{-1}), high in available P (24.5 kg ha^{-1}) and medium in available K (150 kg ha^{-1}).

One gram of Azolla fern was grown in a tray with a dimension of 23 x 15x 6 cm^3 filled with 1.5 litres of potassic solutions (Plate 1) and the fern was collected on 7th, 15th, 30th, 60th, 90th and 120th day after incubation / culturing, rinsed with distilled water and analysed for various biometric and biochemical parameters. The collected Azolla was incorporated with soil at 10 t/ha and maintained at two moisture condition such as 60 and 100 percent. The Azolla incorporated soil was collected after 30 days of incubation, processed and analyzed for various available plant nutrients and organic carbon status. Organic carbon present in soil /wasoxidied/oxidized by chromic acid ($K_2Cr_2O_7$) in the presence of conc. H_2SO_4. Potassium dichromate on reaction with H_2SO_4 provided nascent oxygen which combined with carbon to form CO_2. The excess chromic acid left unused by the organic matter /was determined by back titration with 0.5 N ferrous sulphate or ferrous ammonium sulphate using diphenylamine indicator (Walkley and Black, 1934). Available nitrogen in the soil was estimated by alkaline permanganate method (Subbiah and Asija, 1956). Available phosphorus extracted with 0.03 N NH_4F and 0.025 N HCl. The amount of P extracted wastreated with ammonium molybdate and antimony potassium tartarate and developed colour with ascorbic acid. The intentensity of blue colour /was determined colorimetrically at 660nm (Bray*et al.*, 1945). The soil was leached with neutral normal ammonium acetate and the K^+ ions in the exchange sites were replaced by NH_4^+ ions. The K^+ ions in solution was then determined with the flame photometer (Standford and English (1949).

Data Collection The soil samples were collected from the incubation bottle ffter 90 days of incubation were shade dried, processed and sieved through 0.5 and 2 mm sieve for estimating organic carbon and available NPK respectively.

Statistical Analysis The data were analysed statistically using computer software IRRISTAT (IRRI, 1993). Differences among the mean values of the treatment were compared by the LSD test when the F test from the analysis of variance was significant at the

Comment [EA6]: Any photo to depicts the setup of materials and methods well to aid in future repetition by others

Comment [EA7]: References cannot be CROSS check

Comment [EA8]: What parameter(s) were under study;how was the data collected

Comment [EA9]: How was the data analysed and with what statistical tool?

Formatted: Indent: First line: 0", Space After: 0 pt, Line spacing: Double, No widow/orphan control, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers

Formatted: Font: 12 pt

p= 0.005 level.

3. Results

3.1.Organic carbon

The effect of K fertilizer, its concentration and their interaction on the organic carbon content of Azolla incorporated into the soil at 60 %and 100 % moisture content is presented in Table 1 and 2. The concentration of K fertilizer alone influenced the organic carbon content of the Azolla incorporated at both 60 % and 100 % moisture content.The mean organic carbon content of the soil was 0.657 and 0.525% by K enriched Azollaat 60 and 100% moisture contents respectively. Azollaenriched with 40ppm K solution recordedsignificantly higher soil organic carbon content of 0.763and 0.623% respectively at 60 and 100 % soil moisture content (Table 1) which was 42 and 56% more than the Azolla grown under 0ppm K. However, it was on par with 50 and 30ppm of K solution in both the moisture-maintained soil.The 20ppm K registered 0.697 and 0.567% of organic carbon content in soil maintained at 60 and 100% moisture content which were 9 to 10% less than the highest organic carbon maintained by 40ppm of K.

3.2.Available N

The main effect ofAzolla grown under K fertilizer and its concentration alone significantly influenced the available N content at both 60 and 100% moisture content. On an average 291.9 and 256.9 $kgha^{-1}$ of available N was maintained by the incorporation of K enrichedAzollain soilmaintained at 60 and 100% moisturecontent respectively (Table 1). Among the Kfertilizer, K_2SO_4 significantly maintained higher available N status of 297.2 and 262.2 $kgha^{-1}$ which was 4 to 5% higher than the KClwhich registered 286.6 and 251.6 $kgha^{-1}$ of soil available nitrogen content at 60 and 100% moisture content respectively. The available

N content ranged from 216.2 to 327.3 and 191.1 to 285.3kgha^{-1}from 0 to 40ppm of K concentration at 60 and 100% moisture content respectively. Though, the 40 and 50ppm of K were on par with each other, they however, maintained significantly higher available N status of 327 and 285kg/ha kgha^{-1} at 60 and 100% moisture content respectively and it was 4.0% higher than 30ppm of K enriched Azolla.

3.3.Available P

The main and interaction of fertilizer and their concentration significantly influenced the available P status of Azolla incorporated into the soil maintained at 100% moisture content where as the main effect was alone significantly influenced the available P content at 60% moisture condition. On an average, the K enriched Azollaregistered59.63 and 42.13 kgha^{-1}ofavailable P at 60 and 100% soil moisture respectively. Among the fertilizers used, the K_2SO_4 maintained higher available P content of 61.71 and 44.21 kgha^{-1} in the soil incorporated with K enriched Azolla at 60 and 100 % moisture content respectively followed by KCl which registered the available P content of 57.54 and 40.04 kgha^{-1}(Table 1). Among the various concentration, 40 and 50ppm K was significantly maintained higher and equal soil available P status of 75.17 and 77.33 kgha^{-1} respectively which was 8.6% more than the 30ppm K (71.17 kgha^{-1}) at 60% moisture content.

Table 1. Soil fertility of K enriched Azolla as influenced by main effect of fertilizer and concentration at 60 and 100% moisture content

Soil Fertility Parameters	Organic Carbon (%)		Available N (kgha^{-1})		Available P (kgha^{-1})		Available K (kgha^{-1})	
Moisture	60%	100%	60%	100%	60%	100%	60%	100%
Fertilizer								
F_1 (KCl)	0.644	0.513	286.6^b	251.6^b	57.54^b	40.04^b	205.1^b	160.9^b
F_2 (K_2SO_4)	0.670	0.538	297.2^a	262.2^a	61.71^a	44.21^a	215.5^a	171.3^a
Mean	**0.657**	**0.525**	**291.9**	**256.9**	**59.63**	**42.13**	**210.3**	**166.1**
SEd	0.006	0.006	1.17	1.16	0.85	0.85	1.02	1.02

CD (0.05)	NS	NS	5.02	5.0	3.68	3.68	4.40	4.4
				Concentration				
C1 (0 ppm)	0.518^{f}	0.398^{e}	216.2^{g}	191.2^{g}	34.50^{g}	22.50^{f}	157.8^{g}	125.8^{f}
C2 (2 ppm)	0.552^{e}	0.422^{e}	266.8^{f}	236.8^{f}	44.50^{f}	29.50^{e}	172.5^{f}	135.5^{e}
C3 (5 ppm)	0.587^{d}	0.457^{d}	280.0^{e}	250.0^{e}	51.67^{e}	36.67^{d}	190.2^{e}	153.2^{d}
C4 (10 ppm)	0.632^{c}	0.502^{c}	296.2^{d}	259.2^{d}	57.83^{d}	39.83^{c}	204.5^{d}	162.5^{c}
C5 (20 ppm)	0.697^{b}	0.567^{b}	307.5^{c}	270.5^{c}	64.83^{c}	46.83^{b}	219.3^{c}	177.3^{b}
C6 (30 ppm)	0.755^{a}	0.622^{a}	314.0^{b}	277.0^{b}	71.17^{b}	53.17^{a}	238.7^{b}	189.7^{a}
C7 (40 ppm)	0.763^{a}	0.623^{a}	327.3^{a}	285.3^{a}	75.17^{a}	53.17^{a}	250.0^{a}	195.0^{a}
C8 (50 ppm)	0.753^{a}	0.613^{a}	327.2^{a}	285.2^{a}	77.33^{a}	55.33^{a}	249.5^{a}	189.5^{a}
Mean	**0.657**	**0.525**	**291.9**	**256.9**	**59.63**	**42.13**	**210.3**	**166.1**
SEd	0.011	0.013	2.96	3.0	1.08	1.07	2.89	2.89
CD (0.05)	0.023	0.235	6.06	6.1	2.20	2.19	5.93	5.92

Table 2. Soil fertility of K enriched Azolla as influenced by interaction effect between fertilizer Vs. concentration at 60 and 100% moisture content

Soil Fertility Parameters	Organic Carbon (%)		Available N (kgha^{-1})		Available P (kgha^{-1})		Available K (kgha^{-1})	
Moisture	60%	100%	60%	100%	60%	100%	60%	100%
F_1C_1	0.510	0.390	214.3	189.3	33.33^{j}	21.33^{j}	156.3^{f}	124.3^{h}
F_1C_2	0.543	0.413	263.7	233.7	44.00^{i}	29.00^{i}	171.3^{e}	134.3^{f}
F_1C_3	0.577	0.447	278.0	248.0	51.67^{h}	36.67^{h}	187.3^{d}	150.3^{e}
F_1C_4	0.617	0.487	287.0	250.0	56.67^{g}	38.67^{gh}	195.3^{d}	153.3^{e}
F_1C_5	0.670	0.540	302.7	265.7	61.00^{f}	43.00^{f}	207.0^{c}	165.0^{d}
F_1C_6	0.727	0.593	307.3	270.3	65.00^{e}	47.00^{e}	226.0^{b}	177.0^{c}
F_1C_7	0.757	0.617	319.7	277.7	73.00^{c}	51.00^{d}	249.3^{a}	194.3^{ab}
F_1C_8	0.753	0.613	320.3	278.3	75.67^{bc}	53.67^{cd}	248.3^{a}	188.3^{b}
F_2C_1	0.527	0.407	218.0	193.0	35.67^{j}	23.67^{j}	159.3^{f}	127.3^{h}
F_2C_2	0.560	0.430	270.0	240.0	45.00^{i}	30.00^{i}	173.7^{e}	136.7^{f}
F_2C_3	0.597	0.467	282.0	252.0	51.67^{h}	36.67^{h}	193.0^{d}	156.0^{e}
F_2C_4	0.647	0.517	305.3	268.3	59.00^{fg}	41.00^{fg}	213.7^{c}	171.7^{cd}
F_2C_5	0.723	0.593	312.3	275.3	68.67^{d}	50.67^{d}	231.7^{b}	189.7^{b}
F_2C_6	0.783	0.650	320.7	283.7	77.33^{ab}	59.33^{a}	251.3^{a}	202.3^{a}
F_2C_7	0.770	0.630	335.0	293.0	77.33^{ab}	55.33^{bc}	250.7^{a}	195.7^{ab}
F_2C_8	0.753	0.613	334.0	292.0	79.00^{a}	57.00^{ab}	250.7^{a}	190.7^{b}
Mean	0.657	0.525	291.9	256.9	59.63	42.13	210.3	166.1
SEd								
F at C	0.016	0.017	4.08	4.09	1.66	1.65	3.96	3.96
C at F	0.016	0.016	4.18	4.19	1.52	1.51	4.09	4.09
CD (0.05)								
F at C	NS	NS	NS	NS	4.39	4.39	8.71	3.9
C at F	NS	NS	NS	NS	3.11	3.11	8.38	4.1

But in 100% soil moisture contentat 30, 40 and 50ppm, K was significantly maintained higher and on par soil available P status followed by 20ppm K. The lowest available P of 34.5 and 22.5 kgha^{-1} was registered by soil incorporated with Azolla enriched with 0ppm of K at 60 and 100% moisture content. Under the interaction between K fertilizer and its concentration, K_2SO_4 at 50 and 30 ppm were significantly superior in maintaining higher available P status at 100 % moisture content followed by 40 ppm $K_2SO_4$and it was on par with 50 ppm of KCl (Table 2). However, the 0 ppm of both the fertilizer produced lowest available P content (21.33 and 23.67 kgha^{-1}) of Azolla incorporated soil.

3.4.Available K

The available K content of soil incorporated with K enriched Azolla was significantly influenced by the main and interaction effect of K fertilizer and its concentration. Irrespective of the fertilizer and their concentration about 210.3 and 166.1 kgha^{-1}of available K was maintained by the incorporation of K enriched Azolla in soil at 60 and 100 % moisture respectively (Table 1).Among the K fertilizer, K_2SO_4 was superior in maintaining available K content (215.5 and 171.3 kgha^{-1}) in Azolla incorporated soil which was 5 -7 % more than the KCl (205.1 and 160.9 kgha^{-1}). With respect to concentration of K, 40 and 50 ppm of K at 60 % moisture and 30.40 and 50 ppm of K at 100 % moisture registered significantly higher and equal available K in Azolla incorporated soil followed by 30 and 20 ppm of Kat 60 and 100 % moisture respectively. The Azolla grown under 0 ppm of K registered the lowest available K content of 157.8 and 125.8 kgha^{-1}at 60 and 100 % moisture content respectively. Under the interaction between K fertilizer and its concentration, K_2SO_4 with 30, 40 and 50 ppm at 60 % moisture content and 40 and 50 ppm of$K_2SO_4$and 40 ppm ofKCl produced statistically higher and equal available K in Azolla incorporated soil maintained at 60 and 100 %

moisture respectively (Table 2).The 0 ppm of both the fertilizer registered the lowest available K content (159.3 and 156.3 kgha^{-1}) in soil incorporated with Azolla grown under these concentrations at both moisture content.

4. Discussion

Soil fertility is influenced by the humicsubstances formed during the decomposition ofAzolla (Bhardwaj and Gaur 1970). Incorporation of Azolla enhanced the soil nutrients availability by their biological activity. The decomposed organic matter from Azolla biomass played an active role in the development of microbial population. Similarly, Kannaiyan and Subramani (1992) showed the increased of cellulolyticand urea hydrolyzing activities in addition to significant increase in the population of heterotrophic bacteria by the addedAzolla. Soil incorporation of Azollaalso increased urease and phosphatase activity (Thanikachalam*et al*.1984).Azolla contains macro, secondary and micronutrients that is important for quality rice production (Kumar and Shahi, 2016).Sutanto (2002) stated that, the use of 7.5 ton ha^{-1}Azolla to paddy field increased soil organic matter (C-organic) 0.09 times of control (without Azolla).Syamsiyah*et al*. (2015) proved that, application of Azolla at 2 ton ha^{-1} could increase the organic matter up to 3.69% compare to the field without Azolla.The increasing of organic C is caused by the high content of organic C in Azolla. The incorporated Azolla into soil would soon be mineralized. Watanabe *et al*, (1991) stated that 90% of Azolla was decomposed in 4 weeks and releases humic substances in to the soil.The increase in grain yield might be due to build up of soil organic carbon and more nitrogen through the integrated use of NPK and green manuring with Azolla.

Biological Nitrogen Fixation (BNF) is a major source of fixed N for plant lifeand to sustain production and productivity of agricultural crops. Estimate of global terrestrial showed that the BNF ranged from 100 to 290 million tonnes of N/ year. Of this, 40–48 million tonnes is

203 estimated to be biologically fixed in agricultural crops and fields. BNF is one of the natural
204 sources of nitrogen for rice and *Azolla*-cyanobacteria biomass has been identified aspotential
205 source of nitrogen. The glutamate synthase enzyme dominated in ammonia assimilation
206 followed by glutamine synthates and glutamate dehydrogenase in Azolla (Fig, 1).The activity
207 of all the three enzymes were more at the 40 ppm of K followed by 30 and 50 ppm of K.
208 Incorporation of 40 ppm Keither as KCl and K_2SO_4 incubated Azolla enhanced ammonia
209 assimilation and improved soil fertility (Fig. 2) on 30 dayswhich may help to reduce nitrogen
210 demand for rice crop(Muruganayaki, 2017).



Figure 1. Nitrogen assimilating enzymes as influenced by K fertilizer



Figure 2. Nitrogen fixing enzyme as influenced
by concentration of fertilizers

211

212 According to Roy, (1981), incorporation of 6 t of *Azolla* ha^{-1}, equivalent to 36 kg of N ha^{-1}
213 before planting and incorporation of 1 ton ha^{-1}*Azolla*, equivalent to 24 kgN ha^{-1} after 3-4
214 days of planting. Fogg *et al*.(1973) have found that *Azolla*and cyanobacteria bio-fertilizer can
215 add as much as 30-120 kg N ha^{-1} per crop. Many researchers have considered cyanobacteria
216 as apromising source of nitrogen in tropical rice soils.(Valiente*et al*., 1998) investigated the
217 potential contribution of N_2 fixation by indigenous cyanobacteria to rice soil with increasing
218 amount of fertilizers. Ventura *et al.* (2012) concluded that about 50% of the N inAzolla was

mineralizedafter 2 weeks of incubation with more than 3% N content. Use of Azolla as green manuring and as intercrop proved beneficial and significant result over control in respect of P content. The highest available phosphorus (29.6 kg ha^{-1}) was recorded with 100% NPK + green manuring of Azolla (Kumar and Shahi, 2016).Azolla also contributes to the supply of Phosphorus, Potassium, Sulfur, Zinc, Iron and Molybdenum in sufficient amounts in addition to other micronutrients besides addition of Nitrogen. Similarly, the soil biological health due toapplication of Azolla has resulted in improving mineralization and consequent increase in the soil microbial status (Yadav *et al*. 2014).

5. Conclusion

Soil biological health, mineralization and consequent increase in nutrient status by the application of K enriched Azolla was more under 60 % soil moisture content than fully saturated soil. Further, enrichment of Azolla with 40 ppm of K_2SO_4 enhanced nutrient content in Azolla and ~~subsequently~~ in soil.

References

Awodun~~,~~ M~~.~~ A. ~~2008.~~ Effect of Azolla (*Azolla species*) on physicochemical properties of the soil**.** ***World J.Agril. Sci***.2008;~~,~~**4** (2)**:** 157-160.

Bhardwaj K~~.~~ K~~.~~ R~~.~~, Gaur A~~.~~ C. ~~1970.~~ Effect of humic and pulvic acid on growth and efficiency of nitrogen fixation by *Azotobacterchroococcum*, ***Folia Microbiology***~~,~~ 1970; **15**: 364.

Bray~~,~~ R~~.~~A,~~, and~~ Kurt~~,~~ L~~.~~T. ~~1945.~~ Determination of Total Organic and Available Form of Phosphorus in Soil. ***Soil Sci***., 1945; 59: 34-35.

Corazon M~~.~~ Ramirez. ~~2012.~~ Relationship between soil phosphorus availability and azolla growth, ***Soil Sci. Plant Nut.***~~,~~ 2012; **30** (4) : 595-598.

Debjani Halder, ~~and~~ Shyamal Kheroar. ~~2013.~~Mineralization and Availability of *Azolla*and Cyanobacteria Biomass Nutrients in Rice Soil***.J. Agrl. Sci. and Tech***. 2013; **3**: 782-789.

De Datta~~,~~S~~.~~K,~~ and~~ Buresh~~,~~ R~~.~~J. ~~1989.~~ Integrated nitrogen management in irrigated rice, ***Adv. Soil Sci***.~~,~~ 1989; **10** : 143-169.

Fogg~~,~~ G~~.~~E~~,~~. Steward~~,~~ W~~.~~D~~.~~P~~.~~, Fay~~.~~P~~.~~ ~~and~~ Walsky~~,~~ A~~.~~E. ~~1973.~~ The Blue Green Algae, Academic Press, New York~~.~~;1973

Comment [EA10]: 5. Conclusion part should be written as the conclusion in the abstract part

Comment [EA11]: All references should bewritten according to the referencing style of ASRJ as outlined in the Authors Guide.E.g.: *Hilly M, Adams ML, Nelson SC. A study of digit fusion in the mouse embryo. Clin Exp Allergy. 2002;32(4):489-98*

Comment [EA12]: Reference cannot be CROSS check

Comment [EA13]: Reference cannot be CROSS check

Comment [EA14]: Reference cannot be CROSS check

Halder~~,~~;D, ~~and~~ ; Kheroar~~,~~S. ~~2013.~~ Mineralization and Availability of *Azolla*and Cyanobacteria Biomass Nutrients in Rice Soil.***J.Agrl. Sci. Tech***.~~,~~2013; **3** : 782-789

IRRI, IRRISTAT, Biometric unit, IRRI, Los Banos, Philippines;1993

Kannaiyan S,~~.~~ Subramani S. ~~1992.~~ Use of Azolla as biofertilizer for rice crop, cyanobacterial nitrogen fixation, (**Ed**: Kaushik B. D.), ~~pp.~~ 1992; 281-289, Indian Agricultural Research Institute,New Delhi

Kumar~~,~~ B ~~and~~ Shahi~~,~~D~~.~~K.. ~~2016.e~~Effect of azolla as green manure on soil properties and grain yield of rice in acid soil of jharkhand. ***Annals plant soil Res***., 2016; **18** (3): 214-218.

Michihiko~~,~~ Y~~.~~, Naoki~~,~~T~~.~~, Noriyo, H~~.~~D~~.~~A ~~and~~ Katsunori~~,~~N~~. 2012.~~ Nitrogen fixation in azolla-anabaena symbiosis as affected by mineral nutrient status. ***Soil Sci. Plant Nut***~~.,~~ 2012; **26** (3) : 415-426.

Muruganayaki~~,~~ S ~~and A.~~ Jeyachitra A. ~~(2017)~~ Nitrogen assimilating enzyme of azolla as influenced by potassic fertilizer, ***Inter. J, Curr. Adv. Res***.~~,~~2017; **6**(~~0~~3), pp. 3012-3016.

Roy~~,~~ B. ~~1981.~~ Manuring of rice with Azolla, ***Oryza***, 1981; **21** : 238-241.

Sahrawat~~,~~K~~.~~L. ~~2000.~~ Macro and micronutrients removed by upland and lowland rice cultivars in West Africa, ***Comm. Soil Sci. Plant Analysis***, 2000; **31**:717-723.

Singh~~.~~ A~~.~~ L,~~.~~ ~~and P. K.~~ Singh PK.~~1987.~~ "Influence of Azolla management on the growth, yield of rice and soil fertility": II. N and P contents of plant and soil. ***Plant Soil***. 1987; **102** (1) : 49-54.

Singh P~~.~~K.~~1978.~~ "Nitrogen Economy of Rice Soils in Relation to Nitrogen Fixation by Blue-Green Algae and Azolla. In Increasing Rice Yield in Kharif" Central Rice Research Institute: Cuttack, India. 1987; 221–239.

Singh~~,~~ P~~.~~K,~~.~~ ~~and~~ Subudhi~~,~~ B~~.~~P. ~~1981.~~ Utilize Azolla in poultry feed, in: Subjected Matter Training-Cum-Discussion Seminar on Use of Bio-fertilizer with Special Reference to Azolla, Held from 16th to 21st Feb., CRRI, Orissa, 1981.

Stanford~~,~~ S. ~~and~~ English. ~~1949.~~ Use of flame photometer in rapid soils tests for Potassium and calcium. ***Agron. J***~~.,~~, 1949; **41:** 446-447.

Subba Rao~~,~~ A, ~~and~~ Sammi Reddy~~,~~ K. ~~2005.~~ Integrated nutrient management vis-à-vis crop production/productivity, nutrient balance, farmer livelihood and environment: India. Proc. Regional Workshop Beijing, China, ~~12-16 December~~ 2005.

Subbiah~~,~~ B~~.~~V, ~~and G. L.~~ Asija. ~~1956.~~ A rapid procedure for estimation of available nitrogen in soils. ***Curr. Sci.***, 1956; **25:** 259-260.

Subedi P, ~~and J.~~ Shrestha J. ~~2015.~~ "Improving soil fertility through Azolla application in low land rice": A review. ***Azarian J. Agri.***, 2015; **2** (2): 35-39.

Sutanto~~,~~ R. ~~2002.~~ Penerapanpertanianorganik. PenerbitKanisius. Yogyakarta 2002.

Comment [EA15]: Kannaiyan S, SubramaniS. Use of Azolla as biofertilizer for rice crop, Cyanobacterial Nitrogen Fixation. Ed: KaushikB D, Indian Agricultural Research Institute, New Delhi. 1992, pp. 281–289

Comment [EA16]: Reference cannot be CROSS check

Comment [EA17]: References cannot be CROSS check

Formatted: Font: Not Bold

Comment [EA18]: References cannot be CROSS check

Formatted: Font: Not Bold

Comment [EA19]: References cannot be CROSS check

Syamsiyah, J.,Sunarminto,B.H.. and Mujiyo. 2015. Changes in soil chemical properties of organic paddy field with azolla application.***J. Soil Sci.Agroclim***.,2015;**13** (1): , 2016, 68-73

Thanikachalam A.,.Rajakannu K., , Kannaiyan S. (1984) Effect of Neem cake, Carbofuran and Azolla application on phosphatase activity in soil. 25th Annual Conference of Association of Microbiologists of India, GB Pant University ofAgriculture and Technology, Pant Nagar, India. 1984.

Ventura,W., Watanabe, I, and Mascariña, G.B. 2012. Mineralization of Azolla N and its availability to wetland rice. ***Soil Sci. Plant Nutr***., 2012; **38**:(3),: 505-516

Valiente, E.F, .Ucha, A., Quesada, A., Leganes, F., and Carreres, R. 1998. Contribution of N2-fixing cyanobacteria to rice production, 5th International Symposium, Inorganic N Assimilation, Luso, Portugal, ***Plant Soil,*** 1998; **1** :107-112.

Walkley, A, and C.A. Black CA. 1934. An estimation of methods for determining organic carbon and nitrogen in the soils. ***J. Agric. Sci.,*** 1934; **25:** 598-609.

Watanabe I., Benjamin Padre, Jr., and Corazon Ramirez. 1991. "Mineralization of Azolla N and Its Availability to Wetland Rice". ***Soil Sci. Plant Nutr.,*** 1991; **37** (4): 679-688.

Yadav R. K., Abraham G., Singh Y. V., and P. K. Singh PK. 2014. "Advancement in the utilization of Azolla-Anabaena system in relation to sustainable agricultural practices". Proc. Indian National Science Academy, 2014; **80**:(2): 301-316.

Plate 1. View of Incubation Experiment

Comment [EA20]: Need to cross check year of publication

Comment [EA21]: References cannot be CROSS check

Comment [EA22]: References cannot be CROSS check

Formatted: Left, Indent: Left: 0", First line: 0", Line spacing: Multiple 1.15 li

Formatted: Centered, Space After: 0 pt, Line spacing: single

Formatted Table

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font color: Background 1

Formatted: Left, Indent: Left: 0", First line: 0.5", Line spacing: Multiple 1.15 li