

Original Research Article

EVALUATION OF THE CARDIOPROTECTIVE EFFECT OF *CITRULLUS LANATUS*(*C. LANATUS*) SEEDS IN STREPTOZOTOCIN INDUCED DIABETIC ALBINO RATS.

ABSTRACT

This pharmacological study was carried out to examine the potential of the ethanolic extract of *C. lanatus* in ameliorating cardiovascular dysfunctions in streptozotocin induced diabetic albino rats. The rats were induced with streptozotocin (50mg/kg) and those rats confirmed to be diabetic, were divided into five groups- Normal control, Diabetic control, Diabetic rats treated with *C. lanatus*(100mg/kg, 400mg/kg, 800mg/kg). Analyses were carried out in two phases –At the end of two weeks(14days) and at the end of four weeks(28 days). Biochemical analysis was performed on the blood and heart tissues. At the end of two weeks and four weeks. The data obtained were compared using one way analysis of variance (ANOVA) and the difference between means was obtained using Tukey's multiple comparison test. Analysis showed that cardiac enzymes: creatine kinase (CK-MB) and lactate dehydrogenase (LDH) which are markers of cardiac injury were significantly decreased ($p < 0.05$) which were prior to treatment raised. These changes were observed in a dose dependent manner. *Citrullus lanatus* presents significant potentials of maintaining structural integrity of the heart.

Keyword: *C.lanatus*, *Citrullus lanatus*, Diabetes, Streptozotocin, Cardioprotective, Cardiac enzymes

1. INTRODUCTION

Diabetes is a group of metabolic diseases which is due to a defect or a malfunction in insulin production or insulin action and is characterized by hyperglycaemia (High glucose level in blood). It results from impaired function in carbohydrate, protein and lipid metabolism leading to long term health complications (5)[1]. Cardiovascular complication has been highlighted as one of such. The incidence of myocardial dysfunction is higher in diabetic patients as compared to non-diabetic patients (18)[2] and the mechanism underlying the development of cardiovascular dysfunction seems to look quite complex and uncertain though various hypothesis including hypoinsulinemia, deregulated carbohydrate and lipid metabolism, formation of advanced glycation end products and oxidative stress has been suggested to explain the relationship between diabetes and cardiovascular disease. Hyperglycemia which is the metabolic hallmark of diabetes inflicts cellular injury with oxidative stress playing a supportive role (19)[3]. Antioxidants acts as defense to minimize the destructive effects of the free radicals and are grouped into non enzymatic and enzymatic antioxidants. The non enzymatic antioxidants include glutathione, lycopene, vitamins like vitamins A and C, while the enzymatic include catalase, peroxidase, and glutathione S transferase e.t.c. *Citrullus lanatus* which is grouped amongst the curcubitaceae family is rich in flavonoids, alkaloids, tannins, phenols, glycosides and these phytochemical are very useful to human health.(3)[4]. *C.lanatus seed* is said to possess antihyperlipidemic, antihyperglycemic.(12)[5], antiulcerogenic,(11)[6], antioxidative and hepatoprotective properties (7)[7]

2. MATERIALS AND METHODS

2.1 Plant materials and extraction

Fresh watermelon fruits were purchased from a watermelon depot at oil-mill market, Rivers state. They were halved and the seeds collected, washed, air dried for a week and then ground into fine powder. 250g of this powdered material was macerated in 1 litres of ethanol (250g/l) for 48 hours and then filtered using a muslin cloth. The filtrate was subjected to extraction by the use of soxhlet extractor and later concentrated using the rotary evaporator.

2.2 Experimental design.

Sixty albino rats (180-220g) were purchased and allowed to acclimatize for a period of two weeks during which they were fed *Ad libitum* on commercial dry pellet feed (top feed) and had free access to water. All animals were obtained from the Animal house, Department of Pharmacology, University of Port Harcourt and were housed in cages with saw dust bedding.

The animals had an overnight fast (about 12-14 hours) after which diabetes was induced by injecting intraperitoneally, a single dose of streptozotocin (STZ) (50mg/kg) dissolved in ice cold sodium citrate buffer, (0.1M, pH 4.5). Streptozotocin is an anti microbial agent with selective cytotoxicity to pancreatic β -cell that has the ability to induce a specific necrosis causing destruction of pancreatic β -cells. After 48 hours of induction, Accu-check Glucometer was used to determine the fasting blood glucose level. Rats with glucose concentration $\geq 250\text{mg/dl}$ were selected for the experiment. The rats were randomly divided into five groups of 12. Group A (Normal control) - consisted of non diabetic rats, Group B (Diabetic control) - consisted of STZ induced diabetic rats, Group C - consisted of diabetic rats treated with 100mg/kg of *C.lanatus* seeds extract, Group D - consisted of diabetic rats treated with 400mg/kg of *C.lanatus* seeds extract, and Group E - consisted of diabetic rats treated with 800mg/kg of *C.lanatus* seeds extract.

Treatment commenced on the third day after diabetes induction. A syringe with a specially designed metal ball-ended needle was used to feed the rats by oral gavage. At the end of first 14 days from the onset of treatment with *C. lanatus* seeds extract, six of the rats were sacrificed under Diethyl ether anaesthesia. This was on the 15th day after an overnight fast. Both blood and heart were collected. The remaining six continued in their treatment regime for the next 14 days. At the end of the period (that is on the 29th day), the animals were sacrificed under Diethyl ether anaesthesia after an overnight fast and the blood. Blood samples used for biochemical analysis were collected by cardiac thoracic puncture of the animals. About 5ml of the blood was collected into lithium heparin (anticoagulant) bottles and the plasma was used for assay of cardiac enzymes (Creatine kinase and Lactate dehydrogenase) and glucose was separated by centrifuging the whole blood at 5000rpm for about 10 minutes in an automated bench top centrifuge (Hettich Universal11). The plasma obtained was dispensed into plain bottles which were properly labeled and stored at -20°C until ready for use.

2.4 Animal Ethical Compliance

During the experimental period, there was strict adherence to ethical regulations required for handling experimental animal in accordance with National and institutional guidelines for protection of Animal welfare (15) [8]

3.0 METHODOLOGY

Determination of Creatine Kinase (CK-MB) activity by Kinetic Method

Procedure

1ml of working reagent was added to 40µl of sample mixed properly and incubated at 37⁰c for 100seconds. The changes in absorbance per minute ($\Delta OD/min$) for 3 mins was measured.

Calculation

Creatine kinase activity (U/L) = $\Delta OD/min \times 4127$

Lactate dehydrogenase determination by kinetic method

Procedure

10µl of plasma was added to 1000µl of the working reagent and then homogenized and incubated at 37⁰c for 1 min. The changes in absorbance per minute ($\Delta OD/min$) during a 3 minute period was measured.

Calculation

LDH-P activity (U/L) = $\Delta OD/min \times 1603$

3.1 Statistical analysis

Values obtained are presented as mean \pm standard error of mean (SEM). The statistical tool employed for analysis is one way analysis of variance (ANOVA) followed by Tukey's multiple comparison test using the Graph pad Prism Version 7.0 Software, developed by Graph Pad software, Inc, California. Values of $p < 0.05$ was considered as statistically significant

4.0 RESULTS

The results obtained from this experiment for a period of 14 and 28 days are listed in the tables below.

Table 4.1: THE EFFECT OF C .lanatus SEEDS EXTRACT ON CARDIAC ENZYMES ACTIVITY AFTER 14 DAYS ADMINISTRATION

Groups/ Parameters	CK-MB (U/L)	LDH (U/L)
Group A(NC)	9.500 \pm 0.82	53.60 \pm 2.66
Group B (DC)	21.14 \pm 1.02	68.40 \pm 2.14
Group C (100mg/kg)	11.11 \pm 3.27* ^{ce}	64.00 \pm 6.83
Group D (400mg/kg)	8.44 \pm 0.71	59.40 \pm 4.53* ^{bg}
Group E(800mg/kg)	7.34 \pm 1.41* ^{df}	57.20 \pm 3.42* ^{efh}
P-value	0.001	0.02

F – value

17.72

6.628

(*) –significant. **Legends showing the variation in groups using Tukey test are:**

a= Group A versus Group D, **b** = Group B versus Group D, **c**= Group A versus Group C, **d**= Group A versus Group E, **e**= Group B versus Group C, **f** = Group B versus E, **g** = Group C versus Group D and **h** = Group D versus Group E, **I** = Group A versus Group B, **J**= Group C versus Group E

Table 4.1 showed the effect of *C. lanatus* seeds extract on CK-MB and LDH levels in the rats after 14 days. The results were expressed as mean± SEM. The data was analyzed using ANOVA then Tukey's comparison test. Significance between the means ($p < 0.05$) were observed as indicated in some groups.

Table 4.2: THE EFFECT OF *C. lanatus* SEEDS EXTRACT ON CARDIAC ENZYMES ACTIVITY AFTER 28 DAYS OF ADMINISTRATION.

Groups/ Parameters	CK-MB (U/L)	LDH (U/L)
Group A(NC)	5.34±0.41	46.80 ±3.28
Group B (DC)	13.18±7.96* ^l	67.20± 4.70
Group C (100mg/kg)	9.54±1.76	35.80 ± 5.00
Group D (400mg/kg)	6.68 ±0.30	24.60 ± 1.63* ^b
Group E (800mg/kg)	6.06 ±0.62	22.60 ±2.99* ^d
P-Value	0.007	0.0005
F - Value	4.708	8.100

(*) -significant

Table 4.2 showed the effect of *C. lanatus* seeds extract on CK-MB and LDH levels in the rats after 28 days. The results were expressed as mean± SEM. The data was analyzed using ANOVA then Tukey's comparison test. Significance between the means ($p < 0.05$) were observed as indicated in some groups.

DISCUSSION.

The increase in serum activities of Creatine kinase and lactate dehydrogenase in experimental diabetic rats may indicate compromise on the cardiomyocytes integrity which agrees with reports that serum Creatine kinase-MB (CK-MB) and lactate dehydrogenase (LDH) activities are associated with a rise in cardiac muscular damage (14)[9]. In this study there was a decrease in lactate dehydrogenase (LDH) and CK-MB activities on treatment with *C. lanatus* seed extract which would indicate that intake of these seed was able to improve on cardiac cells integrity and may have good potentials in reducing cardiovascular risk in diabetes mellitus.

CONCLUSION

The wholesome and regular consumption of *C. lanatus* fruit (rind, pulp and seeds) should therefore be encouraged amongst diabetic and non diabetic individuals as often times the seeds and rind are discarded.

REFERENCES

1. Ansley, D. M., & Wang, B. (2013). Oxidative stress and myocardial injury in the diabetic heart. *The Journal of pathology*, 229(2), 232-241.
2. Clairborne A. (1995). Catalase activity. In Handbook of Methods for Oxygen Radical Research, Greewald AR (ed.). CRC Press: Florida, 237–242.
3. Deshmukh, C. D., & Jain, A. (2015). Hypoglycemic effect of methanolic extract of citrullus lanatus seeds. *International journal of pharmaceutical, Chemical & Biological Sciences*, 5(4).
4. Etuk, E. U. (2010). Animals models for studying diabetes mellitus. *Agric Biol JN Am*, 1(2), 130-134.
5. Gezginici-Oktayoglu, S., Basaraner, H., Yanardag, R., & Bolkent, S. (2009). The effects of combined treatment of antioxidants on the liver injury in STZ diabetic rats. *Digestive diseases and sciences*, 54(3), 538-546.
6. Ilchukwu, C. C., Ebenebe, U. E., Ubajaka, C. F., Ilika, A. L., Emelumadu, O. F., & Nwabueze, S. A. (2014). The Role of Oxidative Stress in Diabetes Mellitus: A 24-year Review. *Afrimed J ournal*, 5(1), 1-7.
7. Madhavi P, Kamala V., & Habibur R.,(2012).Hepatoprotective Activity of Citrullus Lanatus Seed Oil on CCl4 Induced Liver Damage in Rats.*Scholars Academic Journal of Pharmacy*,1 (1):30-33.
8. Misra, H. P., & Fridovich, I. (1972). The role of superoxide anion in the autoxidation of epinephrine and a simple assay for superoxide dismutase. *Journal of Biological chemistry*, 247(10), 3170- 3175.
6. Nakhaee A.,Bokaeian M.,Saravani.,Farhangi A.,&Akbarzadeh A,(2009),Attenuation of oxidative stress in streptozotocin induced diabetic rats by Eucalyptus globulus.*Indian J.clinical biochemistry*, 24(4),419-425
7. Ohkawa, H., Ohishi, N., & Yagi, K. (1979). Assay for lipid peroxides in animal tissues by thiobarbituric acid reaction. *Analytical biochemistry*, 95(2), 351-358.
8. Okunrobo O. Lucky, Uwaya O. John, Imafidon E.Kate, Osarumwense O. Peter, Omorodion E. Jude,(2012)Quantitative determination, Metal analysis andAntiulcer evaluation of Methanol seeds extract of Citrullus lanatus Thunb (Cucurbitaceae) in Rats.*Asian Pacific Journal of Tropical Disease*. S1:261-S1265
9. Omigie, I. O., & Agoreyo, F. O. (2014). Effects of watermelon (Citrullus lanatus) seed on blood glucose and electrolyte parameters in diabetic wistar rats. *Journal of Applied Sciences and Environmental Management*, 18(2), 231-233.
10. Oseni, O. A., & Okoye, V. I. (2013). Studies of Phytochemical and Antioxidant properties of the fruit of watermelon (Citrullus lanatus).(Thunb.). *J. Pharm. Biomed. Sci*, 27(27), 508-514.

11. Patel, S. S., & Goyal, R. K. (2011). Cardioprotective effects of gallic acid in diabetes-induced myocardial dysfunction in rats. *Pharmacognosy research*, 3(4), 239.
12. PHS [Public Health Service]. (1996). Public health service policy on humane care and use of laboratory animals
13. Sani, U. M. (2015). Phytochemical screening and antidiabetic effect of extracts of the seeds of *Citrullus lanatus* in alloxan-induced diabetic albino mice. *Journal of Applied Pharmaceutical Science*, 5(3), 51-54.
14. Sedlak, J., & Lindsay, R. H. (1968). Estimation of total, protein-bound, and nonprotein sulfhydryl groups in tissue with Ellman's reagent. *Analytical biochemistry*, 25, 192-205.
15. Thom, T., Haase, N., Rosamond, W., Howard, V. J., Rumsfeld, J., Manolio, T., ... & Lloyd-Jones, D. (2006). Heart disease and stroke statistics--2006 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation*, 113(6), e85
16. Tiwari, B. K., Pandey, K. B., Abidi, A. B., & Rizvi, S. I. (2013). Markers of oxidative stress during diabetes mellitus. *Journal of biomarkers*, 2013.
17. Wei Bhaar D.(1975). *Med Weit*,26: 387.