

2
3 **Occurrence of Bluntnose Sixgill Shark**
4 ***Hexanchus griseus* (Bonnaterre,**
5 **1788) in Yeşilovacık Bay,**
6 **Northeastern Mediterranean**

7
8
9
10 **ABSTRACT**
11

Aims: In this study, an individual of *H. griseus* was caught from the Mersin Bay and this ichthyologic note presents a new female record of *H. griseus* for Mersin Bay, Mediterranean coast of Turkey. This study aims to contribute to the chronological records of the species in the Mediterranean Sea.

Place and Duration of Study: Yeşilovacık bay is a small bay in the west of Mersin Bay. Yeşilovacık bay which lies at approximately 36°07'n latitude, 33°37'e, longitude and about 143 km western of in Mersin province, Turkey. Sample: On 19 March 2018 blunthose sixgill shark specimen was measured and then deposited in the Museum of the Systematic, Faculty of Fisheries, University of Mersin.

Methodology: A female specimen of *H. griseus* with a total length (TL) of 350 cm (400 kg) was captured by a commercial trawl at a depth around of 280 m off the Yeşilovacık Bay (Northeastern Mediterranean Sea), Turkey. Total length was measured to the nearest 1 mm and the weight of specimen was determined to the nearest kilogram.

Results: Measurements of the specimen are presented and compared with the previous records of *H. griseus* in the Eastern Mediterranean coast of Turkey (Iskenderun Bay), which are given in Table. The identification was carried out according to previous report.

Conclusion: To date specific conservation measures are not known for this species for the Northeastern Mediterranean coast of Turkey. Thus, the effect of coastal fishery on sharks should be regularly monitored and protection measures should be taken for this region in order for this shark species to continue their conservation for the next generation.

12
13 *Keywords: Hexanchidae; Shark; Mersin Coast; Mediterranean Sea; Turkey*
14

15
16 **1. INTRODUCTION**
17

18 The bluntnose sixgill shark, *Hexanchus griseus* (Bonnaterre, 1788), apex predator of the
19 deep waters, located at the upper trophic level of the food chain, belongs to the family
20 Hexanchidae [1]. It is known that the species inhabits depths of 2500 m in the subtropical
21 regions of the oceans and seas. *H. griseus* usually lives at depths between 180-1100 m
22 [2,3]. It is a deep-water species, which lives in the outer continental and insular shelves and
23 upper slopes. They are found in pelagic areas and upper slopes [4]. Adult specimens are
24 usually below 91 m, while juveniles can be found near the coast [3]. They are found on the
25 bottom during the day. However, they are close to the surface at night to feed. They usually
26 feed on sharks, rays, chimaeras, bony fish, squids, crabs, shrimps, carrion, and even seals
27 [5]. *H. griseus* is an oceanodrom fish species [6]. This circumglobal species is distributed in
28 tropical and temperate waters, including Eastern Atlantic, Western Atlantic, Western Pacific

29 and Eastern Pacific [7]. The maximum weight reported in the literature is 590 kg [8].
30 Although maximum total length (TL) reported in the literature is 482 cm according to
31 Compagno [9] and Cervigón et al. [10]. However, the maximum total length of the species
32 reported in the records is 650 cm in the Turkish coast by Kabasakal [11]. Compagno [9]
33 reported individuals of this species reach reproduction maturity in the range of 300-482 cm
34 in TL. Although studies on the biological properties of the species are limited, it has been
35 reported that they are ovoviviparous and their fecundity vary between 22 to 108 [1].

36 The bluntnose sixgill shark, *H. griseus* has been reported from France [12,13], from Italy
37 [14,15], from Israel [16], from Tunisia [17], from Greece [18] in the Mediterranean Sea.
38 There are reports of *H. griseus* from the Pacific [1,19], from the Indian and Atlantic Oceans
39 [20]. It has been also reported from Turkey waters of the Mediterranean, the Aegean Sea,
40 the Sea of Marmara and Black Sea [21]. According to Kabasakal [11]; first record of this
41 species in the waters of Turkey was reported in 1923 by [22], and the following studies:
42 Devedjian [23]; Konsuloff and Drensky [24]; Ben-Tuvia [16]; Akşiray [25]; Meriç [26]; Goren
43 and Galil [27]; Jones et al. [28]; Kabasakal [21]; Sion et al. [29]; Golani et al. [30]; Kabasakal
44 [31]; Kabasakal [11]; Basusta and Basusta [32].

45 In this study, an individual of *H. griseus* was caught from the Mersin Bay and this
46 ichthyologic note presents a new female record of *H. griseus* for Mersin Bay, Mediterranean
47 coast of Turkey. This study aims to contribute to the chronological records of the species in
48 the Mediterranean Sea.

49
50

51 **2. MATERIAL AND METHODS**

52

53 **2.1 Study Area**

54

55 Yeşilovacık Bay is a small bay in the west of Mersin Bay. (Fig. 1). Yeşilovacık Bay which lies
56 at approximately 36°07'N latitude, 33°37'E, longitude and about 143 km western of in Mersin
57 Province, Turkey. The bay is the spawning and breeding area for many bony fish species
58 due to the wide continental shelf. The biodiversity of the Yeşilovacık Bay is affected by the
59 migration of the alien species. Both sides of the bay are surrounded by littoral shores, while
60 sandy beach is located inside. It is covered with sandy muddy sediment on the sea floor of
61 the bay. The bay has an anticyclonic gyre opposite to the cyclonic gyre, which the main gyre
62 of the Mediterranean Sea. The maximum depth of the bay is 180 m. The physical and
63 chemical parameters of the Bay are similar to those of the Mediterranean Sea. Tourism,
64 agriculture, fishing and ship transportation are the main activities in the bay.

65

66 **2.2. Fish Sampled**

67

68 On 19 March 2018, a female specimen of *H. griseus* with a total length (TL) of 350 cm (400
69 kg) was captured by a commercial trawl at a depth around of 280 m off the Yeşilovacık Bay
70 (Northeastern Mediterranean Sea), Turkey (Fig. 1).

71 Total length was measured to the nearest 1 mm and the weight of specimen was determined
72 to the nearest kilogram. The head and fins of the specimen were deposited in the Museum
73 of the Systematic, Faculty of Fisheries, Mersin University, (MEUFC-18-11-058), (Fig. 2).

74 **3. RESULTS**

75

76 In this study, a female individual was captured from the Mediterranean waters off Turkish
77 coast (Mersin Bay). The identification was carried out according to Compagno [9]. Some
78 measurements of the specimen are presented and compared with the previous records of *H.*
79 *griseus* in the Eastern Mediterranean coast of Turkey (Iskenderun Bay), which are given in
80 Table 1.

81

82 The species possesses six pairs of gill slits, a subterminal mouth and a short, broadly
83 rounded blunt snout. It is grey with or without irregular brown spots. Also, the distance from
84 the dorsal fin to the caudal fin is equal to the length of the dorsal fin base [9].

85

86 The color of fresh specimen was dark grey without spots on the back and white on the belly,
87 with no clear boundary between the two colors.

88

89

90 **4. DISCUSSION**

91

92 The bluntnose sixgill shark, *H. griseus* is not a commercial species due to the fact that it is a
93 deep-water species and it is difficult to capture. However, it can also be sold for commercial
94 purposes when caught by commercial fishermen [1]. Although *H. griseus* is found in deep
95 waters (1-2500 m), it can come up to shallow waters (50-100 m) [11]. Individuals of this
96 species are usually caught during the fishing activities in coastal waters. In this study, the
97 reported individual was caught at a depth of 280 m.

98 The size at birth is 56 to 70 cm and the total length of the species is usually 300 cm [10].
99 Females mature at about 300 to 482 cm and males at about 300 to 350 cm [9,33,34].
100 Kabasakal [11] reported 6 female individuals of *H. griseus* with a total length of 350 cm were
101 caught between in October-April from Taşucu and Kaş. In the same period, the presence of
102 a large number of mature female individuals in the same region may indicate that they came
103 for breeding purposes. The total length of the individual caught in this study was 350 cm and
104 it was a mature individual. It was evaluated that this individual may have reached shallow
105 waters in the same period with the purpose of reproduction.

106 The bluntnose sixgill shark commonly feeds on *Squalus* spp., sharks, rays, bony fish, crabs,
107 shrimps, carrion, and seals [9,20,35]. In this study, a large number of bony fish species were
108 found in the stomach contents. In addition, shark (*Squalus* spp.), *Galeus melastomus* and
109 some species of rays (*Dipturus oxyrinchus*), as well as cans and plastic remains were found.
110 Similar Barrull and Mate [36] reported that *Scyliorhinus canicula*, *G. melastomus*, *Merluccius*
111 *merluccius* and *Phycis blennoides* were found in the stomach contents of six shark samples
112 caught in the Catalan Sea. Celona et al. [37] reported the stomach contents of 23 specimens
113 captured in Eastern Sicilian waters mainly consist of bony fish (60.87%), cephalopods
114 (13.04%), decapod crustaceans (8.7%), cartilaginous (4.35%) and echinoderms (4.35%).
115 According to Ebert [20] the main prey of individuals over 200 cm of *H. griseus* in the coasts
116 of South Africa is composed of bony fish. Consumption of sharks and sea mammals has
117 been reported to be associated with their lengths. In this study, the main prey of the
118 captured individual (350 cm, TL) formed bony fish. This finding is consistent with the
119 literature.

120 The bluntnose sixgill shark has several records in the Mediterranean Sea, Turkey [11]. 150
121 individuals of *H. griseus* were caught from the Turkish coastal waters between 1963-2013
122 years by commercial fishermen. 90 specimens from the Marmara Sea, 41 specimens from
123 the Turkish coastal waters of Aegean Sea, 15 specimens from the Turkish coastal waters of

124 Mediterranean Sea and 3 specimens from the Turkish coastal waters of Black Sea and 1
125 specimen from the Dardanelles were reported [11]. After 2013, 2 individuals have been
126 reported from Mediterranean coasts of Turkey [32,38].

127 To date there is little information available about habitat, ecology, and population of *H.*
128 *griseus*. This species is listed as Near Threatened (NT) in the Global Red List by the
129 International Union for Conservation of Nature, IUCN [39,40] and also by CITES [41].
130 Therefore, this record is significant and the species could be considered as exceptionally
131 rare in the Mediterranean, Turkey.

132 The bluntnose sixgill shark is a species that has not been economically important because it
133 is not a species offered for human consumption but marketed for human consumption when
134 it is caught to the nets by fishermen in some regions. The fishermen do not like this shark
135 species because they can damage the economically important seafood. The pressure of the
136 species on the economic species is neglected in the scientific researches. However,
137 fishermen should be encouraged to release the sharks caught by mistake. This is an
138 important move in terms of conservation of species and improvement of fisheries
139 management.

140

141 **5. CONCLUSION**

142 This predator species in the upper trophic levels of the food chain are found to be few in
143 number, their distribution in deep waters and the difficulties in catching them therefore limit
144 the biological studies related to the species. These individuals, which are accidentally caught
145 by the fishermen and come to the shallow waters for breeding purposes, enable the study of
146 the species for scientific purposes. This situation is not preferred for commercial fishermen
147 because they cannot market this species economically to the people of our country.
148 Therefore, fishermen can be more easily encouraged to release the captured *H. griseus* into
149 the water.

150 To date specific conservation measures are not known for this species for the Northeastern
151 Mediterranean coast of Turkey. Thus, the effect of coastal fishery on sharks should be
152 regularly monitored and protection measures should be taken for this region in order for this
153 shark species to continue their conservation for the next generation.

154

155 **COMPETING INTERESTS**

156

157 The authors declare no competing of interests.

158

159

160 **ETHICAL APPROVAL**

161

162 As per international standard or university standard written ethical approval has been
163 collected and preserved by the authors.

164

165 **REFERENCES**

166

167 1. Ebert DA. Biological aspects of the sixgill shark, *Hexanchus griseus*. Copeia. 1986;1:131-
168 135.

- 169 2. Castro JI, Woodley CM, Brudek RL. A preliminary evaluation of the status of shark
170 species. FAO Fish. Tech. P. 380, Rome: FAO; 1999.
- 171 3. Mundy BC. Checklist of the fishes of the Hawaiian Archipelago. Bishop Mus. Bull. Zool.
172 2005;(6):1-704.
- 173 4. Last PR, Stevens, J.D. 1994. Sharks and rays of Australia. Australia: CSIRO; 1994.
- 174 5. Hart JL. Pacific fishes of Canada. Fish. Res. Bd. Canad. Bull. 1973;180:1-740.
- 175 6. Riede K, Global register of migratory species - from global to regional scales. Final Report
176 of the R&D-Projekt 808 05 081. Germany: Federal Agency for Nature Conservation, Bonn;
177 2004.
- 178 7. Compagno LJV, Niem VH. Hexanchidae. Cowsharks, sixgill, and sevengill sharks. In:
179 Carpenter KE, Niem VH, editors, FAO identification guide for fishery purposes. The Living
180 Marine Resources of the Western Central Pacific. Rome: FAO; 1998.
- 181 8. Lamb A, Edgell P. Coastal fishes of the Pacific Northwest. Madeira Park, (BC, Canada):
182 Harbour Publishing Co. Ltd; 1986.
- 183 9. Compagno LJV. FAO Species Catalogue. Vol. 4. Sharks of the world. An annotated and
184 illustrated catalogue of shark species known to date. Part 1 - Hexanchiformes to
185 Lamniformes. FAO Fish. Synop. Rome: FAO. 1984;125(4/1):1-249.
- 186 10. Cervigón F, Cipriani R, Fischer W, Garibaldi L, Hendrickx M, Lemus AJ, Márquez R,
187 Poutiers JM, Robaina G, Rodriguez B. Fichas FAO de identificación de especies para los
188 fines de la pesca. Guía de campo de las especies comerciales marinas y de aquassalobres
189 de la costa septentrional de Sur América. Preparado con el financiamiento de la Comisión de
190 Comunidades Europeas y de NORAD. Rome: FAO; 1992. Spain.
- 191 11. Kabasakal H. Bluntnose Sixgill Shark, *Hexanchus griseus* (Chondrichthyes:
192 Hexanchidae), caught by commercial fishing vessels in the seas of Turkey between 1967
193 and 2013. Annales, Ser. Hist. Nat. 2013;23(1):33-48.
- 194 12. Risso A. Ichthyologie de Nice ou Histoire Naturelle des Poissons. F. Schoell, Paris;
195 1810.
- 196 13. Moreau E. 1881. Histoire Naturelle des Poissons de la France. Vol. II. Paris: Masson;
197 1881.
- 198 14. Tortonese E. Fauna d'Italia. Leptocardia, Ciclostomata, Selachii. Bologna: Calderini;
199 1956.
- 200 15. Bini D. Atlantedei Pesci delle Coste Italiane. Vol. I. Leptocardi-Ciclostomi-Selachi.
201 Mondo Sommerso, Roma; 1967. Italian.
- 202 16. Ben-Tuvia A. Revised list of the Mediterranean fishes of Israel. Isr. J. Zool. 1971;20:1-
203 39.
- 204 17. Quignard JP, Capapé C. Complement à la listecommentée des selaciens de Tunisie.
205 Bull. Inst. Océanogr. Pêche, Salammbô. 1972;2:445-447.
- 206 18. Papaconstantinou C. Fauna Graeciae IV. Check-list of Marine Fishes of Greece.
207 Athens: Hellenic Zoological Society; 1988.
- 208 19. Dunbrack R, Zeilinski R. Seasonal and diurnal activity of sixgill sharks (*Hexanchus*
209 *griseus*) on a shallow water reef in the Strait of Georgia, British Columbia. Can. J. Zool.
210 2003;81(6):1107-1111.
- 211 20. Ebert DA. Diet of the sixgill shark *Hexanchus griseus* off southern Africa. S. Afr J. Mar.
212 Sci. 1994;14:213-218.
- 213 21. Kabasakal H. Preliminary observations on the reproductive biology and diet of the
214 bluntnose sixgill shark, *Hexanchus griseus* (Bonnaterre, 1788) (Chondrichthyes:
215 Hexanchidae), in Turkish Seas. Acta Adriat. 2004;45:187-196.
- 216 22. Ninni E. Primo contributo allo studio dei pesci della pescanella acquadell'impero
217 Ottomano. Missione Italiana per l'Esplorazione dei Mari di Levante: Premiate Officine
218 Grafiche Carlo Ferrari; 1923.
- 219 23. Devedjian K. Pêche et Pêcheries en Turquie. Istanbul: Imprimerie de l'Administration de
220 la Dette Publique Ottomane; 1926.

221 24. Konsuloff S, Drensky P. Die fischfauna der Aegäis. Ann. Univ. Sofia Fac. Sci. Nat.
222 1943;39:293-308. Germany.

223 25. Akşiray F. Türkiye deniz balıkları ve tayin anahtarı. 2nd ed. Publications of Istanbul
224 University, No. 3490, Istanbul; 1987. Turkish.

225 26. Meriç N. 1995. A study on existence of some fishes on the continental slope of the Sea
226 of Marmara. Turk. J. Zool. 1995;19:191-198.

227 27. Goren M, Galil BS. Records of *Cataetys laticeps* and *Ophidion barbatum* (Ophidiiformes)
228 in the eastern Mediterranean, with comments on the deep sea ichthyofauna. Cybium.
229 2002;26:150-152.

230 28. Jones EG, Tselepidis A, Bagley PM, Collins MA, Priedel G. Bathymetric distribution of
231 some benthic and benthopelagic species attracted to baited cameras and traps in the deep
232 eastern Mediterranean. Mar. Ecol. Prog. Ser. 2003;251:75-86.

233 29. Sion L, Bozzano A, D'Onghia G, Capezzuto F, Panza M. Chondrichthyes species in
234 deep waters of the Mediterranean Sea. Sci. Mar., 2004;68(Suppl. 3):153-162.

235 30. Golani D, Öztürk B, Başusta N. Fishes of the eastern Mediterranean. Turkish Marine
236 Research Foundation, Publication no. 24. Turkey, Istanbul; 2006.

237 31. Kabasakal H. Türk sularında köpekbalıkları (Sharks of Turkish Waters). Istanbul: Deniz
238 Yayınları; 2011. Turkish.

239 32. Başusta N, Başusta A. Additional record of the bluntnose six-gill shark, *Hexanchus*
240 *griseus* (Bonnaterre, 1788) from Iskenderun Bay with its morphometric measurements. J.
241 Black Sea/Med. Environ. 2015;21(2):224-226.

242 33. Branstetter S, Mceachran JD. A first record of *Odontaspis noronhai* (Lamniformes:
243 Odontaspidae) for the western North Atlantic, with notes on two uncommon sharks from
244 the Gulf of Mexico. Northeast Gulf Sci. 1986;8(2):153-160.

245 34. Capapé C, Hemida F, Guélorget O, Barrull J, Mate I, Ben Souissi J, Bradaï MN.
246 Reproductive biology of the bluntnose sixgill shark *Hexanchus griseus* (Bonnaterre, 1788)
247 (Chondrichthyes: Hexanchidae) from the Mediterranean Sea: a review. Acta Adriat.,
248 2004;45:95-106.

249 35. Boseman, M. Hexanchidae. In: Whitehead PJP, Bauchot ML, Hureau JC, Nielsen J,
250 Tortonese E, editors. Fishes of the North-eastern Atlantic and the Mediterranean. Vol. 2.
251 Paris: UNESCO; 1984.

252 36. Barrull J, Mate I. 2000. Biología de la cañabota, *Hexanchus griseus* (Bonnaterre, 1788)
253 en el Mar Mediterraneo. Bol. Asoc. Esp. Elasm. 2000;3:13-20. Spanish.

254 37. Celona A, De Maddalena A, Romeo T. Bluntnose sixgill shark, *Hexanchus griseus*
255 (Bonnaterre, 1788), in the Eastern North Sicilian Waters. Bol. Mus. Storianat. Venezia.
256 2005;56:137-151.

257 38. Bayhan YK, Ergüden D, Cartes JE. Deep sea fisheries in Mersin Bay, Turkey, Eastern
258 Mediterranean: Diversity and abundance of shrimps and benthic fish fauna. Acta Zool. Bulg.
259 2018;70(2):259-268.

260 39. Cook SF, Compagno LVJ. *Hexanchus griseus*. The IUCN Red List of threatened
261 species, 2009. Version 2018-2. Accessed 20 February 2019.
262 Available: <https://www.iucnredlist.org/species/10030/3155348>.

263 40. IUCN. IUCN Red List of Threatened Species (Version 2009-2). 2009. Accessed
264 February 2019.
265 Available: <https://www.iucnredlist.org>.

266 41. CITES, UNEP-WCMC. The Checklist of CITES Species Website. Appendices I, II and III
267 valid from 04 April 2017. CITES Secretariat, Geneva, Switzerland. Compiled by UNEP-
268 WCMC, Cambridge, UK. 2017. Accessed 21 February 2019.
269 Available: www.cites.org/eng/app/appendices.php.

270

271


272

273

274 **Table 1. Comparison of measurements of the bluntnose sixgill shark specimen (*H.***
 275 ***griseus*) from Mersin Bay (Northeastern Mediterranean, Turkey) with previous record**
 276 **in the Iskenderun Bay (Eastern Mediterranean, Turkey)**
 277

	This study		Başusta and Başusta (2015)	
Specimen	n=1		n=1	
Sex	Female		Male	
Measurements	Length (cm)	Ratio (TL%)	Length (cm)	Ratio (TL%)
TL-Total Length)	350.0	100.00	213.0	100.00
FL-Fork Length	271.8	77.66	165.4	77.65
PCL-Precaudal Length	245.1	70.04	149.2	70.05
HDL-Head Length)	53.2	15.20	32.4	15.21
PNL-Prenarial Length	18.5	5.30	11.3	5.31
EYL -Eye Length	12.0	3.41	7.3	3.43
CDM-Dorsal Caudal Margin	101.8	29.09	62.0	29.11
DL-Dorsal Length	22.1	6.32	13.5	6.34
DH-Dorsal Height	23.0	6.55	14.0	6.57
CLI-Clasper Inner Length	-	-	7.3	3.43
BH-Body Height	37.7	10.68	23.0	10.80
Weight (kg)	400	-	480	-

278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300



301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320

Fig. 1. The shaded area indicates the location where the specimen of *H. griseus* was caught from Mersin Bay

UNDER PEE



321
322
323
324
325
326
327
328
329

Fig. 2. Female specimen of *H. griseus* caught from Mersin Bay

UNDER PEER REVIEW