

Comparative Study on the Degree of Randomness of the Random Number Tables

ABSTRACT

In the field of statistics as well as in the different branches of experimental sciences, random number tables have been playing a vital role for the purpose of selecting random samples. Among the existing different random number tables, four tables namely, Tippet's random number table, Fisher and Yates random number table, Kendall and Smith's random number table and random number table of RAND Corporation are of most frequent use. The current study aims at attempting to make a comparative review on the degree on randomness of these four most frequently used random number tables based on χ^2 test, run test and deviation test. From the findings based on χ^2 test, highest degree of randomness is observed in random number table due to RAND Corporation followed by due to Kendall and Smith, Tippet, Fisher and Yates, respectively. In case of run test, highest degree of randomness is noticed in random number table due to Fisher and Yates followed by due to Tippet, RAND Corporation, Kendall and Smith, respectively. However, from the findings based on the deviation test, it is clear that highest degree of randomness is observed in random number table due to Kendall and Smith followed by due to Fisher and Yates, RAND Corporation, Tippet respectively. It can be observed that the findings obtained in the studies based on different tests are not alike. Consequently, there is necessity to search for the reasons of the difference between these findings. Moreover, it can also be concluded that attempts should be made by the researchers to construct new random numbers table with enhanced degree of randomness than that of the existing tables.

Keywords: Fisher and Yates, Kendall and Smith, RAND Corporation, Randomness, Random number table, Tippet.

1. INTRODUCTION

Random number tables have been playing a vital role in statistics as well as in the different branches of experimental sciences for the purpose of selecting random samples. Use of these tables are much more effective than selecting the random samples manually with dice, cards etc. Several random number tables have already been constructed by the renowned researchers. Those contributions are mainly due to Tippet [1], Fisher and Yates [2], Kendall and Smith [3, 4], Mahalanobis [5], Quenouille [6], Rand Corporation [7], Snedecor and Cochran [8], Hald [9], Royo and Ferrer [10], Moses and Oakford [11], Rohlf and Sokal [12], Manfred [13], Rao, Mitra and Matthai [14] etc. Methods of drawing of random four-digit numbers, random five-digit numbers, random six-digit numbers and random seven-digit numbers from a combination of independent tables of random two-digit numbers and random three-digit numbers have also already been developed. [15, 16, 17, 18, 19, 20, 21, 22]. However, usage of computational random number generators have also been observed to be emerging. If carefully prepared, the process of filtering and testing can eliminate any

noticeable bias or asymmetry from the numbers such that the tables provide the most 'reliable' random numbers available to the casual user. Among these different random number tables, four tables namely, Tippet's random number table, Fisher and Yates random number table, Kendall and Smith's random number table and random number table of RAND Corporation are of most frequent use [23]. The current study aims at attempting to make a comparative review on the degree on randomness of these four most frequently used random number tables.

2. FREQUENTLY USED RANDOM NUMBER TABLES

2.1 Tippet's random number table

This table consists of 10,400 four-digit random numbers. Karl Pearson emphasized on testing statistical theories by sampling experiments. Tippet's random number could put to use for this purpose [24].

2.2 Fisher and Yates random number table

From the 10th to 19th digits of A.S. Thompson's 20-figure logarithmic tables, Fisher and Yates obtained the random numbers. In choosing from those digits, An element of randomness was introduced by using playing cards for the selection of half pages of the tables and of a column between 10th to 19th and finally for allotting these digits to the 50th place in a block [25].

2.3 Kendall and Smith's random number table

In the year of 1939, a set of 100,000 digits were published by M.G. Kendall and B. Babington Smith produced by a specialized machine in conjunction with a human operator [26].

2.4 Random number table of RAND Corporation

In the mid-1940s, development of a large table of random number table was set about by the RAND Corporation with the Monte Carlo method. With the help of a hardware random number generator, 'A Million Random Digits' with 100,000 Normal Deviates were produced. The RAND table used electronic simulation of a roulette wheel attached to a computer, the results emanated from which were then filtered and tested with substantial care before being used to generate the table [27].

3. TESTS USED FOR CHECKING RANDOMNESS

3.1 χ^2 Test

Pearson's chi-square test has been used in order to test whether the occurrences of the numbers appeared in the table is random or not. [28, 29, 30, 31]. This is equivalent to test that equal numbers of 0s, 1s, 2s, 3s, ... , 9s are present in the table or not.

Let N be the number of occurrences of the ten digits in the table and O_i = Observed frequency of the digit i, E_i = Expected frequency of the digit i ($i = 0, 1, 2, \dots, 9$) among the N occurrences. Then the χ^2 statistic for testing the null hypothesis, "the occurrences of the digits in the table is random" i.e. "each digit has the probability 0.1 to occur in any position", which is equivalent to testing "the discrepancy between the observed frequencies and the corresponding expected frequencies of the digits is insignificant" [23] is

$$\chi^2 = \sum_{i=1}^9 \frac{(O_i - E_i)^2}{E_i}, \text{ which follows } \chi^2 \text{ distribution with 9 degrees of freedom.}$$

This statistic can be employed to examine the randomness of the whole table as well as of any part of the table provided that the test satisfies the necessary assumptions of simple

random sample, sample size, expected cell count and independence [32, 33, 34]. The frequency test has been applied to each 100 occurrences.

3.2 Run Test

The run test is a non-parametric test to test the randomness for a two valued data sequence [35]. A run test is based on the null hypothesis that from the same distribution, each element in the sequence has been drawn independently.

Let us consider the following hypothesis:

H_0 : The occurrences of numbers in a table are in random manner.

H_1 : The occurrences of the numbers in the table are not in random manner.

Let, U = Number of observed runs yielded by n successive numbers in a table. Then, U follows a binomial distribution with expectation $E(U)$ and variance $V(U)$ given by $\frac{(n+2)}{2}$ and

$\frac{n(n-1)}{4(n-1)}$, respectively.

Then for large n , under H_0 , the test statistic $Z = \frac{U - E(U)}{\sqrt{V(U)}} \sim N(0, 1)$

One has to accept or reject the null hypothesis H_0 on comparing the values of $|Z|$ with the corresponding theoretical value of $|Z|$ namely 1.96 (at 5% level of significance) and 2.58 (at 1% level of significance).

3.3 Deviation Test

The statistic t can be considered as the ratio of the departure of the estimated value of a parameter from its hypothesized value to its standard error. The t -test is any statistical hypothesis test in which the test statistic follows a Student's t -distribution under the null hypothesis [36, 37].

Let $d_i = d_i(N)$ be the deviation of the observed number of occurrences of the digit i from its theoretical number of occurrences among N occurrences of the 10 digits ($i = 0, 1, 2, 3, 4, 5, 6, 7, 8, 9$). Then among the 10 deviations, independent values can be assumed by any nine. Now, if the occurrences of the 10 digits are random, then $d_i = 0$ in the ideal situation. However, due to chance error, d_i may assume non-zero value.

Thus, d_i 's chance errors but not assignable error if the occurrences of the 10 digits in the set of the N occurrences. The chance variables are assumed to be independently & identically distributed as $N(0, \sigma^2)$. Testing of randomness of occurrences of the 10 digits is equivalent to testing the hypothesis H_0 that $E(d_i) = 0$ for $i = 0, 1, 2, 3, 4, 5, 6, 7, 8, 9$.

Test statistic can be expressed as $t = \frac{\bar{d}}{s/\sqrt{n}} \sim t_{n-2}$, where $\bar{d} = \frac{1}{n} \sum_{i=1}^n d_i$ and $s^2 =$

$\frac{1}{n-1} \sum_{i=1}^n (d_i - \bar{d})^2$

H_0 is rejected at the significance level α if the calculated value of t is found to be exceeding its corresponding theoretical value that corresponds to the level of significance α with $(n-2)$ degrees of freedom.

4. FINDINGS OF THE RANDOMNESS TESTS

4.1 Findings of the χ^2 Test

It is reported for Tippet's random number table that the highest observed chi-square value with 9 degrees of freedom is 15.814, whereas the theoretical value of chi-square with 9 degrees of freedom at 5% level of significance is 16.919. Thus, the lack of randomness of Tippet's random number table was found insignificant at 5% significance level. However, the observed chi-square value corresponds to its theoretical value at 7.5% level of significance. In other words, the lack of randomness of Tippet's random number table can be regarded as

significant at the level of significance $>7.5\%$ and insignificant at the level of significance $< 7.5\%$ [23].

In case of Fisher and Yates random number table, it was observed that the highest observed chi-square value with 9 degrees of freedom is 26.118, which is higher than the theoretical value of chi-square with 9 degrees of freedom at both 5% and 1% significance level (16.919 and 21.666, respectively). Thus, the lack of randomness of Fisher & Yates random number table can be regarded as significant not only at 5% level of significance but also at 1% level. However, the observed chi-square value corresponds to its theoretical value at 0.055% level of significance. In other terms, the lack of randomness of Fisher and Yates random number table can be regarded as significant at the level of significance $> 0.055\%$ and insignificant at the level of significance $< 0.055\%$ [25].

It was mentioned for Kendall and Smith's random number table that the highest observed chi-square value with 9 degrees of freedom is 13.4, which is less than the corresponding theoretical value of chi-square at 5% level of significance. Thus, the lack of randomness of Kendall and Smith's random number table was found insignificant at 5% significance level. However, the observed chi-square value with 9 degrees of freedom namely 13.4 corresponds to the theoretical value of chi-square with 9 degrees of freedom at 18.1% level of significance. Thus, the lack of randomness of Kendall and Smith's random number table can be regarded as significant at the level of significance $>18.1\%$ and insignificant at the level of significance $< 18.1\%$ [23].

For the random number table due to Rand Corporation, it was observed that the highest observed chi-square value with 9 degrees of freedom is 12.518, which is less than the corresponding theoretical value of chi-square at 5% significance level. Thus, the lack of randomness of random number table due to Rand Corporation can be regarded as insignificant at 5% significance level. However, the observed chi-square value with 9 degrees of freedom namely 12.518 corresponds to the theoretical value of chi-square with 9 degrees of freedom at 24% significance level. In other words, the lack of randomness of random number table due to Rand Corporation can be regarded as significant at the level of significance $>24\%$ and insignificant at the level of significance $< 24\%$ [23].

From the findings based on χ^2 test, it is clear that highest degree of randomness is present in random number table due to RAND Corporation followed by due to Kendall and Smith, Tippet, Fisher and Yates, respectively.

4.2 Findings of the Run Test

On comparing the observed values with the corresponding theoretical Z values, the lack of randomness in the three parts containing 19th, 21st and 25th 200 trials respectively in Tippet's random number table can be regarded as significant at 5% level but not at 1% level while the lack of randomness in the other parts can be treated as insignificant [38].

The lack of randomness in Fisher and Yates random number table was found to be non-significant at 5% level by comparing the observed values with the corresponding theoretical Z values [38].

The lack of randomness in the parts containing 1st, 2nd, 26th, 33rd, 36th, 45th, 46th, 48th, 65th, 68th and 70th 200 trials respectively in Kendall and Smith's random number table was found significant at 5% significance level but not at 1%, while the lack of randomness in the other parts of the table can be treated as insignificant [26].

It was found on comparing the observed values with the corresponding theoretical Z values, that the lack of randomness in the four parts containing 35th, 52nd, 73rd, 94th 200 trials respectively in Rand Corporation random number table can be regarded as significant at 5% significance level but not at 1% level, while the lack of randomness in the other parts of the table can be considered as insignificant [38].

From the findings based on run test, it is clear that highest degree of randomness is present in random number table due to Fisher and Yates followed by due to Tippet, RAND Corporation, Kendall and Smith, respectively.

4.3 Findings of the Deviation Test

On comparing the observed values with the corresponding theoretical t values, that the lack of randomness of Tippet's random number table can be treated to be highly significant i.e. significant at both 5% and 1% significance level, except the four parts corresponding to the four sets of trials specifically 1st 2000, 10th 2000, 17th 2000 and last 1600 trials. However, the lack of randomness in these four parts of the table was found significant at 5% level [39].

By comparing the observed values with the corresponding theoretical t values, that the lack of randomness of Fisher and Yates random number table was found to be highly significant i.e. significant at both 5% and 1% significance level, except the two parts corresponding to the two sets of 11th and 13th 1000 trials. However, the lack of randomness in these two parts of the table was found at 5% level of significance [39].

The lack of randomness of Kendall and Smith's random number table can be treated to be highly significant i.e. significant at both 5% and 1% level of significance except the part corresponding to the set of 5th 2000 trials, by comparing the observed values with the corresponding theoretical values of t. However, the lack of randomness in this particular part of the table is significant at 5% level [39].

On comparing the observed values with the corresponding theoretical t values, the lack of randomness of Rand Corporation random number table was found significant both at 5% and 1% level of significance except the five parts corresponding to the five sets viz. 3rd, 7th, 20th, 23rd and 25th sets of 2000 trials. However, the lack of randomness in these five parts of the table was significant at 5% level [39].

From the findings based on the deviation test, it is clear that highest degree of randomness is present in random number table due to Kendall and Smith followed by due to Fisher and Yates, RAND Corporation, Tippet respectively.

5. CONCLUSION

Degree of randomness present in different random number tables (Tippet's random number table, Fisher and Yates random number table, Kendall and Smith's random number table and Random number table of RAND Corporation) based on χ^2 test, run test and deviation test are presented in table 1.

Table 1. Ranks of the four random number tables

Random Number Table	Rank with respect to the degree of presence of randomness based on χ^2 test	Rank with respect to the degree of presence of randomness based on run test	Rank with respect to the degree of presence of randomness based on deviation test
Tippet's random number table	3	2	4
Fisher and Yates random number table	4	1	2
Kendall and Smith's random number table	2	4	1
Random number table due to RAND Corporation	1	3	3

It can be observed that the findings obtained in the studies based on different tests are not alike. Consequently, there is a necessity to search for the reasons of the difference between

these findings. Moreover, it can also be concluded that attempts should be made by the researchers to construct new random numbers table with enhanced degree of randomness than that of the existing tables.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Tippett LHC. 1927. Random Sampling Numbers, Tracts for Computers No 15. Cambridge: Cambridge University Press; 1927.
2. Fisher RA, Yates F. Statistical Tables for Biological, Agricultural and Medical Research. 6th Edition. England: Longman Group Limited; 1982.
3. Kendall MG, Smith BB. Randomness and random sampling numbers. Journal of Royal Statistical Society. 1938; 101(1): 147–166.
4. Kendall MG, Smith BB. A Table of Random Sampling Numbers, Tracts for Computers No 24. Cambridge: Cambridge University Press; 1939.
5. Mahalanobis PC. Tables of random samples from a normal population. Sankhya. 1934; 1: 289–328.
6. Quenouille MH. (1959). Tables of random observations from standard distributions. Biometrika. 1959; 46: 178 – 202.
7. Rand Corporation. A Million Random Digits. Glenoe: Free Press; 1955.
8. Snedecor GW, Cochran WG. Statistical Methods. 6th Edition. Iowa: Iowa State University Press; 1967.
9. Hald A. Statistical Tables and Formulas. Wiley; 1952.
10. Royo J, Ferrer S. (1954). Tables of Random Numbers Obtained from Numbers in the Spanish National Lottery. Trabajos de Estadística. 1954; 5: 247–256.
11. Moses EL, Oakford VR. Tables of Random Permutations. George Allen & Unwin; 1963.
12. Rohlf FJ, Sokal RR. Ten Thousand Random Digits. Freeman; 1969.
13. Manfred M. Le Petit Livre de Nombres au Hasar. Paris: Édition d'artiste; 1971.
14. Rao CR, Mitra SK, Matthai A. Random Numbers and Permutations. Calcutta: Statistical Publishing Society; 1966.
15. Chakrabarty D. Drawing of Random Five-Digit Numbers from Tables of Random Two-Digit and Three-Digit Numbers. International Journal of Advanced Research in Science, Engineering and Technology. 2016; 3(7): 2385 – 2306.
16. Chakrabarty D. Drawing of Random Six-Digit Numbers from Tables of Random Three-Digit Numbers. International Journal of Advanced Research in Science, Engineering and Technology. 2016; 3(8): 2507 – 2515.
17. Chakrabarty D. Drawing of Random Six-Digit Numbers from Tables of Random Two-Digit Numbers. International Journal of Advanced Research in Science, Engineering and Technology. 2016; 3(9): 2643 – 2655.

266 18. Chakrabarty D. Drawing of Random Six-Digit Numbers from a Single Table of
267 Random Two-Digit Numbers. *International Journal of Advanced Research in*
268 *Science, Engineering and Technology*. 2016; 3(10): 2743 – 2753.

269 19. Chakrabarty D. Drawing of Random Six-Digit Numbers from a Single Table of
270 Random Three-Digit Numbers. *International Journal of Advanced Research in*
271 *Science, Engineering and Technology*. 2016; 3(11): 2905 – 2914.

272 20. Chakrabarty D. Drawing of Random Seven-Digit Numbers from Tables of
273 Random Two-Digit Numbers and of Three-Digit Numbers. *International Journal of*
274 *Advanced Research in Science, Engineering and Technology*. 2016; 3(12): 3003 –
275 3012.

276 21. Chakrabarty D. Drawing of Random Four-Digit Numbers from Independent
277 Tables of Random Two-Digit Numbers in Selection of Random Sample. *Biometrics*
278 *& Biostatistics International Journal*. 2016; 4(7): 1-6.

279 22. Chakrabarty D. Drawing of Random Four-Digit Numbers from a Single Table of
280 Random Two-Digit Numbers. *International Journal of Advanced Research in*
281 *Science, Engineering and Technology*. 2017; 4(2): 3877 – 3887.

282 23. Chakrabarty D, Sarma BK. Comparison of degree of randomness of the tables of
283 random numbers due to Tippet, Fisher & Yates, Kendall & Smith and RAND
284 Corporation. *Journal of Reliability and Statistical Studies*. 2017; 10(1): 27-42.

285 24. Nair KR. On Tippetts ‘Random Sampling Numbers’. *Sankhyā: The Indian*
286 *Journal of Statistics*. 1938; 4(1): 65-72.

287 25. Sarmah BK, Chakrabarty D. Testing of Randomness of the Number Generated
288 by Fisher and Yates. *International Journal of Engineering Sciences & Research*
289 *Technology*. 2014; 3(11): 632-636.

290 26. Sarmah BK and Chakrabarty D. Examination of Proper Randomness of Numbers
291 of M. G. Kendall and B. Babington Smith’s Random Numbers Table: Run Test.
292 *International Journal of Multidisciplinary Research and Modern Education*. 2015;
293 1(2): 223-226.

294 27. Moore PG. Reviews: A Million Random Digits with 100,000 Normal Deviates.
295 *Biometrika*. 1955; 42(3-4): 543–543.

296 28. Yates F. Contingency table involving small numbers and the chi square test.
297 *Journal of Royal Statistical Society*. 1934; 1(2): 217–235.

298 29. Corder GW, Foreman DI. *Nonparametric Statistics: A Step-by-Step Approach*.
299 New York: Wiley; 2014.

300 30. Greenwood PE, Nikulin MS. *A Guide to Chi-Squared Testing*. New York:
301 Wiley; 1996.

302 31. Chernoff H, Lehmann EL. The use of maximum likelihood estimates in χ^2 tests
303 for goodness of fit. *The Annals of Mathematical Statistics*. 1954; 25(3): 579–586.

304 32. Gupta SC, Kapoor VK. *Fundamentals of Mathematical Statistics*. 10th ed. New
305 Delhi: Sultan Chand & Sons; 2014.

306 33. Zar JH. *Biostatistical analysis*. 5th ed. Upper Saddle River, N.J.: Prentice-
307 Hall/Pearson; 2010.

308 34. Rangaswamy R. *A Textbook of Agricultural Statistics*. 2nd ed. New Delhi: New
309 Age International Publishers; 2010.

- 310 35. Bradely JV. Distribution Free Statistical Tests. 1st ed. USA: Prentice Hall; 1968.
311 36. Sahu PK, Pal SS, Das AK. Estimation and Inferential Statistics. New Delhi:
312 Springer (India) Pvt. Ltd.; 2015.
313 37. Sahu PK. Applied Statistics for Agriculture, Veterinary, Fishery, Dairy and
314 Allied Fields. Springer Nature: New Delhi; 2016.
315 38. Chakrabarty D. Random Numbers Tables Due to Tippet, Fisher & Yates, Kendall
316 & Smith and Rand Corporation: Comparison of Degree of Randomness by Run Test.
317 Journal of Biostatistics and Biometric Applications. 2018; 3(1): 1-8.
318 39. Chakrabarty D. Deviation Test: Comparison of Degree of Randomness of the
319 Tables of Random Numbers due to Tippet, Fisher & Yates, Kendall & Smith and
320 Rand Corporation. SM Journal of Biometrics & Biostatistics. 2017; 2(3): 1-6.
321