

Analytical study of the Difference Equation

$$x_{n+1} = \frac{x_n x_{n-6}}{x_{n-5} (\pm 1 - x_n x_{n-6})}$$

ABSTRACT

This paper is devoted to find the form of the solutions of the following rational difference equations:

$$x_{n+1} = \frac{x_n x_{n-6}}{x_{n-5} (\pm 1 - x_n x_{n-6})}, \quad n = 0, 1, \dots,$$

where the initial conditions $x_{-6}, x_{-5}, x_{-4}, x_{-3}, x_{-2}, x_{-1}, x_0$ are arbitrary positive real numbers. Also, we give specific form of the solutions of two special cases of this equation.

Keywords: Difference equations, Recursive sequences, Stability, Boundedness.

2000 Mathematics Subject Classification: 39A10

1. INTRODUCTION

Our aim in this paper is to investigate the behavior of the solution of the following nonlinear difference equation

$$x_{n+1} = \frac{x_n x_{n-6}}{x_{n-5} (\pm 1 - x_n x_{n-6})}, \quad n = 0, 1, \dots, \quad (1)$$

where the initial conditions $x_{-6}, x_{-5}, x_{-4}, x_{-3}, x_{-2}, x_{-1}, x_0$ are arbitrary positive real numbers.

The study and solution of nonlinear rational recursive sequence of high order is quite challenging and rewarding. Recently, there has been a lot of interest in studying the qualitative properties of rational recursive sequences, Furthermore diverse nonlinear trend occurring in science and engineering can be modeled by such equations and the solution about such equations offer prototypes towards the development of the theory. However, there have not been any suitable general method to deal with the global behavior of rational difference equations of high order so far. Therefore, the study of rational difference equations of order greater than one is worth further consideration.

Many researchers have investigated the behavior of the solution of difference equations, for example,

Elsayed et al. [23] has obtained results concerning the dynamics and solution of the rational difference equation

$$x_{n+1} = \frac{x_{n-1} x_{n-5}}{x_{n-3} (\pm 1 \pm x_{n-1} x_{n-5})}.$$

Elsayed et al. [27] has obtained results concerning The dynamics and the solutions of the rational difference equation

$$x_{n+1} = \frac{x_n x_{n-4}}{x_{n-3} (\pm 1 \pm x_n x_{n-4})}.$$

Aloqeili [3] has obtained the solutions of the difference equation

$$x_{n+1} = \frac{x_{n-1}}{a - x_n x_{n-1}}.$$

Simsek et al. [37] obtained the solution of the difference equation

$$x_{n+1} = \frac{x_{n-3}}{1 + x_{n-1}}$$

Çinar [7,8,9] got the solutions of the following difference equation

$$x_{n+1} = \frac{x_{n-1}}{1 + ax_n x_{n-1}}, x_{n+1} = \frac{x_{n-1}}{-1 + ax_n x_{n-1}}, x_{n+1} = \frac{ax_{n-1}}{1 + bx_n x_{n-1}}$$

In [28], Ibrahim got the form of the solution of the rational difference equation

$$x_{n+1} = \frac{x_n x_{n-2}}{x_{n-1} (a + bx_n x_{n-2})}$$

Karatas et al. [30] got the solution of the difference equation

$$x_{n+1} = \frac{x_{n-5}}{1 + x_{n-2} x_{n-5}}$$

Here, we recall some notations and results which will be useful in our investigation. Let I be some interval of real numbers and let

$$f : I^{k+1} \rightarrow I,$$

be a continuously differentiable function. Then for every set of initial conditions $x_{-k}, x_{-k+1}, x_{-k+2}, \dots, x_0 \in I$, the difference equation

$$x_{n+1} = f(x_n, x_{n-1}, \dots, x_{n-k}), \quad n = 0, 1, \dots, \quad (2)$$

has a unique solution $\{x_n\}_{n=-k}^{\infty}$.

Definition 1. (Equilibrium Point) A point $\bar{x} \in I$ is called an equilibrium point of Eq. (2) if $\bar{x} = f(\bar{x}, \bar{x}, \dots, \bar{x})$. That is, $x_n = \bar{x}$ for $n \geq 0$, is a solution of Eq. (2), or equivalently, \bar{x} is a fixed point of f .

Definition 2. (Stability)

- The equilibrium point \bar{x} of Eq. (2) is locally stable if for every $\varepsilon > 0$, there exists $\delta > 0$ such that for all $x_{-k}, x_{-k+1}, x_{-k+2}, \dots, x_0 \in I$, with

$$|x_{-k} - \bar{x}| + |x_{-k+1} - \bar{x}| + |x_{-k+2} - \bar{x}| + \dots + |x_0 - \bar{x}| < \delta,$$

we have $|x_n - \bar{x}| < \varepsilon$, for all $n \geq -k$.

- The equilibrium point \bar{x} of Eq. (2) is locally asymptotically stable if \bar{x} is locally stable solution of Eq. (2) and there exists $\gamma > 0$, such that for all $x_{-k}, x_{-k+1}, x_{-k+2}, \dots, x_0 \in I$, with

$$|x_{-k} - \bar{x}| + |x_{-k+1} - \bar{x}| + |x_{-k+2} - \bar{x}| + \dots + |x_0 - \bar{x}| < \delta,$$

we have $\lim_{n \rightarrow \infty} x_n = \bar{x}$.

- The equilibrium point \bar{x} of Eq. (2) is global attractor if for all $x_{-k}, x_{-k+1}, \dots, x_0 \in I$ we have

$$\lim_{n \rightarrow \infty} x_n = \bar{x}.$$

- The equilibrium point \bar{x} of Eq. (2) is globally asymptotically stable if \bar{x} is locally stable, and \bar{x} is also a global attractor of Eq. (2).
- The equilibrium point \bar{x} of Eq. (2) is unstable if \bar{x} is not locally stable.

The linearized equation of Eq. (2) about the equilibrium \bar{x} is the linear difference equation

$$y_{n+1} = \sum_{i=0}^k \frac{\partial f(\bar{x}, \bar{x}, \dots, \bar{x})}{\partial x_{n-i}} y_{n-i}$$

Theorem 1. Assume that $p, q \in \mathbb{R}$ and $k \in \{0, 1, 2, \dots\}$. Then $|p| + |q| < 1$ is a sufficient condition for the asymptotic stability of the difference equation

$$x_{n+1} + px_n + qx_{n-k} = 0, n = 0, 1, \dots$$

Remark 1. The theorem can be easily extended to a general linear equations of the form

$$x_{n+k} + p_1 x_{n+k-1} + \dots + p_k x_n = 0, \quad n = 0, 1, \dots, \quad (3)$$

where $p_1, p_2, \dots, p_k \in \mathbb{R}$ and $k \in \{0, 1, 2, \dots\}$. Then Eq. (3) is asymptotically stable provided that $\sum_{i=0}^k |p_i| < 1$.

Our goal in this section is to find a specific form of the solutions of some special cases of Eq. (1) when a, b, c and d are integers and give numerical examples of each case.

2. FIRST CASE : ON THE DIFFERENCE EQUATION $X_{N+1} = \frac{X_N X_{N-6}}{X_{N-5}(1 - X_N X_{N-6})}$

In this subsection we study the following special case of Eq. (1):

$$x_{n+1} = \frac{x_n x_{n-6}}{x_{n-5}(1 - x_n x_{n-6})}, \quad n = 0, 1, \dots, \quad (4)$$

where the initial conditions $x_{-6}, x_{-5}, x_{-4}, x_{-3}, x_{-2}, x_{-1}, x_0$ are arbitrary nonzeros real numbers.

Theorem 2. Let $\{x_n\}_{n=-4}^{\infty}$ be a solution of Eq. (4). Then for $n = 0, 1, \dots$,

$$\begin{aligned} x_{12n-6} &= x_{-6} \prod_{i=0}^{n-1} \left(\frac{1 - (12i)x_{-6}x_0}{1 - (12i+6)x_{-6}x_0} \right), & x_{12n-5} &= x_{-5} \prod_{i=0}^{n-1} \left(\frac{1 - (12i+1)x_{-6}x_0}{1 - (12i+7)x_{-6}x_0} \right), \\ x_{12n-4} &= x_{-4} \prod_{i=0}^{n-1} \left(\frac{1 - (12i+2)x_{-6}x_0}{1 - (12i+8)x_{-6}x_0} \right), & x_{12n-3} &= x_{-3} \prod_{i=0}^{n-1} \left(\frac{1 - (12i+3)x_{-6}x_0}{1 - (12i+9)x_{-6}x_0} \right), \\ x_{12n-2} &= x_{-2} \prod_{i=0}^{n-1} \left(\frac{1 - (12i+4)x_{-6}x_0}{1 - (12i+10)x_{-6}x_0} \right), & x_{12n-1} &= x_{-1} \prod_{i=0}^{n-1} \left(\frac{1 - (12i+5)x_{-6}x_0}{1 - (12i+11)x_{-6}x_0} \right), \\ x_{12n} &= x_0 \prod_{i=0}^{n-1} \left(\frac{1 - (12i+6)x_{-6}x_0}{1 - (12i+12)x_{-6}x_0} \right), & x_{12n+1} &= \frac{x_{-6}x_0}{x_{-5}(1 - x_{-6}x_0)} \prod_{i=0}^{n-1} \left(\frac{1 - (12i+7)x_{-6}x_0}{1 - (12i+13)x_{-6}x_0} \right), \\ x_{12n+2} &= \frac{x_{-6}x_0}{x_{-4}(1 - 2x_{-6}x_0)} \prod_{i=0}^{n-1} \left(\frac{1 - (12i+8)x_{-6}x_0}{1 - (12i+14)x_{-6}x_0} \right), & x_{12n+3} &= \frac{x_{-6}x_0}{x_{-3}(1 - 3x_{-6}x_0)} \prod_{i=0}^{n-1} \left(\frac{1 - (12i+9)x_{-6}x_0}{1 - (12i+15)x_{-6}x_0} \right), \\ x_{12n+4} &= \frac{x_{-6}x_0}{x_{-2}(1 - 4x_{-6}x_0)} \prod_{i=0}^{n-1} \left(\frac{1 - (12i+10)x_{-6}x_0}{1 - (12i+16)x_{-6}x_0} \right), & x_{12n+5} &= \frac{x_{-6}x_0}{x_{-1}(1 - 5x_{-6}x_0)} \prod_{i=0}^{n-1} \left(\frac{1 - (12i+11)x_{-6}x_0}{1 - (12i+17)x_{-6}x_0} \right). \end{aligned}$$

Proof: We use an inductive proof for this rational recursive sequences. It is easy to see that for $n = 0$, the result holds. Suppose that $n > 0$ and that the assumption is satisfied for $n - 1$. That is;

$$\begin{aligned}
 x_{12n-18} &= x_{-6} \prod_{i=0}^{n-2} \left(\frac{1 - (12i)x_{-6}x_0}{1 - (12i+6)x_{-6}x_0} \right), & x_{12n-17} &= x_{-5} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+1)x_{-6}x_0}{1 - (12i+7)x_{-6}x_0} \right), \\
 x_{12n-16} &= x_{-4} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+2)x_{-6}x_0}{1 - (12i+8)x_{-6}x_0} \right), & x_{12n-15} &= x_{-3} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+3)x_{-6}x_0}{1 - (12i+9)x_{-6}x_0} \right), \\
 x_{12n-14} &= x_{-2} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+4)x_{-6}x_0}{1 - (12i+10)x_{-6}x_0} \right), & x_{12n-13} &= x_{-1} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+5)x_{-6}x_0}{1 - (12i+11)x_{-6}x_0} \right), \\
 x_{12n-12} &= x_0 \prod_{i=0}^{n-2} \left(\frac{1 - (12i+6)x_{-6}x_0}{1 - (12i+12)x_{-6}x_0} \right), & x_{12n-11} &= \frac{x_{-6}x_0}{x_{-5}(1 - x_{-6}x_0)} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+7)x_{-6}x_0}{1 - (12i+13)x_{-6}x_0} \right), \\
 x_{12-10} &= \frac{x_{-6}x_0}{x_{-4}(1 - 2x_{-6}x_0)} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+8)x_{-6}x_0}{1 - (12i+14)x_{-6}x_0} \right), & x_{12n-9} &= \frac{x_{-6}x_0}{x_{-3}(1 - 3x_{-6}x_0)} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+9)x_{-6}x_0}{1 - (12i+15)x_{-6}x_0} \right), \\
 x_{12n-8} &= \frac{x_{-6}x_0}{x_{-2}(1 - 4x_{-6}x_0)} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+10)x_{-6}x_0}{1 - (12i+16)x_{-6}x_0} \right), & x_{12n-7} &= \frac{x_{-6}x_0}{x_{-1}(1 - 5x_{-6}x_0)} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+11)x_{-6}x_0}{1 - (12i+17)x_{-6}x_0} \right).
 \end{aligned}$$

Now, using the main Eq. (4), one has

$$\begin{aligned}
 x_{12n-6} &= \frac{x_{12n-7}x_{12n-13}}{x_{12n-12}(1 - x_{12n-7}x_{12n-13})} \\
 &= \frac{\frac{x_{-6}x_0}{x_{-1}(1 - 5x_{-6}x_0)} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+11)x_{-6}x_0}{1 - (12i+17)x_{-6}x_0} \right) x_{-1} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+5)x_{-6}x_0}{1 - (12i+11)x_{-6}x_0} \right)}{x_0 \prod_{i=0}^{n-2} \left(\frac{1 - (12i+6)x_{-6}x_0}{1 - (12i+12)x_{-6}x_0} \right) \left(1 - \frac{x_{-6}x_0}{x_{-1}(1 - 5x_{-6}x_0)} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+11)x_{-6}x_0}{1 - (12i+17)x_{-6}x_0} \right) x_{-1} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+5)x_{-6}x_0}{1 - (12i+11)x_{-6}x_0} \right) \right)} \\
 &= \frac{\frac{x_{-6}}{(1 - 5x_{-6}x_0)} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+5)x_{-6}x_0}{1 - (12i+17)x_{-6}x_0} \right)}{\prod_{i=0}^{n-2} \left(\frac{1 - (12i+6)x_{-6}x_0}{1 - (12i+12)x_{-6}x_0} \right) \left(1 - \frac{x_{-6}x_0}{(1 - 5x_{-6}x_0)} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+5)x_{-6}x_0}{1 - (12i+17)x_{-6}x_0} \right) \right)} \\
 &= \prod_{i=0}^{n-2} \left(\frac{1 - (12i+12)x_{-6}x_0}{1 - (12i+6)x_{-6}x_0} \right) \frac{\frac{x_{-6}}{(1 - 5x_{-6}x_0)} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+5)x_{-6}x_0}{1 - (12i+17)x_{-6}x_0} \right)}{\left(1 - \frac{x_{-6}x_0}{(1 - 5x_{-6}x_0)} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+5)x_{-6}x_0}{1 - (12i+17)x_{-6}x_0} \right) \right)} \\
 &= \prod_{i=0}^{n-2} \left(\frac{1 - (12i+12)x_{-6}x_0}{1 - (12i+6)x_{-6}x_0} \right) \frac{1}{\left(1 - \frac{x_{-6}x_0}{(1 - (12n-7)x_{-6}x_0)} \right)} \\
 &= \prod_{i=0}^{n-2} \left(\frac{1 - (12i+12)x_{-6}x_0}{1 - (12i+6)x_{-6}x_0} \right) \frac{x_{-6}}{1 - (12n-6)x_{-6}x_0}.
 \end{aligned}$$

Hence, we have

$$x_{12n-6} = x_{-6} \prod_{i=0}^{n-1} \left(\frac{1 - (12i)x_{-6}x_0}{1 - (12i+6)x_{-6}x_0} \right).$$

Similarly, using the main Eq. (4), one has

$$\begin{aligned} x_{12n-5} &= \frac{x_{12n-6}x_{12n-12}}{x_{12n-11}(1 - x_{12n-6}x_{12n-12})} \\ &= \frac{x_{-6} \prod_{i=0}^{n-1} \left(\frac{1 - (12i)x_{-6}x_0}{1 - (12i+6)x_{-6}x_0} \right) x_0 \prod_{i=0}^{n-2} \left(\frac{1 - (12i+6)x_{-6}x_0}{1 - (12i+12)x_{-6}x_0} \right)}{\frac{x_{-6}x_0}{x_{-5}(1 - x_{-6}x_0)} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+7)x_{-6}x_0}{1 - (12i+13)x_{-6}x_0} \right) \left(1 - x_{-6} \prod_{i=0}^{n-1} \left(\frac{1 - (12i)x_{-6}x_0}{1 - (12i+6)x_{-6}x_0} \right) x_0 \prod_{i=0}^{n-2} \left(\frac{1 - (12i+6)x_{-6}x_0}{1 - (12i+12)x_{-6}x_0} \right) \right)} \\ &= x_{-5}(1 + x_{-6}x_0) \prod_{i=0}^{n-2} \left(\frac{1 - (12i+13)x_{-6}x_0}{1 - (12i+7)x_{-6}x_0} \right) \frac{\left(\frac{1 - (12n-12)x_{-6}x_0}{1 - (12n-6)x_{-6}x_0} \right) \prod_{i=0}^{n-2} \left(\frac{1 - (12i)x_{-6}x_0}{1 - (12i+12)x_{-6}x_0} \right)}{\left(1 - x_{-6}x_0 \left(\frac{1 - (12n-12)x_{-6}x_0}{1 - (12n-6)x_{-6}x_0} \right) \prod_{i=0}^{n-2} \left(\frac{1 - (12i)x_{-6}x_0}{1 - (12i+12)x_{-6}x_0} \right) \right)} \\ &= x_{-5} \prod_{i=0}^{n-2} \left(\frac{1 - (12i+1)x_{-6}x_0}{1 - (12i+7)x_{-6}x_0} \right) \frac{1 - x_{-6}x_0}{1 - (12n-5)x_{-6}x_0}. \end{aligned}$$

Hence, we have

$$x_{12n-5} = x_{-5} \prod_{i=0}^{n-1} \left(\frac{1 - (12i+1)x_{-6}x_0}{1 - (12i+7)x_{-6}x_0} \right).$$

Similarly, one can easily obtain the other relations. Thus, the proof is completed.

Theorem 3. Eq. (4) has one equilibrium point $\bar{x} = 0$ and this equilibrium point is not locally asymptotically stable.

Proof. In this section we investigate the local stability character of the solutions of Eq. (4). Equation (4) has a unique positive equilibrium point and is given by

$$\bar{x} = \frac{\bar{x}^2}{\bar{x}(1 - \bar{x}^2)} = \frac{\bar{x}}{1 - \bar{x}^2},$$

or also

$$1 = 1 - \bar{x}^2,$$

then the unique equilibrium point is given by $\bar{x} = 0$.

Define the following function

$$\begin{aligned} f &: (0, \infty)^3 \rightarrow (0, \infty) \\ f(u, v, w) &= \frac{uw}{v(1 - uw)}. \end{aligned}$$

Therefore it follows that

$$f_u(u, v, w) = \frac{w}{v(1-uw)^2}, \quad f_v(u, v, w) = -\frac{uw}{v^2(1-uw)}, \quad f_w(u, v, w) = \frac{u}{v(1-uw)^2}.$$

Then

$$f_u(\bar{x}, \bar{x}, \bar{x}) = \frac{1}{(1-\bar{x}^2)^2} = 1, \quad f_v(\bar{x}, \bar{x}, \bar{x}) = -\frac{1}{(1-\bar{x}^2)} = -1, \quad f_w(\bar{x}, \bar{x}, \bar{x}) = \frac{1}{(1-\bar{x}^2)^2} = 1.$$

The linearized equation of Eq. (4) about \bar{x} is

$$y_{n+1} - y_{n-6} + y_{n-5} - y_n = 0. \quad (5)$$

It follows from Theorem 1 that Eq. (5) is not asymptotically stable. The proof is complete.

For confirming the results of this section, we consider numerical example for ,(See Figure 1).




Figure 1. Left $x_{-6} = 1, x_{-5} = 2, x_{-4} = 3, x_{-3} = 4, x_{-2} = 5, x_{-1} = 6, x_0 = 7$ and right $x_{-6} = 11, x_{-5} = 12, x_{-4} = 13, x_{-3} = 4, x_{-2} = 15, x_{-1} = 6, x_0 = 17$

3. SECOND CASE : ON THE DIFFERENCE EQUATION $X_{N+1} = \frac{X_N X_{N-6}}{X_{N-5}(-1 - X_N X_{N-6})}$

In this subsection we study the following special case of Eq. (1):

$$x_{n+1} = \frac{x_n x_{n-6}}{x_{n-5}(-1 - x_n x_{n-6})}, \quad n = 0, 1, \dots, \quad (6)$$

where the initial conditions $x_{-6}, x_{-5}, x_{-4}, x_{-3}, x_{-2}, x_{-1}, x_0$ are arbitrary nonzeros real numbers.

Theorem 4. Let $\{x_n\}_{n=-4}^{\infty}$ be a solution of Eq. (6). Then the solution of Eq. (6) is bounded and periodic of period 12 given by:

$$\begin{aligned} x_{12n-6} &= x_{-6}, & x_{12n-5} &= x_{-5}, & x_{12n-4} &= x_{-4}, & x_{12n-3} &= x_{-3}, \\ x_{12n-2} &= x_{-2}, & x_{12n-1} &= x_{-1}, & x_{12n} &= x_0, & x_{12n+1} &= \frac{x_0 x_{-6}}{x_{-5}(-1 - x_0 x_{-6})}, \\ x_{12n+2} &= \frac{x_0 x_{-6}}{x_{-4}}, & x_{12n+3} &= \frac{x_0 x_{-6}}{x_{-3}(-1 - x_0 x_{-6})}, & x_{12n+4} &= \frac{x_0 x_{-6}}{x_{-2}}, & x_{12n+5} &= \frac{x_0 x_{-6}}{x_{-1}(-1 - x_0 x_{-6})}. \end{aligned}$$

Proof. For $n = 0$, the result holds. Now suppose that our assumption holds for $n - 1$. That is;

$$\begin{aligned} x_{12n-18} &= x_{-6}, & x_{12n-17} &= x_{-5}, & x_{12n-16} &= x_{-4}, & x_{12n-15} &= x_{-3}, \\ x_{12n-14} &= x_{-2}, & x_{12n-13} &= x_{-1}, & x_{12n-12} &= x_0, & x_{12n-11} &= \frac{x_0 x_{-6}}{x_{-5}(-1 - x_0 x_{-6})}, \\ x_{12n-10} &= \frac{x_0 x_{-6}}{x_{-4}}, & x_{12n-9} &= \frac{x_0 x_{-6}}{x_{-3}(-1 - x_0 x_{-6})}, & x_{12n-8} &= \frac{x_0 x_{-6}}{x_{-2}}, & x_{12n-7} &= \frac{x_0 x_{-6}}{x_{-1}(-1 - x_0 x_{-6})}. \end{aligned}$$

Now it follows from Eq. (6) that

$$\begin{aligned} x_{12n-6} &= \frac{x_{12n-7} x_{12n-13}}{x_{12n-12}(-1 - x_{12n-7} x_{12n-13})} = \frac{\frac{x_0 x_{-6}}{x_{-1}(-1 - x_0 x_{-6})} x_{-1}}{x_0 \left(-1 - \frac{x_0 x_{-6}}{x_{-1}(-1 - x_0 x_{-6})} x_{-1} \right)} \\ &= \frac{\frac{x_{-6}}{(-1 - x_0 x_{-6})}}{\left(-1 - \frac{x_0 x_{-6}}{(-1 - x_0 x_{-6})} \right)} = \frac{\frac{x_{-6}}{(-1 - x_0 x_{-6})}}{\left(\frac{1}{(-1 - x_0 x_{-6})} \right)} = x_{-6} \end{aligned}$$

Similarly

$$\begin{aligned} x_{12n-5} &= \frac{x_{12n-6} x_{12n-12}}{x_{12n-11}(-1 - x_{12n-6} x_{12n-12})} = \frac{\frac{x_{-6} x_0}{\frac{x_0 x_{-6}}{x_{-5}(-1 - x_0 x_{-6})}(-1 - x_{-6} x_0)}}{x_{-5}(-1 - x_0 x_{-6})} = x_{-5} \\ x_{12n-4} &= \frac{x_{12n-5} x_{12n-11}}{x_{12n-10}(-1 - x_{12n-5} x_{12n-11})} = \frac{\frac{x_{-5} \frac{x_0 x_{-6}}{x_{-5}(-1 - x_0 x_{-6})}}{\frac{x_0 x_{-6}}{x_{-4}} \left(-1 - x_{-5} \frac{x_0 x_{-6}}{x_{-5}(-1 - x_0 x_{-6})} \right)}}{\frac{1}{\frac{x_{-4}}{(-1 - x_0 x_{-6})}}} \\ &= \frac{\frac{1}{\frac{x_{-4}}{(-1 - x_0 x_{-6})}}}{\frac{1}{\frac{x_{-4}}{(-1 - x_0 x_{-6})}}} = \frac{\frac{1}{(-1 - x_0 x_{-6})}}{\left(\frac{1}{(-1 - x_0 x_{-6})} \right)} = x_{-4} \end{aligned}$$

Similarly, one can easily obtain the other relations. Thus, the proof is completed.

Theorem 5. Eq. (6) has a unique equilibrium point which is 0 and this equilibrium point is not locally asymptotically stable.

Proof. As the proof of Theorem 3 and will be omitted.

For confirming the results of this section, we consider the following numerical examples,(See Figure 2).

4. CONCLUSION

This paper discussed local stability, the solutions of some special cases of Eq. (1) and gave numerical examples of each case.




Figure 2. Left $x_{-6} = 1, x_{-5} = 2, x_{-4} = 3, x_{-3} = 4, x_{-2} = 5, x_{-1} = 6, x_0 = 7$ and right $x_{-6} = 5, x_{-5} = 3, x_{-4} = 4, x_{-3} = 1, x_{-2} = 1, x_{-1} = 3, x_0 = 4$

REFERENCES

1. M. Abu Alhalawa and M. Salah, *Dynamics of higher order rational difference equation*, J. Nonlinear Anal. Appl., 8(2)(2017), 363-379.
2. A. M. Ahmed and A. M. Youssef, *A solution form of a class of higher-order rational difference equations*, J. Egyptian. Math. Soc., 21(2013), 248-253.
3. M. Aloqeili, *Dynamics of a rational difference equation*, Appl. Math. Comp., 176(2)(2006), 768-774.
4. N. Battaloglu, C. Cinar, and I. Yalçinkaya, *The Dynamics of the Difference Equation*, Ars Combinatoria, 97(2010), 281-288.
5. F. Belhannache, N. Touafek and R. Abo-Zeid, *On a higher-order rational difference equation*, J. Appl. Math. & informatics, 34(5-6)(2016), 369-382.
6. F. Bozkurt, I. Ozturk and S. Ozen, *The global behavior of the difference equation*, Stud. Univ. Babeş-Bolyai Math., 54(2)(2009), 3-12.
7. C. Cinar, *On the positive solutions of the difference equation $x_{n+1} = \frac{x_{n-1}}{1+ax_nx_{n-1}}$* , Appl. Math. Comp., 158(3)(2004), 809-812.
8. C. Cinar, *On the solutions of the difference equation $x_{n+1} = \frac{x_{n-1}}{-1+ax_nx_{n-1}}$* , Appl. Math. Comp., 158(3)(2004), 793-797.
9. C. Cinar, *On the positive solutions of the difference equation $x_{n+1} = \frac{ax_{n-1}}{1+bx_nx_{n-1}}$* , Appl. Math. Comp., 156(2)(2004), 587-590.
10. E. M. Elabbasy and E. M. Elsayed, *Dynamics of a rational difference equation*, Chinese Annals of Mathematics, Series B, 30(2009), 187-198.
11. E. M. Elabbasy, H. El-Metwally, E. M. Elsayed, *On the difference equations $x_{n+1} = \frac{\alpha x_{n-k}}{\beta + \gamma \prod_{i=0}^k x_{n-i}}$* , J. Conc. Appl. Math., 5(2)(2007), 101-113.
12. E. M. Elabbasy, H. El-Metwally, E. M. Elsayed, *On the difference equation $x_{n+1} = ax_n - \frac{bx_n}{cx_n - dx_{n-1}}$* , Adv. Differ. Eq., 2006(2006), Article ID 82579, 1-10.
13. M. M. El-Dessoky, *On the difference equation $x_{n+1} = ax_{n-1} + bx_{n-k} + \frac{cx_{n-s}}{dx_{n-s} - e}$* , Math. Meth. Appl. Sci., 40(2017), 535-545.
14. H. El-Metwally and E. M. Elsayed, *Form of solutions and periodicity for systems of difference equations*, Journal of Computational Analysis and Applications, 15(5) (2013), 852-857.
15. E. M. Elsayed, *Behavior and Expression of the Solutions of Some Rational Difference Equations*, Journal of Computational Analysis and Applications, 15 (1) (2013), 73-81.
16. E. M. Elsayed, *Qualitative properties for a fourth order rational difference equation*, Acta Applicandae Mathematicae, 110(2)(2010), 589-604.

17. E. M. Elsayed, *On the Global attractivity and the solution of recursive sequence*, Studia Scientiarum Mathematicarum Hungarica, 47(3)(2010), 401-418.
18. E. M. Elsayed, *Qualitative behavior of difference equation of order three*, Acta Scientiarum Mathematicarum (Szeged), 75(1-2)(2009), 113-129.
19. E. M. Elsayed, *Dynamics of a rational recursive sequence*, Inter. J. Differ. Equations, 4(2)(2009), 185-200.
20. E. M. Elsayed, *Qualitative behavior of s rational recursive sequence*, Indagationes Mathematicae, New Series, 19(2)(2008), 189-201.
21. E. M. Elsayed, F. Alzahrani and H. S. Alayachi, *Formulas and properties of some class of nonlinear difference equations*, Journal of Computational Analysis and Applications, 24 (8) (2018), 1517-1531.
22. E. M. Elsayed and A. Khaliq, *The Dynamics and Global attractivity of a Rational Difference Equation*, Advanced Studies in Contemporary Mathematics, 26(1)(2016), 183-202.
23. E. M. Elsayed and A. Khaliq, *The dynamics and solution of some difference equations*, J. Nonlinear Sci. Appl, 9(2016), 1052-1063.
24. E. M. Elsayed and T. F. Ibrahim, *Periodicity and solutions for some systems of nonlinear rational difference equations*, Hacettepe Journal of Mathematics and Statistics, 44 (6) (2015), 1361-1390.
25. E. M. Elsayed and T. F. Ibrahim, *Solutions and Periodicity of a Rational Recursive Sequences of Order Five*, Bulletin of the Malaysian Mathematical Sciences Society, 38 (1) (2015), 95-112.
26. E. M. Elsayed and A. M. Ahmed, *Dynamics of a three-dimensional systems of rational difference equations*, Mathematical Methods in The Applied Sciences, 39 (5) (2016), 1026-1038.
27. E. M. Elsayed, S.R. Mahmoud and A.T. Ali, *The Dynamics and the Solutions of some Rational Difference Equations*, Journal of Computational Analysis and Applications, 18 (3) (2015), 430-439.
28. T. F. Ibrahim, *Closed Form Expressions of some systems of Nonlinear Partial Difference Equations*, Journal of Computational Analysis and Applications, 23 (3) (2017), 433-445.
29. R. Karatas, *On solutions of the difference equation $x_{n+1} = \frac{(-1)^n x_{n-4}}{1 + (-1)^n x_n x_{n-1} x_{n-2} x_{n-3} x_{n-4}}$* , Selcuk J. Appl. Math., 8(1)(2007), 51-56.
30. R. Karatas, C. Cinar and D. Simsek, *On positive solutions of the difference equation $x_{n+1} = \frac{x_{n-5}}{1 + x_{n-2} x_{n-5}}$* , Int. J. Contemp. Math. Sci., 1(10)(2006), 495-500.
31. V. L. Kocic and G. Ladas, *Global Behavior of Nonlinear Difference Equations of Higher Order with Applications*, Kluwer Academic Publishers, Dordrecht, 1993.
32. M. R. S. Kulenovic and G. Ladas, *Dynamics of Second Order Rational Difference Equations with Open Problems and Conjectures*, Chapman & Hall / CRC Press, 2001.
33. K. Liu, P. Li, F. Han and W. Zhong, *Global Dynamics of Nonlinear Difference Equation $x_{n+1} = \frac{x_n}{x_{n-1} x_{n-2}}$* , Journal of Computational Analysis and Applications, 24 (6) (2018), 1125-1132.
34. O. Ocalan, H. Ogunmez and M. Gumus, *Global behavior test for a non-linear difference equation with a period-two coefficient*, Dynamics of Continuous, Discrete and Impulsive Systems Series A: Mathematical Analysis, 21 (2014), 307-316.
35. M. Saleh and S. Abu-Baha, *Dynamics of a higher order rational difference equation*, Appl. Math. Comp., 181(1)(2006), 84-102.
36. D. Simsek and F. Abdullayev *On the recursive sequence $x_{n+1} = \frac{x_{n-(4k+3)}}{1 + \prod_{t=1}^2 x_{n-(k+1)t-k}}$* , J. Math. Sci., 6(222)(2017), 762-771.
37. D. Simsek, C. Cinar and I. Yalcinkaya, *On the recursive sequence $x_{n+1} = \frac{x_{n-3}}{1 + x_{n-1}}$* , Int. J. Contemp. Math. Sci., 1(10)(2006), 475-480.
38. T. Sun, Y. Zhou, G. Su and Bin Qin, *Eventual periodicity of a max-type difference equation system*, Journal of Computational Analysis and Applications, 24 (5) (2018), 976-983.
39. C. Y. Wang, X. J. Fang and R. Li, *On the solution for a system of two rational difference equations*, Journal of Computational Analysis and Applications, 20 (1) (2016) 175-186.

40. C. Y. Wang, X. J. Fang and R. Li, *On the dynamics of a certain four-order fractional difference equations*, Journal of Computational Analysis and Applications, 22 (5) (2017), 968-976.
41. C. Wang, Y. Zhou, S. Pan and R. Li, *On a system of three max-type nonlinear difference equations*, Journal of Computational Analysis and Applications, 25 (8) (2018), 1463-1479.
42. L. Xianyi and Z. Deming, *Global asymptotic stability in a rational equation*, J. Differ. Equations Appl., 9(9)(2003), 833-839.
43. X. Yan and W. Li, *Global attractivity for a class of nonlinear difference equations*, Soochow J. Math., 29(3)(2003), 327-338.
44. Y. Yazlik, E. M. Elsayed and N. Taskara, *On the Behaviour of the Solutions of Difference Equation Systems*, Journal of Computational Analysis and Applications, 16 (5) (2014), 932-941.
45. T. Yi and Z. Zhou, *Periodic solutions of difference equations*, J. Math. Anal. Appl., 286(2003), 220-229.
46. E. M. E. Zayed and M. A. El-Moneam, *On the qualitative study of the nonlinear difference equation $x_{n+1} = \frac{\alpha x_n - \sigma}{\beta + \gamma x_{n-\tau}^p}$* , Fasciculi Mathematici, 50(2013), 137-147.
47. E. M. E. Zayed and M. A. El-Moneam, *On the global attractivity of two nonlinear difference equations*, J Math. Sci. 177(2011) 487.