

2

3 **Response of different irrigation levels on vegetative parameters of Sweet**
4 **Cherry grown in a high density planting system**

5

6 **Abstract**

7 A field experiment was conducted on four year old plants of Sweet cherry cv. Regina grafted
8 on Gisela- 5 rootstock at SKUAST-K Shalimar Srinagar Jammu and Kashmir during the year
9 2016-17 and 2017-18. The experiment consisted of four irrigation treatment combinations I₀
10 (0%), I₁ (50%), I₂ (75%) and I₃ (100%) based on Class A Pan Evaporation percentages (0%,
11 50%, 75% and 100%) that were applied at four growth stages viz Fruit set stage (T₁), Pit
12 hardening stage (T₂), Fruit growth stage (T₃) and Fruit bud differentiation stage (T₄). The
13 quantity of water required was applied through drip irrigation on daily basis as per crop
14 evapotranspiration. The difference between water levels was 25%. The vegetative parameters
15 such as plant height, plant girth, trunk cross-sectional area (TCSA) and annual shoot
16 extension growth of young dwarf sweet cherry plants cv. Regina on Gisela-5 were
17 investigated in temperate climate. The experiment was laid out in RBD design with three
18 replications. Maximum average values of vegetative growth parameters were obtained in I₃
19 treatment followed by I₂. Highest plant height and annual shoot extension growth was
20 recorded at T₁ stage however maximum plant girth was recorded at T₃ stage and highest
21 TCSA was obtained at T₄ stage. Furthermore the highest plant girth was recorded in I₃T₃
22 combination. Plants treated with 100% and 75% ETc level of irrigation excelled in vegetative
23 growth parameters.

24 **Keywords:** Evapotranspiration; irrigation; sweet cherry; gisela; regina and vegetative.

25

26 **Introduction**

27 Irrigation is one of the major agricultural activities throughout the production season. Its
28 importance increases as climate gets drier (Naor, 2006) [1]. Proper timing of water
29 applications during appropriate periods can increase the yield and quality of most
30 horticultural crops. The inadequate irrigation levels or disproportionate ratios often cause a
31 reduction in yields and a decrease in fruit quality (Koszański et al. 2006) [2]. Recent
32 advances in agro technology makes it possible to apply irrigation to the root zone only

33 thereby making efficient water utilization. This is especially true with drip irrigation system
34 where only a portion of ground is irrigated. The research intends to review the critical growth
35 stages of water requirement in Sweet Cherry under High Density Plantation that play an
36 important role in determining the overall fruit quality. It is pertinent to mention here that in
37 Kashmir Sweet cherry is mainly grown on Karewas which are totally rain fed with poor water
38 holding capacity. Very meagre amount of rainfall (~700 mm) coupled with its erratic
39 distribution results in deficient water supply at flowering and fruit development stage which
40 causes severe pollination problems, poor fruit set, low productivity and inferior fruit quality
41 that ultimately gets reflected by striking drop in economic well being of farmers and also
42 sweet cherry can't withstand the water stagnation conditions, moreover there is an increasing
43 shift from traditional plantation to High Density Plantation for which water management is
44 most essential.

45 Cherries need irrigation or adequate soil moisture to ensure good fruit fill. The most critical
46 stages of irrigation in fruit development of cherry is fruit growth stage followed by fruit set
47 and fruit bud differentiation stage. During fruit growth stage, rapid growth takes place
48 through cell expansion that is dependent upon available water. Uneven precipitation can
49 cause plant stress during critical growth periods, which will affect both crop
50 productivity and quality produce. Dehghanisani *et al.*, (2007) [3] reported that there was a
51 high correlation between the length of young branches and canopy volume on one hand and
52 annual amount of irrigation water applied on the other hand in mature cherry trees. The aim
53 of the present study has been to determine the crop evapotranspiration and effect of different
54 irrigation levels applied at critical growth stages on vegetative growth of sweet cherry under
55 high density plantation through drip irrigation.

56 **Materials and methods**

57 **Experimental site:** This study was carried out in the High Density Experimental Sweet
58 cherry orchard at SKUAST-K, Srinagar (North Kashmir, Jammu and Kashmir), located at an
59 altitude 1588m above sea level on latitude $34^{\circ} 5'$ N and longitude $74^{\circ} 47'$ E during the year
60 2016-17 and 2017-18 (Fig 1). The local climate is temperate with hot and dry, summers and
61 winters cold with snow for almost three months. Meteorological data for the experimental
62 year was measured on a daily basis at the SKUAST-K Agro meteorological Station (Fig 2 &
63 Fig 3). Month wise crop water requirement for cherry at 100% ETc is given in Table 1 which
64 was obtained from Pan Evaporimeter located at the experimental area installed at Agro-
65 metrological station of SKUAST-K Shalimar.

66 **Experimental design and irrigation treatments:** The study material consisted of Sweet
67 cherry plants cv Regina (*Prunus cerasus* x *Prunus canescens*,) on Gisela-5 dwarf rootstock.
68 The trees were planted in 2013 spaced 4 x 2 m apart (Fig 1). The irrigation was applied at
69 four fruit development stages viz T₁ (fruit set), T₂ (pit hardening), T₃ (fruit development) and
70 T₄ (fruit bud differentiation) stages and the amount of irrigation was programmed at I₀ (0%),
71 I₁(50%), I₂ (75%) and I₃ (100%) levels of recorded evapotranspiration as per Class A Pan
72 evaporation. I₀ was used as the control treatment and in this treatment no irrigation was
73 applied at any stage. The amount of irrigation water to be applied on daily basis was
74 calculated from the daily pan evaporation values (Epan) measured in the Class A Pan.



75

76 **Fig 1: Experimental High density Sweet cherry (cv. Regina on Gisela 5)orchard at SKUAST-Kashmir.**



77

78

79

Fig 2: A view of Agrometeorological station of SKUAST-Kashmir



80
81
82
83

Fig 3: USWB CLASS(A) Pan Evaporimeter installed at Agromet station of SKUAST-Kashmir

84 **Measurements:** The quantity of irrigation was estimated by the following FAO
85 methodology:

86 **$ET_c \text{ or } WR = DE \times KC \times AA \times PC \times AC \div IE$**

87 Where, ET_c = Crop Evapotranspiration, WR = Water requirement, DE Daily pan evaporation
88 data, KC = Crop coefficient, PC = Pan Coefficient, AA = Area allotted per plant (m^2), AC =
89 Area shaded by canopy at noon (%), IE = Irrigation efficiency of system (taken as 90%)

90 **Irrigation amount/Water to be applied (litres/tree/day) = ET_c – Effective rainfall** (Value
91 of KC was taken as given by FAO for cherry during various growth stages, PC was taken as
92 0.8, however AC was calculated on daily basis).

93 **Vegetative parameters:** In order to determine the effects of different irrigation levels at
94 various phenological stages on vegetative growth, the following measurements were taken :

95 **Plant height :** Height of each experimental plant was measured with the help of a measuring
96 tape from ground level to the tip of the main leader tape before the commencement and at the
97 end of growing season. The increase in height was calculated by subtracting the initial height
98 from the final height and was expressed in centimetres.

99 **Plant girth :** The plant girth of each experimental plant at 30 cm above the graft union was
100 measured with the help of a measuring tape before the commencement and at the end of

101 growing season. The increase in girth was calculated by subtracting the initial girth from the
 102 final girth and was expressed in centimetres.

103 **Trunk cross sectional area (TCSA) :** The Trunk cross sectional area of each experimental
 104 plant was taken from the measurements of plant girth and expressed in cm² by using the
 105 formula as given by Kumar *et al.*, 2008 [4]:

106
$$\text{TCSA} = \text{Girth}^2 \div 4\pi$$

107 **Shoot extension growth :** The shoot extension growth of each experimental plant was
 108 obtained by measuring the distance between successive terminal bud scars of the same branch
 109 at the end of the growing season and was expressed in cm.

110 **Results and Discussion**

111 **Results**

112 **Applied irrigation water and evapotranspiration:** Sweet cherry trees were irrigated from
 113 15th of april till 31th august, but the first irrigation was applied during july-august 2016 to
 114 record the effect of irrigation applied at fruit bud differentiation stage during the next year.
 115 Water requirement (lt/tree/day) for Sweet cherry during 2017 and 2018 is given in Table 1.
 116 Highest monthly ETc values for treatment I₃ was estimated as 21.45 lt/tree/day in July and
 117 25.56 lt/tree/day in june during 2017 and 2018 respectively (Table 1). The amounts of applied
 118 water per tree in litres was highest in the month of july (624.63) during 2017 and june
 119 (729.84) during 2018 (Table 1).

120 **Table 1: Month-wise crop water requirement for cherry during the growing season at 100 per**
 121 **cent ETc by pan evaporation method during 2017 and 2018**

Month	2017			2018		
	ETc or Water requirement (lt/tree/day)	Water applied (I ₃) (lt/tree)	Total Rainfall l (mm)	ETc or Water requirement (lt/tree/day)	Water applied (I ₃) (lt/tree)	Total Rainfall (mm)
Apr (15-30)	14.12	156.6	92	12.41	137.55	81
May	16.91	500.81	69	14.40	415.92	50.8
Jun	19.21	503.16	121.9	25.56	729.84	61.6
Jul	21.45	624.63	67.2	22.21	607.99	134.2
Aug	20.72	598.8	38	21.25	565.14	120.6

122 *Based on 90% irrigation efficiency of drip irrigation method
 123 The pan co-efficient (Kp) for experimental farm was taken as 0.8

124
 125 **Irrigation and Vegetative growth relations:** The differences between irrigation levels (I) as
 126 well as the phenological stages (T) were statistically significant (p<0.05) for plant height
 127 (Table 2), plant girth (Table 3), TCSA (Table 4) and shoot extension growth (Table 5). Also
 128 the interaction effect of irrigation levels with that of the phenological stages (IxT) were

129 statistically significant for plant girth (Table 3), however the interaction effect (IxT) for plant
 130 height (Table 2), TCSA (Table 4) and shoot extension growth (Table 5) were non significant.

131 Plant height of sweet cherry increased significantly with increase in irrigation levels
 132 (Table 2). Maximum plant height of 19.16 cm was recorded with highest irrigation level of
 133 100 per cent ET_C (I₃), it was followed by 17.29 cm recorded with 75 per cent ET_C level of
 134 irrigation (I₂) however the lowest plant height of 12.69 cm was recorded with 0 per cent ET_C
 135 level of irrigation (I₀). Similarly highest plant height increment of 19.45 cm was found at T₁
 136 stage, whereas the lowest (13.33 cm) incremental plant height was recorded at T₄ stage
 137 (Table 2).

138 **Table 2: Effect of different irrigation levels at various phenological stages on incremental plant**
 139 **height (cm) of Sweet cherry (cv. Regina)**

Phenological Stages / Levels	T ₁	T ₂	T ₃	T ₄	Mean
I ₀	16.09	13.32	11.74	9.60	12.69d
I ₁	19.16	14.84	14.65	13.19	15.44c
I ₂	20.51	17.74	16.37	14.53	17.29b
I ₃	22.11	20.11	18.44	15.99	19.16a
Mean	19.45a	16.50b	15.30c	13.33d	

140 C.D. (p ≤ 0.05) Irrigation levels (I) : 0.624 ; Stages (T) : 0.624 ; I x T : NS

141 T₁ = Fruit set stage (15April-5May); T₂ = Pit hardening stage (6 May-25May); T₃ = Fruit growth stage (26May-8 june); T₄ = Fruit bud
 142 differentiation stage (july-August);

143 I₀ = 0 % ET_C; I₁ = 50 % ET_C; I₂ = 75 % ET_C; I₃ = 100 % ET_C

144 *values followed by the same letter are not significantly different at p< 0.05.

145
 146
 147 The data pertaining to plant girth increment (Table-3) revealed that plant girth of
 148 sweet cherry increased significantly with irrigation. The highest plant girth (3.22 cm) was
 149 recorded with 100 per cent ET_C (I₃) level of irrigation over plant girth of 2.54 cm recorded in
 150 control trees (I₀). Also the highest plant girth of 3.11 cm was recorded at T₃ phenological
 151 stage followed by 3.05 cm at T₄ stage which was statistically at par with T₃ stage (Table 3).
 152 The combination I₃T₃ recorded the highest plant girth of 3.43 cm which was statistically at
 153 par with I₃T₄ and I₂T₃ (Table 3).

154 **Table 3 : Effect of different irrigation levels at various phenological stages on increase in plant**
 155 **girth (cm) of Sweet cherry cv. Regina**

Phenological Stages / Levels	T ₁	T ₂	T ₃	T ₄	Mean

I₀	2.47	2.27	2.73	2.69	2.54d
I₁	2.52	2.40	3.01	2.98	2.74c
I₂	2.7	2.65	3.27a	3.19	2.59b
I₃	3.08	2.98	3.43a	3.38a	3.22a
Mean	2.69b	2.59c	3.11a	3.05a	

156 C.D. ($p \leq 0.05$) Irrigation levels (I) : 0.105 ; Stages (T) : 0.105 ; I x T : 0.210

157 T₁ = Fruit set stage (15April-5May); T₂ = Pit hardening stage (6 May-25May); T₃ = Fruit growth stage (26May-8 june); T₄ = Fruit bud
158 differentiation stage (july-August);

159 I₀ = 0 % ETc; I₁ = 50 % ETc; I₂ = 75 % ETc; I₃ = 100 % ETc

160 *values followed by the same letter are not significantly different at $p < 0.05$.

161 Significant increase in TCSA was noticed during the research with increased
162 irrigation levels, applied at various phenological stages of growth and development (Table 4).
163 Maximum TCSA of 11.92 cm² was recorded with highest irrigation level of 100 per cent ETc
164 (I₃) while as the lowest TCSA of 10.66 cm² was recorded with 0 per cent ETc level of
165 irrigation (control). Similarly, the maximum TCSA of 11.69 cm² was recorded at T₄
166 phenological stage which was statistically at par with with T₃ stage. However, T₂ stage of
167 growth and development recorded significantly lower TCSA of 10.81 cm² (Table 4).

168 **Table 4: Effect of different irrigation levels at various phenological stages on TCSA (cm²) of**
169 **Sweet cherry cv. Regina.**

Phenological Stages \ Levels	T ₁	T ₂	T ₃	T ₄	Mean
I₀	10.29	10.22	11.09	11.03	10.66d
I₁	10.65	10.50	11.62	11.46	11.08c
I₂	10.99	10.90	12.13	11.94	11.49b
I₃	11.73	11.54	12.07	12.33	11.92a
Mean	10.92b	10.81c	11.73a	11.69a	

170 C.D. ($p \leq 0.05$) Irrigation levels (I) : 0.202 ; Stages (T) : 0.202; I x T : NS

171 T₁ = Fruit set stage (15April-5May); T₂ = Pit hardening stage (6 May-25May); T₃ = Fruit growth stage (26May-8 june); T₄ = Fruit bud
172 differentiation stage (july-August);

173 I₀ = 0 % ETc; I₁ = 50 % ETc; I₂ = 75 % ETc; I₃ = 100 % ETc

174 *values followed by the same letter are not significantly different at $p < 0.05$.

175 Annual shoot extension growth of sweet cherry was positively influenced with
176 different levels of irrigation, applied at various phenological stages (Table 5). The maximum
177 annual shoot extension growth of sweet cherry (50.08 cm) was recorded with 100 per cent
178 ETc (I₃) level of irrigation which was found to be statistically at par with 75 (I₂) and 50 (I₁)
179 per cent ETc level of irrigation whereas 0 per cent ETc (I₀) level of irrigation registered
180 lowest (36.04) shoot extension growth. Similarly the annual shoot extension growth of sweet
181 cherry showed a significant difference at various phenological stages of growth and

182 development. Maximum annual shoot extension growth of 47.95 cm was recorded for T₁
 183 stage which was statistically at par with T₂ stage (Table 5).

184 **Table- 5: Effect of different irrigation levels at various phenological stages on Shoot extension**
 185 **growth (cm) of Sweet cherry cv. Regina**

Phenological Stages / Levels	T ₁	T ₂	T ₃	T ₄	Mean
I ₀	38.00	36.66	35.16	34.33	36.04b
I ₁	50.50	49.00	48.33	47.88	48.91a
I ₂	51.33	49.16	48.50	48.00	49.25a
I ₃	52.00	50.33	49.33	48.66	50.08a
Mean	47.95a	46.29a	45.33b	44.70c	

186 C.D. (p ≤ 0.05) Irrigation levels (I) : 2.05 ; Satges (T) : 2.05; I x T : NS

187 T₁ = Fruit set stage (15April-5May); T₂ = Pit hardening stage (6 May-25May); T₃ = Fruit growth stage (26May-8 June); T₄ = Fruit bud
 188 differentiation stage (July-August);

189 I₀ = 0 % ETc; I₁ = 50 % ETc; I₂ = 75 % ETc; I₃ = 100 % ETc

190 *values followed by the same letter are not significantly different at p< 0.05.

191 Discussion

192 Response of sweet cherry to various irrigation levels with respect to vegetative parameters at
 193 various phenological stages showed a positive effect with an increase in irrigation levels this
 194 may be due to the availability of sufficient moisture for continued growth which probably
 195 lead to a greater development of the overall tree canopy. Applied irrigation water based on
 196 Epan values were in agreement with the results reported by Abrisqueta *et al.*, (2001) [5], but
 197 not in agreement with results of study conducted by Dehghanisani *et al.*, (2007) [3]. This
 198 disagreement could be based on different climate conditions .The differences may also be
 199 attributed to different type and age of fruit trees. Many researchers reported that vegetative
 200 growth significantly increased as the irrigation water applied in different stone fruit trees
 201 (Burak and Senih, 2010 [6]., Fereres & Goldhamer, 1990 [7] and Dehghanisani *et al.*, 2007)
 202 [3]. In this study, the relation between vegetative growth and applied irrigation water for I₃
 203 was in agreement with findings of those researches. Also the maximum growth was recorded
 204 at cell division and expansion stage as the cell expansion is most sensitive to water deficit
 205 during growing season. The amount of water available for cell expansion is therefore an
 206 important factor regulating the vegetative growth of plants. The present findings are also in
 207 agreement with those of Hutmacher *et al.* (1994) [8], who observed reduced trunk growth in
 208 almond due to deficit irrigation. Li (1993) [9] reported that deficit irrigation during fruit
 209 development and post-harvest in peach trees significantly reduced vegetative growth. The
 210 greater trunk girth obtained under present study with availability of water might be due to

211 higher absorption of water and nutrient from soil, better translocation of assimilates and
212 production of hormones from roots and better unloading through phloem.

213

214 **Conclusion**

215 The reason for maximum vegetative growth at higher irrigation levels may be due to
216 adaptation ability of young plants to the root zone and plant characteristics such as shallow
217 root development and dwarf rootstock. Irrigation treatment I₃ and I₂ may be recommended as
218 optimum irrigation treatment for irrigation of Regina on Gisela-5 young sweet cherry trees in
219 the temperate conditions. On the other hand, these irrigation treatments must be re-considered
220 in different conditions and I₃ irrigation level should be verified with yield parameters.

221

222 **Competing interests**

223 Authors have declared that no competing interests exist

224 .

225 **References**

226 1. Naor, A. 2006. Irrigation scheduling and evaluation of tree water status in deciduous orchards.,
227 *In: Horticultural Reviews*, (Ed.): J. Janick. ISBN 0-471-73216-8, **32**: 111-165.

228

229 2. Koszański, Z., Rumas-Rudnicka, E. and Podsiadło, C. 2006. *Wpływ nawadniania kropowego i*
230 *nawożenia mineralnego na jakość owoców truskawki [Effect of sprinkling irrigation and mineral*
231 *fertilization on quality of strawberry fruit]*. *J. Elementol.*, **11**(1): 21-27. (in Polish)

232

233 3. Dehghanisani, H., A. Naseri, H. Anyoji and A.E. Eneji. 2007. Effects of deficit irrigation and
234 fertilizer use on vegetative growth of drip irrigated cherry trees, *J. Plant Nutr.*, **30**: 411-425.

235

236 4. Kumar, D, Pandey, V., Anjaneyulu, K. and Nath, V. 2008. Relationship of trunk cross-sectional
237 area with fruit yield, quality and leaf nutrients status in Allahabad Safeda guava (*Psidium guajava*).
238 *Indian J. Agric. Sci.* **78**: 337-39.

239

240 5. Abrisqueta, J.M., A.J. Ruiz and A. Franco. 2001. Water balance of apricot trees. *Agr. Water*
241 *Manage.*, **50**: 211-227.

242 6. Burak, N., C and Senih, Y. 2010. The effects of different irrigation levels on

243 vegetative growth of young dwarf cherry trees in a sub-humid climate. *Pakistan Journal of Botany.*,
244 **42**(5): 3399-3408.

245

- 246 7. Fereres, F. and D.A. Goldhamer. 1990. Deciduous fruits and nut trees. In: irrigation of Agricultural
247 Crop. (Eds.): B.A. Stewart and D.R. Nielsen. *Amer. Soc. of Agronomy*, Madison, WI, pp 987- 1017.
248
- 249 8. Hutmacher, R.B., Nightingale, H.I., Rolston, D.E., Biggar, J.W., Dale, F., Vail, S.S. and Peters, D.
250 1994. Growth and yield responses of almond (*Prunus amygdalus*) to trickle irrigation. *Irrigation*
251 *Science* **14** : 117-126.
252
- 253 9. Li, S. H., 1993. The Response of Sensitive Periods of Fruit Tree Growth, Yield and Quality to
254 Water Stress and Water Saving Irrigation. *Plant Physiol. Commun.*, 29(1:) 10-16.
255
256
257
258
259
260
261
262

UNDER PEER REVIEW